

Solved Examples

JEE Main/Boards

Example 1: A person sees the point A on the rim at the bottom of a cylindrical vessel when the vessel is empty through the telescope T. When the vessel is completely filled with a liquid of refractive index 1.5, he observes a mark at the center B, of the bottom, without moving the telescope or the vessel. What is the height of the vessel if the diameter of its cross section is 10 cm?


Sol: The vessel is filled with a liquid of refractive index μ when the ray from B reaches the telescope.

$$\therefore \mu \sin i = 1.0 \times \sin r, \quad \dots (i)$$

where $\angle r = \angle ADC$, $\angle i = \angle BDC$

$$\sin i = \frac{R}{\sqrt{R^2 + h^2}}, \quad \sin r = \frac{2R}{\sqrt{h^2 + (2R)^2}}$$

where h is the height of the liquid, and R is the radius of the vessel.



Substituting these values in Eq. (i),

$$\mu \times \frac{R}{\sqrt{R^2 + h^2}} = \frac{2R}{\sqrt{h^2 + (2R)^2}} \Rightarrow \mu = \frac{2\sqrt{R^2 + h^2}}{\sqrt{h^2 + (2R)^2}}$$

$$4R^2(\mu^2 - 1) = h^2(4 - \mu^2) \Rightarrow h = 2R \sqrt{\frac{\mu^2 - 1}{4 - \mu^2}}$$

$$h = 10 \sqrt{\frac{(1.5)^2 - 1}{4 - (1.5)^2}} = 10 \sqrt{\frac{5}{7}} = 8.45 \text{ cm.}$$

Example 2: A parallel beam of light rays is incident on a transparent sphere of radius R and a refractive index

μ in the direction of one of the diameters. At what distance from the center of the sphere, will the rays be focused? Assume that $\mu < 2$.

Sol: For refraction of light ray from surface of sphere, the distance of image is obtained by $\frac{\mu_1}{v} - \frac{\mu_2}{r} = \frac{\mu_1 - \mu_2}{R}$.

The ray refracted at one surface becomes object for the opposite surface.

For the first refraction surface:

$$\mu_1 = 1, \mu_2 = \mu, u = -\infty, v = ?, r = +R$$

$$\therefore \frac{\mu}{v} - \frac{1}{-\infty} = \frac{\mu - 1}{R} \Rightarrow v = \frac{\mu R}{\mu - 1}$$

For refraction at the second surface:

I' serves as an object for the second surface. Now, $\mu_1 = \mu$ (since light travels from sphere to air) and $\mu_2 = 1$,

$$u = BI' = + \left(\frac{\mu R}{\mu - 1} - 2R \right) = \frac{R(2 - \mu)}{\mu - 1}$$


(It is positive because light travels from the left to the right, and the distance is also in the same direction.)
 $v = ?, r = -R$.

$$\therefore \frac{1}{v} - \frac{\mu(\mu - 1)}{R(2 - \mu)} = \frac{1 - \mu}{-R} \text{ or } v = \frac{R(2 - \mu)}{2(\mu - 1)}$$

This distance is positive and is referred from the second surface.

\therefore Distance of the focal point from the center

$$= \frac{R(2 - \mu)}{2(\mu - 1)} + R = \frac{\mu R}{2(\mu - 1)}$$



Example 3: A rectangular glass block of thickness 10 cm and refractive index 1.5 is placed over a small coin. A beaker is filled with water of refractive index $4/3$ to a height of 10 cm and is placed over the block.

(i) Find the apparent position of the object when it is viewed from normal incidence.

(ii) Draw a neat ray diagram.

(iii) If the eye is slowly moved away from the normal, at a certain position, the object is found to disappear due to the TIR. At which surface, does this happen and why?

Sol: The image of coin formed at upper surface of block, becomes object for beaker containing water. The image

thus formed at distance v is given by $\frac{\mu_1}{v} - \frac{\mu_2}{r} = \frac{\mu_1 - \mu_2}{R}$.

For the first surface: $\mu_2 = \frac{4}{3}$, $\mu_1 = 1.5$,

$u = -10\text{ cm}$, $R = \infty$, $v_1 = ?$

$$\frac{\frac{4}{3}}{v_1} - \frac{1.5}{-10} = \frac{\frac{4}{3} - 1.5}{\infty} \Rightarrow v_1 = -\frac{80}{9}\text{ cm}$$

I' serves as an object for the second surface. For the second surface:

Note: This is an alternative to the apparent depth relation.


$$u = -BI' = -(BA + AI') = -\left(10 + \frac{80}{9}\right) = -\frac{170}{9}\text{ cm}$$

$\mu_1 = \frac{4}{3}$, $\mu_2 = 1$, $R = \infty$, $v_2 = ?$

$$\therefore \frac{1}{v_2} - \frac{\frac{4}{3}}{-\frac{170}{9}} = \frac{1 - \frac{4}{3}}{\infty} \Rightarrow v_2 = 14.2\text{ cm}$$

θ_c : critical angle for glass–water interface

$$= \sin^{-1} \frac{1}{\mu_g} \Rightarrow \theta_c = \sin^{-1} \frac{1}{\mu_a \times \mu_g}$$



$$= \sin^{-1} \cdot \frac{1}{\frac{3}{4} \times \frac{3}{2}} = \sin^{-1} \frac{8}{9} = 62.7^\circ$$

The critical angle for water–air interface

$$= \sin^{-1} \frac{1}{\mu_w} = \sin^{-1} \frac{3}{4} = 48^\circ.$$

Obviously, therefore, TIR takes place earlier at the water–air interface.

Example 4: How long will the light take in travelling a distance of 500 m in water? Given that μ for water is $4/3$ and the velocity of light in vacuum is 3×10^{10} cm/s. Also calculate the equivalent path.

Sol: The velocity of light in a medium is given as $v = \frac{c}{\mu}$.

The optical path travelled by light is $\ell = \mu \times d$ where d is the distance travelled in water.

We know that

$$\mu = \frac{\text{Velocity of light in vacuum}}{\text{Velocity of light in water}}$$

$$\frac{4}{3} = \frac{3 \times 10^{10}}{\text{Velocity of light in water}}$$

Velocity of light in water = 2.25×10^{10} cm/s.


$$\text{Time taken} = \frac{500 \times 100}{2.25 \times 10^{10}} = 2.22 \times 10^{-6} \text{ sec.}$$

Equivalent optical path

= $\mu \times$ distance travelled in water

$$= \frac{4}{3} \times 500 = 666.64 \text{ m.}$$

Example 5: In the figure shown, for an angle of incidence i at the top surface, what is the minimum refractive index needed for TIR at the vertical face?



Sol: Total internal reflection of light occurs inside body when the angle of incidence is greater than critical

angle θ_c and according to Snell's law $\theta_c = \sin^{-1} \left(\frac{1}{\mu} \right)$

Applying the Snell's law at the top surface,

$$\mu \sin r = \sin i \quad \dots (i)$$

For TIR, the vertical face

$$\mu \sin \theta_c = 1$$

Using geometry, $\theta_c = 90^\circ - r$

$$\mu \sin(90 - r) = 1 \Rightarrow \mu \cos r = 1 \quad \dots (ii)$$


On squaring and adding Eqs (i) and (ii), we get

$$\begin{aligned} \therefore \mu^2 \sin^2 r + \mu^2 \cos^2 r &= 1 + \sin^2 i \\ \Rightarrow \mu &= \sqrt{1 + \sin^2 i} \end{aligned}$$

Example 6: A point source of light is placed at the bottom of a tank containing a liquid (refractive index = μ) up to a depth h . A bright circular spot is seen on the surface of the liquid. Find the radius of this bright spot.

Sol: The light waves incident on the surface of the water at an incident angle greater than critical angle will get reflected in side water. This light waves forms cone in the volume of the tank. Thus we get relation of critical angle as

$$\tan \theta_c = \frac{R}{h}$$




Rays coming out of the source and incident at an angle greater than θ_c will be reflected back into the liquid; therefore, the corresponding region on the surface will appear dark. As it is obvious from the figure, the radius of the bright spot is given by

$$R = h \tan \theta_c = \frac{h \sin \theta_c}{\cos \theta_c} \Rightarrow R = \frac{h \sin \theta_c}{\sqrt{1 - \sin^2 \theta_c}}$$

$$\text{Since } \sin \theta_c = \frac{1}{\mu}; \quad \therefore R = \frac{h}{\sqrt{\mu^2 - 1}}$$

Example 7: The cross section of the glass prism has the form of an isosceles triangle. One of the equal faces is coated with silver. A ray of light incident normally on the other equal face and after getting reflected twice emerges through the base of prism along the normal. Find the angle of the prism.



Sol: The angles of prism add up to 180° .

From the figure, $\alpha = 90^\circ - A$

$$i = 90^\circ - \alpha; \quad i = 90^\circ - \alpha = A \quad \dots (i)$$

$$\text{Also, } \beta = 90^\circ - 2i = 90^\circ - 2A$$

$$\text{and } \gamma = 90^\circ - \beta = 2A$$

$$\text{Thus, } \gamma = r = 2A$$

From geometry,

$$A + \gamma + \gamma = 180^\circ. \quad \text{or } A = \frac{180}{5} = 36^\circ.$$

Example 8: A lens has a power of +5 dioptre in air. What will be its power if completely immersed in water?

$$\text{Given } \mu_g = \frac{3}{2}; \quad \mu_w = \frac{4}{3}.$$

Sol: According to the lens maker's formula the focal

$$\text{length } 1/f \text{ is } \frac{1}{f} = \left(\frac{\mu_1}{\mu_2} - 1 \right) \left[\frac{1}{R_1} - \frac{1}{R_2} \right]$$

Let f_a and f_w be the focal lengths of the lens in air and water, respectively, then,

$$P_a = \frac{1}{f_a} \quad \text{and} \quad P_w = \frac{\mu_w}{f_w}.$$

$$f_a = 0.2\text{m} = 20\text{cm}.$$

Using the lens maker's formula,

$$P_a = \frac{1}{f_a} = (\mu_g - 1) \left[\frac{1}{R_1} - \frac{1}{R_2} \right] \quad \dots (i)$$

$$\frac{1}{f_w} = \left(\frac{\mu_g}{\mu_w} - 1 \right) \left[\frac{1}{R_1} - \frac{1}{R_2} \right]$$

$$P_w = \frac{1}{f_w} = \left(\frac{\mu_g - \mu_w}{\mu_w} \right) \left[\frac{1}{R_1} - \frac{1}{R_2} \right] \quad \dots(ii)$$

Dividing Eq. (ii) by Eq. (i), we get,

$$\frac{P_w}{P_a} = \frac{(\mu_g - \mu_w)}{\mu_w(\mu_g - 1)} = \frac{1}{3} \cdot \frac{1}{\mu_w}$$

$$\Rightarrow P_w = P_a \left(\frac{1}{3} \right) \left(\frac{3}{4} \right) = \frac{+5}{4} D$$

Example 9: The distance between two point sources of light is 24 cm. Find out where you would place a converging lens of focal length 9 cm, so that the images of both the sources are formed at the same point.


Sol: For lens the distance of the image formed from the lens is given by $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$ where u , v and f are distance of object, distance of image and focal length respectively.

$$\text{For } S_1 : \frac{1}{v_1} - \frac{1}{-x} = \frac{1}{9}$$

$$\therefore \frac{1}{v_1} = \frac{1}{9} - \frac{1}{x} \quad \dots (i)$$

$$\text{For } S_2 : \frac{1}{v_2} - \frac{1}{-(24-x)} = \frac{1}{9} \quad \dots (ii)$$

$$\therefore \frac{1}{v_2} = \frac{1}{9} - \frac{1}{24-x}$$



Since, the sign convention for S_1 and S_2 is just opposite. Hence,

$$v_1 = -v_2$$

$$\Rightarrow \frac{1}{v_1} = -\frac{1}{v_2}$$

$$\therefore \frac{1}{9} - \frac{1}{x} = \frac{1}{24-x} - \frac{1}{9}$$

Solving this equation, we get $x = 6$ cm. Therefore, the lens should be kept at a distance of 6 cm from either of the object.

Example 10: Two equiconvex lenses of focal lengths 30 cm and 70 cm, made of material of refractive index = 1.5, are held in contact coaxially by a rubber band round their edges. A liquid of refractive index 1.3 is introduced in the space between the lenses filling it completely. Find the position of the image of a luminous point object placed on the axis of the combination lens at a distance of 90 cm from it.

Sol: This system is combination of three lenses. Two lenses of glass one lens of liquid. Add the powers to get total power.

$$|R_1| = |R_2| = f_1 = 30 \text{ cm (As } \mu = 1.5).$$

$$\text{Similarly, } |R_3| = |R_4| = f_2 = 70 \text{ cm.}$$

The focal length of the liquid lens (in air),

$$\frac{1}{f_3} = (\mu - 1) \left(\frac{1}{R_2} - \frac{1}{R_3} \right)$$


$$= (1.3 - 1) \left(\frac{1}{-30} - \frac{1}{70} \right) = -\frac{1}{70}$$

$$(b) m = \frac{v}{u}$$

$$\therefore m_1 = \frac{(5.0 - 4.0)}{(-4.0)} = -0.25,$$


$$\text{and } m_2 = \frac{(5.0 - 1.0)}{(-1.0)} = -4.00.$$

Hence, both the images are real and inverted, the first is magnification -0.25 , and the second is -4.00 .



JEE Advanced/Boards

Example 1: A 4-cm-thick layer of water covers a 6-cm-thick glass slab. A coin is placed at the bottom of the slab and is being observed from the air side along the normal to the surface. Find the apparent position of the coin from the surface.



Sol: As the thick layer of water is placed over the glass slab. The coin placed beneath the glass slab will


appear to shift upwards due to both glass and water by distance s . This apparent shift is thus given by

$$s = s_1 + s_2 = \left(1 - \frac{1}{\mu_1}\right) \times h_1 + \left(1 - \frac{1}{\mu_2}\right) \times h_2$$

The total apparent shift is

$$s = h_1 \left(1 - \frac{1}{\mu_1}\right) + h_2 \left(1 - \frac{1}{\mu_2}\right)$$

$$s = 4 \left(1 - \frac{1}{4/3}\right) + 6 \left(1 - \frac{1}{3/2}\right) = 3 \text{ cm.}$$



Thus, $h = h_1 + h_2 - s = 4 + 6 - 3$
 $= 7.0 \text{ cm.}$

Example 2: An achromatic lens of focal length 100 cm is made up of crown and flint glass lenses. Find the focal length of each lens. Given that for crown glass $\mu_v = 1.5245$, $\mu_r = 1.5155$ and for flint glass $\mu'_v = 1.659$ and $\mu'_r = 1.641$.

Sol: The focal length of the combination of lens is given by $\frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2}$.

If μ and μ' are mean refractive indices for crown and flint glasses, respectively, then,

$$\mu = \frac{\mu_v + \mu_r}{2} = \frac{1.5245 + 1.5155}{2} = 1.52.$$

$$\mu' = \frac{\mu'_v + \mu'_r}{2} = \frac{1.659 + 1.641}{2} = 1.65.$$

The dispersive powers ω and ω' for crown and flint glass, respectively, are

$$\omega = \frac{\mu_v - \mu_r}{\mu - 1} = \frac{1.5245 - 1.5155}{1.52 - 1} = \frac{0.009}{0.52} = \frac{9}{520}$$

$$\omega' = \frac{\mu'_v - \mu'_r}{\mu' - 1} = \frac{1.659 - 1.641}{1.65 - 1} = \frac{0.018}{0.65} = \frac{9}{325}$$

To have an achromatic combination,

$$\frac{\text{focal length for crown glass}}{\text{focal length for flint glass}} = \frac{f}{f'} = -\frac{\omega}{\omega'}$$

$$= -\frac{9}{520} \times \frac{325}{9} = -\frac{5}{8}; \quad \frac{1}{f} = -\frac{5}{8f'} \quad \dots (i)$$

As the focal length of the combination is 100 cm,

$$\frac{1}{F} = \frac{1}{f} + \frac{1}{f'} = \frac{1}{f} - \frac{5}{8f} = \frac{3}{8f} = \frac{1}{100}.$$

$$f = \frac{3}{8} \times 100 = 37.5 \text{ cm.}$$

$$f' = \frac{-8}{5} \times f = \frac{-8}{5} \times \frac{75}{2} = -60 \text{ cm.}$$


The achromatic doublet requires a convex lens of focal length 37.5 cm made of crown glass and a concave lens of focal length 60 cm made of flint glass.

Example 3: A 20-cm-thick slab of glass of refractive index 1.5 is placed in front of a plane mirror and a pin is placed in front of it in the air at a distance of 40 cm from the mirror. Find the position of the image.

Sol: The slab of thickness t forms the image of the object O at the point O' .

The slab shifts the image of object by $(t - t/\mu)$. O' serves as an object for the plane mirror.

Object distance for the mirror is MO .



$$= 40 - \frac{20}{3} = \frac{100}{3} \text{ cm.}$$

Since in a plane mirror, object distance = image distance.

$$MI' = MO' = \frac{100}{3} \text{ cm.}$$


Now, I' serves as an object for the slab again.

$$I' \text{ is shifted to } I'' \text{ by } 20 - \frac{20}{1.5} = \frac{20}{3} \text{ cm.}$$

$$\therefore \text{Distance of the final image } (I'') \text{ from the mirror}$$

$$= \frac{100}{3} - \frac{20}{3} = \frac{80}{3} \text{ cm.}$$

Example 4: The convex surface of a thin concavo-convex lens of glass of refractive index 1.5 has a radius of curvature 20 cm. The concave surface has a radius of curvature of 60 cm. The convex side is coated with silver and placed at a horizontal surface as shown in the figure.



(a) Where a pin should be placed on the optical axis such that its image is formed at the same place?

(b) If the concave part is filled with water of refractive index $4/3$, find the distance through which the pin should be moved, so that the image of the pin again coincides with pin.

Sol: When the convex side of the concavo-convex lens is coated with silver, the combination becomes a mirror. This combination consist two lenses and one mirror placed close to each other. The powers of all the three will be added. When the water is filled on concave side, we get plano-convex water lens whose focal length is found by lens makers formula. This combination consists of four lenses (two lenses of glass and two lenses of water) and one mirror.

(a) The refraction takes place from the first surface, reflection from the lower surface and finally refraction from the first surface of focal lengths f_g , f_m and f_g , respectively. The combined focal length F is given by

$$\frac{1}{F} = \frac{1}{f_g} + \frac{1}{f_m} + \frac{1}{f_g} = \frac{2}{f_g} + \frac{1}{f_m} \quad \dots (i)$$

$$f_m = R_2 / 2 = 20 / 2 = 10 \text{ cm.}$$

The value of f_g can be obtained by using the formula

$$\frac{1}{f_g} = ({}^a\mu_g - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$= (1.5 - 1) \left(\frac{1}{20} - \frac{1}{60} \right)$$

$$f_g = 60 \text{ cm.}$$

Substituting these values in Eq. (i),

$$\frac{1}{F} = \frac{2}{60} + \frac{1}{10} = \frac{2}{15}$$

$$F = \frac{15}{2} = 7.5 \text{ cm.}$$

For the image to be informed at the same point as the object

$$u = 2F = 2 \times 7.5 = 15 \text{ cm.}$$

The object should be placed at a distance of 15 cm from the lens on the optical axis.

(b) If f_w is the focal length of lens in water, the focal length F' of this combination is given by

$$\frac{1}{F'} = \frac{1}{f_w} + \frac{1}{f_g} + \frac{1}{f_m} + \frac{1}{f_g} + \frac{1}{f_w}$$

$$\frac{1}{F'} = \frac{2}{f_w} + \frac{2}{f_g} + \frac{1}{f_m} \quad \dots (ii)$$

$$f_g = 60 \text{ cm, } f_m = 10 \text{ cm.}$$

The value of f_w is calculated by using the relation,

$$\frac{1}{f_w} = ({}^a\mu_w - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$\Rightarrow \frac{1}{f_w} = \left(\frac{4}{3} - 1 \right) \left(\frac{1}{60} \right) = \frac{1}{180}$$

$$f_w = 180 \text{ cm.}$$

Substituting these values in Eq. (ii), we get

$$\frac{1}{F'} = \frac{2}{180} + \frac{2}{60} + \frac{1}{10}; F' = \frac{90}{13}$$

$$u' = 2F' = \frac{2 \times 90}{13} = \frac{180}{13} \text{ cm.}$$

Displacement of the pin

$$= u - u' = 15 - \frac{180}{13} = \frac{15}{13} = 1.14 \text{ cm.}$$

Example 5: The radius of curvature of the convex face of plano-convex lens is 12 cm, and its $\mu = 1.5$.

(a) Find the focal length of the lens.

The plane surface of the lens is coated with silver.

(b) At what distance from the lens, will the parallel rays incident on the convex surface converge?

(c) Sketch the ray diagram to locate the image, when a point object is placed on the axis at a distance of 20 cm from the lens.

(d) Calculate the image distance when the object is placed as in (c).

Sol: Use the lens maker's formula to find the focal length of the plano-convex lens. When the plane side of the lens is coated with silver, the combination becomes

a mirror. This combination consist two lenses and one mirror placed close to each other. The powers of all the three will be added.

(a) The focal length f of lens of refractive index μ is given by the formula.

$$\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right).$$

In case of plano-convex lens,

$$R_1 = \infty, R_2 = -12 \text{ cm}, \mu = 1.5$$

$$\therefore \frac{1}{f} = (1.5 - 1) \left(\frac{1}{12} \right)$$


$$f = 24 \text{ cm}.$$

(b) The focal length, f_m , of plane-silvered surface is infinity. The focal length F of the plano-convex lens, when the plane surface is coated with silver is given by


$$\frac{1}{F} = \frac{1}{f} + \frac{1}{f_m} + \frac{1}{f} = \frac{2}{f} + \frac{1}{\infty}$$

$$\therefore F = \frac{f}{2} = \frac{24}{2} = 12 \text{ cm}.$$

Such a lens behaves like a concave mirror of focal length 12 cm. The parallel rays converge at a distance of 12 cm from the silvered surface.



(c) The figure shows the ray diagram of the image formed by this lens when the object is placed at a distance of 20 cm from the lens. The light is incident from the right to the left.



(d) For a lens,

$$\frac{1}{v} + \frac{1}{u} = \frac{1}{F}.$$

For $F = 12 \text{ cm}$, $u = +20 \text{ cm}$ and $v = ?$

$$\frac{1}{v} = \frac{1}{12} - \frac{1}{20} = \frac{1}{30}$$

$$v = 30 \text{ cm}.$$

A real image is formed.

Example 6: An object O is placed at a distance of 15 cm from a convex lens A of focal length 10 cm and its image I_1 is formed on a screen on the other side of the lens. A concave lens B is now placed midway between A and I_1 , then the screen is moved back 10 cm to receive a clear image I_2 .

Find (a) The focal length of the concave lens and

(b) Linear magnification of the final image.


Sol: The convex lens forms the image of the object at point I_1 . This image acts as the object for concave lens and the final image is formed at I_1' .

For refraction through convex lens,

$$u_1 = OP = -15 \text{ cm}.$$

$$f = +10 \text{ cm}, v_1 = PI_1.$$

$$\text{Using } \frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$



$$\frac{1}{v_1} = \frac{1}{f} + \frac{1}{u} = \frac{1}{10} + \left(\frac{1}{-15} \right) = \frac{3-2}{30} = \frac{1}{30}$$

$$\Rightarrow v_1 = 30 \text{ cm}.$$

I_1 serves as a virtual object for lens B . For refraction through lens B ,

$$u_2 = QI_1 = PI_1 - 15 = 30 - 15 = +15 \text{ cm},$$

$$v_2 = 15 + 10 = +25 \text{ cm}.$$

$$\frac{1}{f} = \frac{1}{v} - \frac{1}{u} = \frac{1}{25} - \frac{1}{15} = \frac{3-5}{75} = -\frac{2}{75}$$

$$f = -\frac{75}{2} = -37.5 \text{ cm},$$

The negative sign implies it is a concave lens.


$$\begin{aligned} \text{Linear magnification} &= \frac{v_1}{u_1} \times \frac{v_2}{u_2} \\ &= \frac{30}{15} \times \frac{5}{3} = -\frac{10}{3} = -3.33 \end{aligned}$$

The negative sign shows that the final image is inverted.

Example 7: A parallel beam of light travelling in water ($\mu = 4/3$) is refracted by a spherical air bubble of radius 2 mm placed in water. Assume that the light rays to be paraxial.

(a) Find the position of the image due to refraction at the first surface and the position of the final image.

(b) Draw a ray diagram showing the positions of both images.



Sol: The image of the object formed by the first refraction by the water-glass surface acts as the object for the second refraction at glass-water surface.

(a) For the refraction from a single spherical surface, we have

$$\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{(\mu_2 - \mu_1)}{R}$$

Let P_1 be pole at the first surface and P_2 to be pole at the second surface. At P_1 ,

$$\mu_1 = (4/3); \quad \mu_2 = 1; \quad R = +0.2\text{cm}, \quad u_1 = \infty.$$

$$\text{So } \frac{1}{v_1} - \frac{(4/3)}{\infty} = \frac{1 - (4/3)}{+0.2}.$$

$$\therefore v_1 = -0.6\text{cm}.$$

The first surface will form a virtual image I_1 at a distance 0.6 cm to the left of P_1 as shown in the figure.

This image acts as an object for the second surface. So for the second surface at P_2 ,

$$\mu_1 = 1, \quad \mu_2 = (4/3), \quad R = -0.2\text{cm}$$

$$\text{and } \mu_2 = -(0.6 + 0.4) = -1.0\text{cm}$$


$$\frac{(4/3)}{v_2} - \frac{1}{(-1)} = \frac{(4/3) - 1}{-0.2}$$

$$v_2 = -0.5\text{cm}$$

The final image I_2 is formed at a distance of 0.5 cm to the left of the second surface P_2 . The final image is at a distance of $0.5 - 0.4 = 0.1\text{cm}$ to the left of the first surface as shown in figure.

(b) The ray diagram is already shown in the figure drawn.

Example 8: An object is placed at a distance of 12 cm to the left of a diverging lens of focal length -6.0 cm. A converging lens with a focal length of 12.0 cm is placed at a distance d to the right of the diverging lens. Find the distance d that corresponds to a final image at infinity.



Sol: The concave lens forms the image of the object at point say I_1 . This image acts as the object for convex lens and the final image is formed at say I_1' .

Applying lens formula $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$ twice, we have

$$\frac{1}{v_1} - \frac{1}{-12} = \frac{1}{-6} \quad \dots (i)$$

$$\frac{1}{\infty} - \frac{1}{v_1 - d} = \frac{1}{12} \quad \dots (ii)$$

Solving Eqs (i) and (ii), we have


$$v_1 = -4\text{cm}.$$

$$\text{And } d = 8\text{cm}.$$

Example 9: A solid glass sphere with a radius R and a refractive index of 1.5 is coated with silver over a hemisphere. A small object is located on the axis of the sphere at a distance $2R$ to the left of the vertex of the un-silvered hemisphere. Find the position of the final image after all refractions and reflections have taken place.

Sol: The image formed by first refraction at air-glass spherical surface acts as the object for the concave mirror. The image formed by the mirror acts as the object for spherical glass-air spherical surface.

The ray of light first gets refracted, then reflected and then again refracted. For the first refraction and then reflection, the ray of light travels from the left to the right, while for the last refraction it travels from the right to the left. Hence, the sign convention will change accordingly.



First refraction: Using $\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R}$ with proper sign conventions, we have


$$\frac{1.5}{v_1} - \frac{1.0}{-2R} = \frac{1.5 - 1.0}{+R} \quad \therefore v_1 = \infty.$$

Second reflection: Using $\frac{1}{v} + \frac{1}{u} = \frac{1}{f} = \frac{2}{R}$ with proper sign conventions, we have

$$\frac{1}{v_2} + \frac{1}{\infty} = -\frac{2}{R} \quad \therefore v_2 = -\frac{R}{2}$$

Third refraction: Again using $\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R}$

with reversed sign convention, we have



$$\frac{1.0}{v_3} - \frac{1.5}{-1.5R} = \frac{1.0 - 1.5}{-R}$$

$$\Rightarrow v_3 = -2R;$$

i.e. final image is formed on the vertex of the silvered face.

JEE Main/Boards

Exercise 1

Q.1 A ray of light incident on an equilateral glass prism shows a minimum deviation of 30° . Calculate the speed of light through the glass prism.

Q.2 Where an object should be placed from a converging lens of focal length 20 cm so as to obtain a real image of magnification 2?

Q.3 What changes in the focal length of (i) a concave mirror and (ii) a convex lens occur, when the incident violet light on them is replaced with red light?

Q.4 State the conditions for TIR of light to take place. Calculate the speed of light in a medium, whose critical angle is 45° .

Q.5 An object is placed at the focus of concave lens. Where will its image be formed?

Q.6 Draw a graph to show the variation of the angle of deviation ' D ' with that of the angle of incidence ' i ' for a monochromatic ray of light passing through a glass prism refracting angle ' A '. Hence, deduce the relation.

Q.7 Draw a ray diagram of an astronomical telescope in the normal adjustment position. Write down the expression for its magnifying power.

Q.8 A spherical surface of radius of curvature R separates a rarer and a denser medium. Complete the path of the incident ray of light, showing the formation of the real image. Hence, derive the relation connecting an object distance ' u ', image distance ' v ', radius of curvature R and the refractive indices n_1 and n_2 of the two media. Briefly explain how the focal length of a convex lens changes, with an increase in wavelength of incident light.

Q.9 (a) Draw a labelled ray diagram to show the formation of the image by a compound microscope. Write the expression for its magnifying power.

(b) How does the resolving power of a compound microscope change and when?

(i) Refractive index of the medium between the object and the objective lens increases and

(ii) Wavelength of the radiation used is increased.

Q.10 Draw a labeled ray diagram to show the image formation I of a refractive-type astronomical telescope. Why should the diameter of the objective of a telescope be large?

Q.11 A beam of light converges to a point P. A lens is placed in the path of the convergent beam at a distance of 12 cm from P. At what point, does the beam converge if the lens is

(i) A convex lens of focal length 20 cm.

(ii) A concave lens of focal length 16cm?

Do the required calculations.

Q.12 A double convex lens of glass of refractive index 1.6 has its both surfaces of equal radii of curvature of 30 cm each. An object of height 5 cm is placed at a distance of 12.5 cm from the lens. Calculate the size of the image found.

Q.13 Why does the bluish color predominate in a clear sky?

Q.14 How does the angle of minimum deviation of a glass prism of a refractive index 1.5 change, if it is immersed in a liquid of refractive index 1.3?

Q.15 Draw a labeled ray diagram, showing the image formation of an astronomical telescope in the normal adjustment position. Write the expression for its magnifying power.

Q.16 Derive the lens formula, $\frac{1}{f} = \frac{1}{v} - \frac{1}{u}$ for a concave lens, using the necessary ray diagram.

Two lenses of power 10 D and -5 D are placed in contact.

(i) Calculate the power of the new lens.

(ii) Where should an object be held from the lens so as to obtain a virtual image of magnification 2?

Q.17 Two thin lenses of power $+6$ D and -2 D are in contact. What is the focal length of the combination?

Q.18 Define the refractive index of a transparent medium. A ray of light passes through a triangular prism. Plot a graph showing the variation of the angle of deviation with an angle of incidence.

Q.19 (a) (i) Draw a labeled ray diagram to show the formation of image in an astronomical telescope for a distant object.

(ii) Write three distinct advantages of a reflecting-type telescope over a refracting-type telescope.

(b) A convex lens of focal length 10 cm is placed coaxially 5 cm away from a concave lens of focal length 10 cm. If an object is placed 30 cm in front of the convex lens, find the position of the final image formed by the combined system.

Q.20 A converging lens is kept coaxially in contact with a diverging lens – both the lenses being of equal focal lengths. What is the focal length of the combination?

Q.21 (a) (i) Draw a neat labeled ray diagram of an astronomical telescope in the normal adjustment. Explain briefly its working.

(ii) An astronomical telescope uses two lenses of power 10 D and 1 D. What is its magnifying power in the normal adjustment?

(b) (i) Draw a neat labeled ray diagram of a compound microscope. Explain briefly its working.

(ii) Why must both the objective and the eyepiece of a compound microscope have short focal lengths?

Q.22 Draw a labeled ray diagram of a reflecting telescope. Mention its two advantages over the refracting telescope.

Q.23 (i) An object is placed between two plane mirrors inclined at 60° to each other. How many images do you expect to see?

(ii) An object is placed between two plane parallel mirrors. Why do the distant images get fainter and fainter?

Q.24 The magnifying power of a compound microscope is 20. The focal length of its eye piece is 3 cm. Calculate the magnification produced by the objective lens.

Q.25 An astronomical telescope having a magnifying power of 8 consists of two thin lenses 45 cm apart. Find the focal length of the lenses.

Exercise 2

Single Correct Choice Type

Q.1 Two plane mirrors are inclined at 70° . A ray incident on one mirror at an angle of θ after reflection falls on the second mirror and is reflected from there parallel to the first mirror, θ is:

- (A) 50° (B) 45°
(C) 30° (D) 55°

Q.2 There are two plane mirror with reflecting surfaces facing each other. Both the mirrors are moving with the speed of v away from each other. A point object is placed between the mirrors. The velocity of the image formed due to the n^{th} reflection will be

- (A) nv (B) $2nv$
(C) $3nv$ (D) $4nv$

Q.3 A man of height ' h ' is walking away from a street lamp with a constant speed ' v '. The height of the street lamp is $3h$. The rate at which the length of the man's shadow is increasing when he is at a distance of $10h$ from the base of the street lamp is:

- (A) $v/2$ (B) $v/3$
(C) $2v$ (D) $v/6$


Q.4 A boy of height 1.5 m with his eye level at 1.4 m stands before a plane mirror of length 0.75 m fixed on the wall. The height of the lower edge of the mirror above the floor is 0.8 m. Then,

- (A) The boy will see his full image.
(B) The boy cannot see his hair.

(C) The boy cannot see his feet.

(D) The boy cannot see neither his hair nor his feet.


Q.5 A point source of light S is placed in front of two large mirrors as shown. Which of the following observers will see only one image of S ?



- (A) Only A (B) Only C
(C) Both A and C (D) Both B and C

Q.6 In the figure shown if the object 'O' moves toward the plane mirror, then the image I (which is formed after successive reflections from M_1 & M_2 , respectively) will move:

- (A) Toward right
(B) Toward left
(C) With zero velocity
(D) Cannot be determined



Q.7 A point source of light is 60 cm away from a screen and is kept at the focus of a concave mirror that reflects light on the screen. The focal length of the mirror is 20 cm. The ratio of average intensities of the illumination on the screen when the mirror is present and when the mirror is removed is:


- (A) 36:1 (B) 37:1
(C) 49:1 (D) 10:1

Q.8 A concave mirror is placed on a horizontal table, with its axis directed vertically upward. Let O be the pole of the mirror and C its center of curvature. A point object is placed at C . It has a real image, also located at C (a condition called auto-collimation). If the mirror is now filled with water, the image will be:

- (A) Real and will remain at C
(B) Real and located at a point between C and ∞
(C) Virtual and located at a point between C and O
(D) Real and located at a point between C and O


Q.9 In the diagram shown below, a point source O is placed vertically below the center of a circular plane

mirror. The light rays starting from the source are reflected from the mirror such that a circular area A on the ground receives light. Now, a glass slab is placed between the mirror and the source O. What will be the magnitude of the new area on the ground receiving light?



- (A) A
 (B) Greater than A
 (C) Less than A
 (D) Cannot say, as the information given is insufficient

Q.10 In the figure ABC is the cross section of a right-angled prism, and BCDE is the cross section of a glass slab. The value of θ , so that light incident normally on the face AB does not cross the face BC, is (given $\sin^{-1}(3/5) = 37^\circ$)



- (A) $\theta \leq 37^\circ$ (B) $\theta > 37^\circ$
 (C) $\theta \leq 53^\circ$ (D) $\theta < 53^\circ$

Q.11 A small source of light is 4 m below the surface of a liquid of refractive index $5/3$. In order to cut off all the light coming out of liquid surface, the minimum diameter of the disc placed on the surface of liquid is:

- (A) 3m (B) 4m (C) 6m (D) ∞


Q.12 A point source of light is placed at a distance h below the surface of a large deep lake. What is the percentage of light energy that escapes directly from the water surface? μ of the water = $4/3$? (Neglect partial reflection)

- (A) 50% (B) 25% (C) 20% (D) 17%

Q.13 When the object is at distances u_1 and u_2 , the images formed by the same lens are real and virtual, respectively, and of same size. Then, the focal length of the lens is:


- (A) $\frac{1}{2}\sqrt{u_1 u_2}$ (B) $\frac{1}{2}(u_1 + u_2)$
 (C) $\sqrt{u_1 u_2}$ (D) $2(u_1 + u_2)$

Q.14 Parallel beam of light is incident on a system of two convex lenses of focal lengths $f_1 = 20$ cm and $f_2 = 10$ cm. What should be the distance between the two lenses so that rays after refraction from both lenses get un-deviated?




- (A) 60 cm (B) 30 cm (C) 90 cm (D) 40 cm

Q.15 An object is placed at a distance of 15 cm from a convex lens of a focal length 10 cm. On the other side of the lens, a convex mirror is placed at its focus such that the image formed by the combination coincides with the object itself. The focal length of the convex mirror is



- (A) 20 cm (B) 10 cm (C) 15 cm (D) 30 cm


Q.16 Look at the ray diagram shown, what will be the focal length of the 1st and the 2nd lens, if the incident light ray is parallel to emergent ray.




- (A) -5cm and -10cm (B) $+5\text{cm}$ and $+10\text{cm}$
 (C) -5cm and $+5\text{cm}$ (D) $+5\text{cm}$ and $+5\text{cm}$

Q.17 A point object is kept at the first focus of a convex lens. If the lens starts moving toward right with a constant velocity, the image will

- (A) Always move toward right
 (B) Always move toward left
 (C) First move toward right and then toward left.
 (D) First move toward left and then toward right.



Q.18 A ray incident at an angle 53° on a prism emerges at an angle of 37° as shown. If the angle of incidence is 50° , which of the following is a possible value of the angle of emergence



- (A) 35° (B) 42° (C) 40° (D) 38°

Q.19 A prism has a refractive index $\sqrt{\frac{3}{2}}$ and a refracting angle 90° . Find the minimum deviation produced by the prism.

- (A) 40° (B) 45° (C) 30° (D) 49°

Q.20 A certain prism is found to produce a minimum deviation of 38° . It produces a deviation of 44° when the angle of incidence is either 42° or 62° . What is the angle of incidence when it is undergoing a minimum deviation?

- (A) 45° (B) 49° (C) 40° (D) 55°

Q.21 A thin prism of angle 5° is placed at a distance of 10 cm from the object. What is the distance of the image from the object? (Given μ of prism = 1.5)


- (A) $\frac{\pi}{8}\text{cm}$ (B) $\frac{\pi}{12}\text{cm}$
 (C) $\frac{5\pi}{36}\text{cm}$ (D) $\frac{\pi}{7}\text{cm}$

Q.22 Light ray is incident on a prism of angle $A=60^\circ$ and refractive index $\mu = \sqrt{2}$. The angle of incidence at which the emergent ray grazes the surface is given by

- (A) $\sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right)$ (B) $\sin^{-1}\left(\frac{1-\sqrt{3}}{2}\right)$
 (C) $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$ (D) $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$

Previous Years' Questions

Q.1 A student measures the focal length of convex lens by putting an object pin at a distance ' u ' from the lens and measuring the distance ' v ' of the image pin. The graph between ' u ' and ' v ' plotted by the student should look like **(2002)**



Q.2 An experiment is performed to find the refractive index of glass using a travelling microscope. In this experiment, distances are measured by **(2003)**


- (A) A vernier scale provided on the microscope
 (B) A standard laboratory scale
 (C) A meter scale provided on the microscope
 (D) A screw gauge provided on the microscope

Q.3 Two transparent media of refractive indices μ_1 and μ_3 have a solid lens shaped transparent material of refractive index μ_2 between them as shown in the figures in Column II. A ray traversing these media is also shown in the figures. In Column I, the different

relationships between μ_1, μ_2 and μ_3 are given. Match them to the ray diagram shown in Column II. (2007)

Column I	Column II
(A) $\mu_1 < \mu_2$	(p)
(B) $\mu_1 > \mu_2$	(q)
(C) $\mu_2 = \mu_3$	(r)
(D) μ	(s)
	(t)

Q.4 A ray OP of monochromatic light is incident on the face AB of prism ABCD near vertex B at an incident angle of 60° (see figure) If the refractive index of the material of the prism is $\sqrt{3}$, which of the following is (are) correct? (2009)



- (A) The ray gets totally internally reflected at face CD.
- (B) The ray comes out through face AD.
- (C) The angle between the incident ray and the emergent ray is 90° .
- (D) The angle between the incident ray and the emergent ray is 120° .

Q.5 The focal length of a thin biconvex lens is 20cm. When an object is moved from a distance of 25 cm in front of it to 50 cm, the magnification of its image changes from m_{25} to m_{50} . The ratio $\frac{m_{25}}{m_{50}}$ is (2005)

Q.6 An object at distance 2.4 m in front of a lens forms a sharp image on a film at distance 12 cm behind the lens. A 1-cm-thick glass plate of refractive index 1.50 is interposed between lens and film with its plane faces parallel to film. At what distance (from lens), should the object be shifted to be in sharp focus on film? (2009)

- (A) 7.2 m
- (B) 2.4 m
- (C) 3.2 m
- (D) 5.6 m

Q.7 A spectrometer gives the following reading when used to measure the angle of prism.

Main scale reading: 58.5°

Vernier scale reading: 09 divisions

Given that 1 division on the main scale corresponds to 0.5° . Total divisions on the vernier scale are 30 and match with 29 divisions of the main scale. The angle of the prism from the above data (2011)

- (A) 58.59°
- (B) 58.77°
- (C) 58.65°
- (D) 59°

Q.8 An initially parallel cylindrical beam travels in a medium of refractive index $\mu(I) = \mu_0 + \mu_2 I$, where μ_0 and μ_2 are positive constants, and I is the intensity of the light beam. The intensity of the beam is decreasing with an increasing radius.

As the beam enters the medium, it will (2013)

- (A) Diverge
- (B) Converge
- (C) Diverge near the axis and converge near the periphery
- (D) Travel as a cylindrical beam

Q.9 The initial shape of the wave in front of the beam is (2000)


- (A) Convex

- (B) Concave
 (C) Convex near the axis and concave near the periphery
 (D) Planar

Q.10 An object 2.4 m in front of a lens forms a sharp image on a film 12 cm behind the lens. A glass plate 1 cm thick, of refractive index 1.50 is interposed between lens and film with its plane faces parallel to film. At what distance (from lens) should object be shifted to be in sharp focus on film? **(2012)**

- (A) 7.2 m (B) 2.4 m (C) 3.2 m (D) 5.6 m

Q.11 The graph between angle of deviation (δ) and angle of incidence (i) for a triangular prism is represented by: **(2013)**



Q.12 Diameter of plano-convex lens is 6 cm and thickness at the centre is 3 mm. If speed of light in material of lens is 2×10^8 m/s, the focal length of the lens is: **(2013)**

- (A) 20 cm (B) 30 cm (C) 10 cm (D) 15 cm

Q.13 A thin convex lens made from crown glass ($\mu = \frac{3}{2}$) has focal length f . When it is measured in two different liquids having refractive indices $\frac{4}{3}$ and $\frac{5}{3}$, it has the focal lengths f_1 and f_2 respectively. The correct relation between the focal lengths is: **(2014)**

- (A) $f_2 > f$ and f_1 becomes negative
 (B) f_1 and f_2 both become negative
 (C) $f_1 = f_2 < f$
 (D) $f_1 > f$ and f_2 becomes negative

Q.14 A green light is incident from the water to the air – water interface at the critical angle (θ). Select the correct statement **(2014)**


(A) The spectrum of visible light whose frequency is more than that of green light will come out to the air medium.

(B) The entire spectrum of visible light will come out of the water at various angles to the normal.

(C) The entire spectrum of visible light will come out of the water at an angle of 90° to the normal.

(D) The spectrum of visible light whose frequency is less than that of green light will come out to the air medium.

Q.15 Monochromatic light is incident on a glass prism of angle A . If the refractive index of the material of the prism is μ , a ray, incident at an angle θ , on the face AB would get transmitted through the face AC of the prism provided: **(2015)**



(A) $\theta < \sin^{-1} \left[\mu \sin \left(A - \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$

(B) $\theta > \cos^{-1} \left[\mu \sin \left(A + \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$

(C) $\theta < \cos^{-1} \left[\mu \sin \left(A + \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$

(D) $\theta > \sin^{-1} \left[\mu \sin \left(A - \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$

Q.16 Assuming human pupil to have a radius of 0.25 cm and a comfortable viewing distance of 25 cm, the minimum separation between two objects that human eye can resolve at 500 nm wavelength is: **(2015)**

- (A) 30 μm (B) 100 μm
 (C) 300 μm (D) 1 μm

Q.17 In an experiment for determination of refractive index of glass of a prism by $i - \delta$, plot, it was found that a ray incident at angle 35° , suffers a deviation of 40° and that it emerges at angle 79° . In that case which of the following is closest to the maximum possible value of the refractive index? **(2016)**


- (A) 1.6 (B) 1.7 (C) 1.8 (D) 1.5

JEE Advanced/Boards

Exercise 1


Q.1 Two flat mirrors have their reflecting surfaces facing each other, with an edge of one mirror in contact with an edge of the other so that the angle between the mirrors is 60° . Find all the angular positions of the image with respect to x-axis. Take the case when a point object is between the mirrors at $(1, 1)$. Point of intersections is $(0, 0)$ and 1st mirror is along the x-axis.

Q.2 In the figure shown, AB is a plane mirror of length 40 cm placed at a height 40 cm from ground. There is a light source S at a point on the ground. Find the minimum and maximum height of a man (eye height) required to see the image of the source if he is standing at a point P on the ground as shown in the Fig.



Q.3 A plane mirror of circular shape with radius $r=20$ cm is fixed to the ceiling. A bulb is placed on the axis of the mirror. A circular area of radius $R=1$ on the floor is illuminated after the reflection of light from the mirror. The height of the room is 3 m. What is the maximum distance from the center of the mirror and the bulb so that the required area is illuminated?

Q.4 A concave mirror of focal length 20 cm is cut into two parts from the middle, and the two parts are moved perpendicularly by a distance 1 cm from the previous principal axis AB. Find the distance between the images formed by the two parts?



Q.5 A balloon is rising up along the axis of a concave mirror of radius of curvature 20 m. A ball is dropped from the balloon at a height 15 m from the mirror when the balloon has velocity 20 m/s. Find the speed of the image of the ball formed by a concave mirror after 4 s? [Take: $g = 10 \text{ m/s}^2$]


Q.6 An observer whose least distance of distinct vision is 'd' views his own face in a convex mirror of radius or curvature 'r'. Prove that the magnification produced cannot exceed $\frac{r}{d + \sqrt{d^2 + r^2}}$.

Q.7 A surveyor on one bank of canal observes the images of the 4-inch mark and 17-ft mark on a vertical staff, which is partially immersed in the water and held against the bank directly opposite to him. He sees that the reflected and refracted rays come from the same point which is the center of the canal. If the 17-ft mark and the surveyor's eye are both 6 ft above the water level, estimate the width of the canal, assuming that the refractive index of the water is $4/3$. Zero mark is at the bottom of the canal.

Q.8 A ray of light travelling in air is incident at a grazing angle (incident angle = 90°) on a long rectangular slab of a transparent medium of thickness $t=1.0$ (see figure). The point of incidence is the origin A $(0, 0)$. The medium has a variable index of refraction $n(y)$ given by

$$n(y) = [ky^{3/2} + 1]^{-1/2}, \text{ where } k = 1.0 \text{ m}^{-3/2},$$

the refractive index of air is 1.0



(i) Obtain a relation between the slope (dy/dx) of the trajectory of the ray at a point B (x, y) in the medium and the incident angle at that point.


(ii) Find the value of $n \sin i$.

(iii) Obtain an equation for the trajectory $y(x)$ of the ray in the medium.

(iv) Determine the coordinate (x_1, y_1) of point P, where the ray intersects the upper surface of the slab–air boundary.


(v) Indicate the path of the ray subsequently.

Q.9 A uniform, horizontal beam of light is incident upon a quarter cylinder of radius $R = 5$ cm and has a refractive index $2/\sqrt{3}$. A patch on the table at a distance 'x' from the cylinder is unilluminated. Find the value of 'x'?




Q.10 An opaque cylindrical tank with an open top has diameter of 3.00 m and is fully filled with water. When the sunlight reaches at an angle of 37° above the horizon, sunlight ceases to illuminate any part of the bottom of the tank. How deep the tank is?

Q.11 A beam of parallel rays of width b propagates in a glass at an angle θ to its plane face. The beam width after it enters into air through this face is _____ if the refractive index of glass is μ .




Q.12 A parallel beam of light falls normally on the first face of a prism of a small angle. At the second face, it is partly transmitted and partly reflected, and the reflected beam strikes at the first face again and emerges from it in a direction by making an angle $6^\circ 30'$ with the reversed direction of the incident beam. The refracted beam has undergone a deviation of $1^\circ 15'$ from the original direction. Find the refractive index of the glass and the angle of the prism.

Q.13 A light ray I is incident on a plane mirror M. The mirror is rotated in the direction as shown in the figure by an arrow at a frequency $9/\pi$ rev/sec. The light reflected by the mirror is received on the wall W at a distance 10 m from the axis of rotation. When the angle of incidence becomes 37° , find the speed of the spot (a point) on the wall?




Q.14 The diagram shows five isosceles right-angled prisms. A light ray incident at 90° at the first face emerges at the same angle with the normal from the last face. Find the relation between the refractive indices?

Q.15 Two rays travelling parallel to the principal axis




strike a large plano-convex lens of a refractive index of 1.60. If the convex face is spherical, a ray near the edge does not pass through the focal point (spherical aberration occurs). If this face has a radius of curvature of 20.0 cm and the two rays are at $h_1 = 0.50$ cm and $h_2 = 12.0$ cm from the principal axis, then find the difference in the positions where they cross the principal axis.



Q.16 A room contains air in which the speed of sound is 340m/s. The walls of the room are made of concrete, in which the speed of sound is 1700m/s. (a) Find the critical angle for the TIR of sound at the concrete–air boundary. (b) In which medium must the sound be undergone the TIR?


Q.17 A rod made of glass ($\mu = 1.5$) and of square cross section is bent as shown in the figure. A parallel beam of light falls perpendicularly on the plane flat surface A. Referring to the diagram, d is the width of a side, and R is the radius of inner semicircle. Find the maximum value of ratio $\frac{d}{R}$ so that all the light rays entering the glass through surface A emerge from the glass through surface B.



Q.18 A prism of refractive index $\sqrt{2}$ has a refracting angle of 30° . One of the refracting surfaces of the prism is polished. For the beam of monochromatic light to retrace its path, find the angle of incidence on the refracting surface.

Q.19 An equilateral prism deviates a ray by 23° for two angles of incidence differing by 23° . Find μ of the prism?

Q.20 A ray is incident on a glass sphere as shown in the figure. The opposite surface of the sphere is partially coated with silver. If the net deviation of the ray transmitted at the partially silvered surface is $1/3$ rd of the net deviation suffered by the ray reflected at the partially silvered surface (after emerging out of the sphere), find the refractive index of the sphere.




Q.21 Two thin similar watch glass pieces are joined together front to front, with rear convex portion is coated with silver, and the combination of glass pieces is placed at a distance $\alpha = 60$ cm from a screen. A small object is placed normal to the optical axis of the combination such that its two times magnified image is formed on the screen. If air between the glass pieces is replaced by water ($\mu = 4/3$), calculate the distance through which the object must be displaced so that a sharp image is again formed on the screen.

Q.22 A spherical light bulb with a diameter of 3.0 cm radiates light equally in all directions, with a power of 4.5π W. (a) Find the light intensity at the surface of the bulb. (b) Find the light intensity 7.50 m from the center

of the bulb. (c) At this 7.50-m distance, a convex lens is set up with its axis pointing toward the bulb. The lens has a circular face with a diameter of 15.0 cm and a focal length of 30.0 cm. Find the diameter of the image of the bulb formed on a screen kept at the location of the image. (d) Find the light intensity at the image.

Q.23 A thin plano-convex lens fits exactly into a plano-concave lens with their plane surface parallel to each other as shown in the figure. The radius of curvature of the curved surface $R = 30$ cm. The lenses are made of different material having a refractive index $\mu_1 = 3/2$ and $\mu_2 = 5/4$ as shown in the Fig.




(i) If plane surface of the plano-convex lens is coated with silver, then calculate the equivalent focal length of this system and also the nature of this equivalent mirror.

(ii) An object having a transverse length of 5 cm is placed on the axis of equivalent mirror (in par 1) at a distance 15 cm from the equivalent mirror along the principal axis. Find the transverse magnification produced by the equivalent mirror.

Q.24 Two identical convex lenses L_1 and L_2 are placed at a distance of 20 cm from each other on the common principal axis. The focal length of each lens is 15 cm, and lens L_2 is placed right to lens L_1 . A point object is placed at a distance of 20 cm left to lens L_1 on the common principal axis of two lenses. Find where a convex mirror of radius of curvature 5 cm should be placed to the right of L_2 so that the final image coincides with the object?

Q.25 A thin equiconvex lens of glass of a refractive index $\mu = 3/2$ and a focal length 0.3 m in air is sealed into an opening at one end of a tank filled with water ($\mu = 4/3$). On the opposite side of the lens, a mirror is placed inside the tank on the tank wall perpendicular to the lens axis, as shown in the figure. The distance between the lens and the mirror is 0.8 m. A small object is placed outside the tank in front of the lens at a distance of 0.9 m from the lens along its axis. Find the

position (relative to the lens) of the image of the object formed by the system.



- (A) $f/2, 1/2$ (B) $f, 1/4$
 (C) $3f/4, 1/2$ (D) $f, 3/4$

Exercise 2

Single Correct Choice Type

Q.1 An object is placed in front of a convex mirror at a distance of 50 cm. A plane mirror is introduced covering the lower half of the convex mirror. If the distance between the object and the plane mirror is 30 cm, there is no gap between the images formed by the two mirrors. The radius of the convex mirror is:

- (A) 12.5 cm (B) 25 cm
 (C) 50 cm (D) 100 cm

Q.2 An infinitely long rod lies along the axis of a concave mirror of a focal length f . The near end of the rod is at a distance $u > f$ from the mirror. Its image will have a length.

- (A) $f^2 / (u - f)$ (B) $uf / (u - f)$
 (C) $f^2 / (u + f)$ (D) $uf / (u + f)$

Q.3 A luminous point object is moving along the principal axis of a concave mirror of a focal length 12 cm toward it. When its distance from mirror is 20 cm, its velocity is 4 cm/s. The velocity of the image in cm/s at that instant is:


- (A) 6 toward the mirror
 (B) 6 away from the mirror
 (C) 9 away from the mirror
 (D) 9 toward the mirror

Q.4 A thin lens has a focal length f , and its aperture has a diameter d . It forms an image of intensity I . Now the central part of the aperture up to diameter $(d/2)$ is blocked by an opaque paper. The focal length and image intensity would change to

Q.5 An object is placed in front of a thin convex lens of focal length 30 cm, and a plane mirror is placed 15 cm behind the lens. If the final image of the object coincides with the object, the distance of the object from the lens is

- (A) 60 cm (B) 30 cm
 (C) 15 cm (D) 25 cm

Q.6 A converging lens of a focal length 20 cm and diameter 5 cm is cut along line AB. The part of the lens shaded in the diagram is used to form an image of a point P placed 30 cm away from it on line XY, which is perpendicular to the plane of the lens. The image of P will be formed.



- (A) 0.5 cm above XY (B) 1 cm below XY
 (C) On XY (D) 1.5 cm below XY

Q.7 A screen is placed 90 cm away from an object. The image of the object on the screen is formed by a convex lens at two different locations separated by 20 cm. The focal length of the lens is

- (A) 18 cm (B) 21.4 cm
 (C) 60 cm (D) 85.6 cm

Q.8 In the above problem, if the sizes of the images formed on the screen are 6 cm and 3 cm, then the height of the object is nearly:


- (A) 4.2 cm (B) 4.5 cm
 (C) 5 cm (D) 9 cm

Q.9 A concave mirror cannot form

- (A) A virtual image of a virtual object
 (B) A virtual image of a real object
 (C) A real image of a real object
 (D) A real image of a virtual object


Multiple Correct Choice Type

Q.10 A reflecting surface is represented by the equation $y = \frac{2L}{\pi} \sin\left(\frac{\pi x}{L}\right), 0 \leq x \leq L$. A ray travelling horizontally becomes vertical after reflection. The coordinates of the point(s) where this ray is incident is




- (A) $\left(\frac{L}{4}, \frac{\sqrt{2L}}{\pi}\right)$ (B) $\left(\frac{L}{3}, \frac{\sqrt{3L}}{\pi}\right)$
 (C) $\left(\frac{3L}{4}, \frac{\sqrt{2L}}{\pi}\right)$ (D) $\left(\frac{2L}{3}, \frac{\sqrt{3L}}{\pi}\right)$

Q.11 In the figure shown, consider the first reflection at the plane mirror and second at the convex mirror. AB is object.



- (A) The second image is real, inverted by $1/5$ th magnification w.r.t. AB.
 (B) The second image is virtual and erect by $1/5$ th magnification w.r.t. AB.
 (C) The second image moves toward the convex mirror.
 (D) The second image moves away from the convex mirror.

Q.12 In the diagram shown, a ray of light is incident on the interface between 1 and 2 at an angle slightly greater than the critical angle. The light undergoes TIR at this interface. After that, the light ray falls at interfaces 1 and 3, and again it undergoes TIR. Which of the following relations should hold true?



- (A) $\mu_1 < \mu_2 < \mu_3$ (B) $\mu_1^2 - \mu_2^2 > \mu_3^2$
 (C) $\mu_1^2 - \mu_3^2 > \mu_2^2$ (D) $\mu_1^2 + \mu_2^2 > \mu_3^2$

Q.13 For refraction through a small angle prism, the angle of deviation:

- (A) Increases with an increase in RI of prism.
 (B) Will decrease with an increase in RI of prism.
 (C) Is directly proportional to the angle of prism.
 (D) Will be $2D$ for a ray of RI = 2.4 if it is d for a ray of RI = 1.2.


Q.14 For the refraction of light through a prism

- (A) For every angle of deviation, there are two angles of incidence.
 (B) The light travelling inside an equilateral prism is necessarily parallel to the base when prism is set for a minimum deviation.
 (C) There are two angles of incidence for a maximum deviation.
 (D) The angle of minimum deviation will increase if refractive index of prism is increased keeping the outside medium unchanged if $\mu_p > \mu_s$.

Q.15 A man of height 170 cm wants to see his complete image in a plane mirror (while standing). His eyes are at a height of 160 cm from the ground.


- (A) Minimum length of the mirror = 80 cm.
 (B) Minimum length of the mirror = 85 cm.
 (C) Bottom of the mirror should be at a height 80 cm or less.
 (D) Bottom of the mirror should be at a height 85 cm.

Q.16 A flat mirror M is arranged parallel to a wall W at a distance l from it. The light produced by a point source S kept on the wall is reflected by the mirror and produces a light spot on the wall. The mirror moves with a velocity v toward the wall.



- (A) The spot of light will move with the speed v on the wall.
 (B) The spot of light will not move on the wall.
 (C) As the mirror comes closer, the spot of light will become larger and shift away from the wall with a speed larger than v .
 (D) The size of the light spot on the wall remains the same.


Q.17 Two reflecting media are separated by a spherical interface as shown in the figure. PP' is the principal axis; μ_1 and μ_2 are the medium of refraction, respectively, then,



- (A) If $\mu_2 > \mu_1$, then there cannot be a real image of a real project.
 (B) If $\mu_2 > \mu_1$, then there cannot be a real image of a virtual object.
 (C) If $\mu_1 > \mu_2$, then there cannot be a virtual image of a virtual object.
 (D) If $\mu_1 > \mu_2$, then there cannot be a real image of a real object.

Q.18 A luminous point object is placed at O , whose image is formed at I as shown in the figure. AB is the optical axis. Which of the following statements are correct?

- (A) If a lens is used to obtain an image, the lens must be converging.
 (B) If a mirror is used to obtain an image, the mirror must be a convex mirror having a pole at the point of intersection of lines OI and AB .
 (C) Position of the principal focus of mirror cannot be found.
 (D) I is a real image.



Q.19 A lens is placed in the XYZ coordinate system such that its optical center is the origin and the principal axis is along the x -axis. The focal length of the lens is 20 cm. A point object has been placed at the point $(-40 \text{ cm}, +1 \text{ cm}, -1 \text{ cm})$. Which of the following are correct about coordinates of the image?

- (A) $x = 40 \text{ cm}$ (B) $y = +1 \text{ cm}$
 (C) $z = +1 \text{ cm}$ (D) $z = -1 \text{ cm}$

Q.20 Which of the following can form a diminished, virtual and erect image of your face.

- (A) Converging mirror (B) Diverging mirror
 (C) Converging lens (D) Diverging lens


Assertion Reasoning Type

Q.21 Statement-I: If a source of light is placed in front of rough wall, its image is not seen.

Statement-II: The wall does not reflect light.

- (A) Statement-I is true, and statement-II is true; statement-II is correct explanation for statement-I
 (B) Statement-I is true, and statement-II is true; statement-II is NOT the correct explanation for statement-I.
 (C) Statement-I is true, and statement-II is false.
 (D) Statement-I is false, and statement-II is true.

Q.22 Statement-I: As the distance x of a parallel ray from axis increases, the focal length decreases



Statement-II: As x increases, the distance from the pole to the point of intersection of a reflected ray with the principal axis decreases.

- (A) Statement-I is true, and statement-II is true; statement-II is correct explanation for statement-I.

(B) Statement-I is true, and statement-II is true; statement-II is NOT the correct explanation for statement-I.

(C) Statement-I is true, and statement-II is false.

(D) Statement-I is false, and statement-II is true.

Q.23 Statement-I: When an object dipped in a liquid is viewed normally, the distance between the image and the object is independent of the height of the liquid above the object.

Statement-II: The normal shift is independent of the location of the slab between the object and the observer.


(A) Statement-I is true, statement-II is true, and statement-II is the correct explanation for statement-I.

(B) Statement-I is true, statement-II is true, and statement-II is NOT the correct explanation for statement-I.

(C) Statement-I is true, and statement-II is false.

(D) Statement-I is false, and statement-II is true.

Q.24 Statement-I: When two plane mirrors are kept perpendicular to each other as shown in the figure (O is the point object), three images will be formed.



Statement-II: In case of a multiple reflection, the image of one surface can act as an object for the next surface.

(A) Statement-I is true, statement-II is true, and statement-II is the correct explanation for statement-I.

(B) Statement-I is true, statement-II is true, and statement-II is NOT the correct explanation for statement-I.

(C) Statement-I is true, and statement-II is false.

(D) Statement-I is false, and statement-II is true.

Q.25 Statement-I: Keeping a point object fixed, if a plane mirror is moved, the image will definitely move.

Statement-II: In case of a plane mirror, the distance between a point object and its image from a given point on mirror is equal.

(A) Statement-I is true, statement-II is true, and statement-II is the correct explanation for statement-I.

(B) Statement-I is true, statement-II is true, and statement-II is NOT the correct explanation for statement-I.

(C) Statement-I is true, and statement-II is false.

(D) Statement-I is false, and statement-II is true.

Comprehension Type

Paragraph 1: Spherical aberration in spherical mirrors is a defect that is due to the dependence of focal length ' f ' on the angle of incidence ' θ ' as shown in the figure is given by

$$f = R - \frac{R}{2} \sec \theta,$$

where R is radius of curvature of mirror and θ is the angle of incidence. The rays that are close to the principal axis are called marginal rays. As a result, different rays focus at different points and the image of a point object is not a point.

Q.26 If f_p and f_m represent the focal length of paraxial and marginal rays, respectively, then the correct relationship is:

(A) $f_p = f_m$ (B) $f_p > f_m$


(C) $f_p < f_m$ (D) none

Q.27 If the angle of incidence is 60° , then the focal length of this ray is:

(A) R (B) $R/2$

(C) $2R$ (D) 0

Paragraph 2: A student is performing Young's double slit experiment. There are two slits S_1 and S_2 . The distance between them is d . There is a large screen at a distance D ($D \gg d$) from the slits. The setup is shown in the following figure. A parallel beam of light is incident on it. A monochromatic light of wavelength λ is used. The initial phase difference between the two slits behave as two coherent sources of light is zero. The intensities of light waves on the screen coming out of S_1 and S_2 are same, i.e. I_0 . In this situation, the principal maximum is formed at point P. At the point on screen where the principal maximum is formed, the phase difference between two interfering waves is zero.



Q.28 The total deviation suffered by the ray falling on the mirror at an angle of incidence 60° is

- (A) 180° (B) 90°
(C) Cannot be determined (D) None

Q.29 For paraxial rays, focal length approximately is

- (A) R (B) $R/2$
(C) $2R$ (D) none

Q.30 Which of the following statements are correct regarding spherical aberration:

- (A) It can be completely eliminated.
(B) It cannot be completely eliminated, but it can be minimized by allowing either paraxial or marginal rays to hit the mirror.
(C) It is reduced by taking mirrors with large aperture.
(D) None of these.

Q.31 Initially, the distance of third minima from principal maxima will be

- (A) $\frac{3\lambda D}{2d}$ (B) $\frac{3\lambda D}{d}$
(C) $\frac{5\lambda D}{4d}$ (D) $\frac{5\lambda D}{2d}$

Q.32 A glass slab of thickness t and refractive index μ is introduced before S_2 . Now, P does not remain the point of principal maximum. Suppose the principal maximum forms at a point P' on screen, then PP' is equal to

- (A) $\frac{tD(\mu-1)}{d}$ (B) $\frac{tD(\mu-1)}{2d}$
(C) $\frac{D(\mu-1)}{t}$ (D) $\frac{D(\mu-1)}{d}$

Q.33 Use the statement given in previous question. Now, a parallel beam is incident at an angle α w.r.t. line OP, such that the principal maximum again comes at

point P (see figure). The value of α is

- (A) $\sin^{-1} \frac{t(\mu-1)}{d}$ (B) $\cos^{-1} \frac{t(\mu-1)}{d}$
(C) $\sin^{-1} \frac{t(\mu-1)D}{d}$ (D) $\sin^{-1} \frac{tD}{d}$

Q.34 Match the Column


Column I	Column II
(A) Converging system	(p) Convex lens
(B) Concave lens	(q) Concave lens
(C) A virtual image is formed by	(r) Concave mirror
(D) Magnification < 1 is possible with	(s) Convex mirror

Previous Years' Questions

Q.1 A student performed the experiment of determination of the focal length of a concave mirror by $u-v$ method using an optical bench of length 1.5 m. The focal length of the mirror used is 24 cm. The maximum error in the location of the image can be 0.2 cm. The five sets of (u, v) values recorded by the student (in cm) are: (42, 56), (48, 48), (60, 40), (66, 33) and (78, 39). The data set (s) that cannot come from experiment and is (are) incorrectly recorded, is (are) **(1999)**





- (A) (42, 56) (B) (48, 48)
(C) (66, 33) (D) (78, 39)

Q.2 A light beam travels from Region I to Region IV (See figure). The refractive index in Regions I, II, III and IV are n_0 , $\frac{n_0}{6}$ and $\frac{n_0}{8}$, respectively. The angle of incidence θ for which the beam misses entering Region IV (as in the figure): **(2004)**



- (A) $\sin^{-1} \left(\frac{3}{4} \right)$ (B) $\sin^{-1} \left(\frac{1}{8} \right)$
(C) $\sin^{-1} \left(\frac{1}{4} \right)$ (D) $\sin^{-1} \left(\frac{1}{3} \right)$

Q.3 An optical component and an object S placed along its optical axis are given in column I. The distance between the object and the component can be varied. The properties of images are given in column II. Match all the properties of images from column II with the appropriate components given in column I. Indicate your answer by darkening the appropriate bubbles of the 4 x 4 matrix given in the ORS. **(2006)**

Column I	Column II
(A) 	(p) Real image
(B) 	(q) Virtual image
(C) 	(r) Magnified image
(D) 	(s) Image at infinity

Q.4 Two beams of red and violet colors are pass separately through a prism (angle of the prism is 60°). In the position of minimum deviation, the angle of refraction will be **(2007)**


- (A) 30° for both the colors
- (B) Greater for the violet color
- (C) Greater for the red color
- (D) Equal but not 30° for both the colors

Q.5 Which one of the following statements is WRONG in the context of X-rays generated from an X-ray tube? **(2001)**

- (A) The wavelength of the characteristics X-rays decreases when the atomic number of the target increases.
- (B) The cutoff wavelength of the continuous X-rays depends on the atomic number of the target.

(C) The intensity of the characteristics X-rays depends on the electrical power given to the X-ray tube.


(D) Cutoff wavelength of the continuous X-rays depends on the energy of the electrons in the X-ray tube.



Paragraph: Most materials have a refractive index $n > 1$. Therefore, when a light ray from air enters a naturally occurring material, then by the Snell's law, $\frac{\sin \theta_1}{\sin \theta_2} = \frac{n_2}{n_1}$, it is understood that the refracted ray bends toward the normal, but it never emerges on the same side of the normal as the incident ray. According to the electromagnetism, the refractive index of the medium is

given by the relation, $n = \left(\frac{c}{v} \right) = \pm \sqrt{\epsilon_1 \mu_1}$, where c is the speed of electromagnetic waves in vacuum, v is its speed in the medium, ϵ_1 and μ_1 are the relative permittivity and permeability of the medium, respectively. In a normal material, both ϵ_1 and μ_1 are positive, implying positive n for the medium. When both ϵ_1 and μ_1 are negative, one must choose the negative root of n . Such materials with negative refractive indices can now be artificially prepared and are called meta-materials. They exhibit a significantly different optical behavior, without violating any physical laws. Since n is negative, it results in a change in the direction of propagation of the refracted light. However, similar to the normal materials, the frequency of light remains unchanged upon refraction even in meta-materials. **(2012)**


Q.6 For light incident from air on a meta-material, the appropriate ray diagram is



Q.7 Choose the correct statement.

- (A) The speed of light in the meta-material is $v = c|n|$.
- (B) The speed of light in the meta-material is $v = \frac{c}{|n|}$.
- (C) The speed of light in the meta-materials is $v = c$.
- (D) The wavelength of the light in the meta-material (λ_m) is given by $\lambda_m = \lambda_{\text{air}}|n|$, where λ_{air} is the wavelength of the light in air.

Q.8 A biconvex lens is formed with two thin plano-convex lenses as shown in the figure. Refractive index n of the first lens is 1.5 and that of the second lens is 1.2. Both curved surfaces are of same radius of curvature $R = 14$ cm. For this biconvex lens, for an object distance of 40 cm, the image distance will be **(2009)**




Q.9 A biconvex lens of focal length 15 cm is in front of a plane mirror. The distance between the lens and the mirror is 10 cm. A small object is kept at a distance of 30 cm from the lens. The final image is **(2011)**

(A) Virtual and at a distance of 16 cm from the mirror

- (B) Real and at a distance of 16 cm from the mirror
- (C) Virtual and at a distance of 20 cm from the mirror
- (D) Real and at a distance of 20 cm from the mirror

Q.10 The image of an object approaching a convex mirror of a radius of curvature 20 m along its optical axis moves from $\frac{25}{3}$ m to $\frac{50}{7}$ m in 30 s. What is the speed of the object in km/h? **(2011)**


Q.11 A bi-convex lens is formed with two thin plano-convex lenses as shown in the figure. Refractive index n of the first lens is 1.5 and that of the second lens is 1.2. Both the curved surfaces are of the same radius of curvature $R = 14$ cm. For this bi-convex lens, for an object distance of 40 cm, the image distance will be – **(2012)**



- (A) – 280.0 cm (B) 40.0 cm (C) 21.5 cm (D) 13.3 cm


Q.12 A transparent slab of thickness d has a refractive index $n(z)$ that increases with z . Here z is the vertical distance inside the slab, measured from the top. The slab is placed between two media with uniform refractive indices n_1 and $n_2 (> n_1)$, as shown in the figure. A ray of light is incident with angle θ_1 from medium 1 and emerges in medium 2 with refraction angle θ_f with a lateral displacement l . **(2016)**

Which of the following statement (s) is (are) true?



- (A) $n_1 \sin \theta_i = n_2 \sin \theta_f$
 (B) $n_1 \sin \theta_i = (n_2 - n_1) \sin \theta_f$
 (C) l is independent of n_2
 (D) l is dependent on $n(z)$

Q.13 A small object is placed 50 cm to the left of a thin convex lens of focal length 30 cm. A convex spherical mirror of radius of curvature 100 cm is placed to the right of the lens at a distance of 50 cm. The mirror is tilted such that the axis of the mirror is at an angle $\theta = 30^\circ$ to the axis of the lens, as shown in the figure. (2015)



If the origin of the coordinate system is taken to be at the centre of the lens, the coordinates (in cm) of the point (x, y) at which the image is formed are (2016)

- (A) $(25, 25\sqrt{3})$ (B) $(125/3, 25/\sqrt{3})$
 (C) $(50 - 25\sqrt{3}, 25)$ (D) $(0, 0)$

Q.14 A ray of light travelling in the direction $\frac{1}{2}(\hat{i} + \sqrt{3}\hat{j})$ is incident on a plane mirror. After reflection, it travels along the direction $\frac{1}{2}(\hat{i} + \sqrt{3}\hat{j})$. The angle of incidence is (2013)


- (A) 30° (B) 45° (C) 60° (D) 75°

Q.15 The image of an object, formed by a plano-convex lens at a distance of 8 m behind the lens, is real and is one-third the size of the object. The wavelength of light inside the lens is $\frac{2}{3}$ times the wavelength in free space.

The radius of the curved surface of the lens is (2013)

- (A) 1 m (B) 2 m (C) 3 m (D) 4 m

Q.16 A right angled prism of refractive index μ_1 , is placed in a rectangular block of refractive index μ_2 , which is surrounded by a medium of refractive index μ_3 , as shown in the figure. A ray of light 'e' enters the rectangular block at normal incidence. Depending upon the relationships between μ_1 , μ_2 and μ_3 , it takes one of the four possible paths 'ef', 'eg', 'eh', or 'ei'.



Match the paths in list I with conditions of refractive indices in list II and select the correct answer using the codes given below the lists: (2013)


	List I		List II
P.	$e \rightarrow f$	1.	$\mu_1 > \sqrt{2} \mu_2$
Q.	$e \rightarrow g$	2.	$\mu_2 > \mu_1$ and $\mu_2 > \mu_3$
R.	$e \rightarrow h$	3.	$\mu_1 = \mu_2$
S.	$e \rightarrow i$	4.	$\mu_2 < \mu_1 < \sqrt{2} \mu_2$ and $\mu_2 > \mu_3$

Codes:

	P	Q	R	S
A	2	3	1	4
B	1	2	4	3
C	4	1	2	3
D	2	3	4	1


Q.17 A transparent thin film of uniform thickness and refractive index $n_1 = 1.4$ is coated on the convex spherical surface of radius R at one end of a long solid glass cylinder of refractive index $n_2 = 1.5$, as shown in the figure. Rays of light parallel to the axis of the cylinder traversing through the film from air to glass

get focused at distance f_1 from the film, while rays of light traversing from glass to air get focused at distance f_2 from the film. Then **(2014)**



- (A) $|f_1| = 3R$ (B) $|f_1| = 2.8R$
 (C) $|f_2| = 2R$ (D) $|f_2| = 1.4R$

Q.18 A point source S is placed at the bottom of a transparent block of height 10 mm and refractive index 2.72. It is immersed in a lower refractive index liquid as shown in the figure. It is found that the light emerging from the block to the liquid forms a circular bright spot of diameter 11.54 mm on the top of the block. The refractive index of the liquid is **(2014)**



- (A) 1.21 (B) 1.30 (C) 1.36 (D) 1.42

Q.19 Four combinations of two thin lenses are given in list I. The radius of curvature of all curved surfaces is r and the refractive index of all the lenses is 1.5. Match lens combinations in List I with their focal length in list II and select the correct answer using the code given below the lists. **(2014)**

	List I		List II
P			$2r$
Q			$r/2$

	List I		List II
R			$-r$
S			r


Code:

- (A) P-1, Q-2, R-3, S-4
 (B) P-2, Q-4, R-3, S-1
 (C) P-4, Q-1, R-2, S-3
 (D) P-2, Q-1, R-3, S-4

Q.20 Consider a concave mirror and a convex lens (refractive index = 1.5) of focal length 10 cm each, separated by a distance of 50 cm in air (refractive index = 1) as shown in the figure. An object is placed at a distance of 15 cm from the mirror. Its erect image formed by this combination has magnification M_1 . When the set-up is kept in a medium of refractive index $7/6$, the magnification becomes M_2 . The magnitude **(2015)**


$\left| \frac{M_2}{M_1} \right|$ is

Q.21 Two identical glass rods S_1 and S_2 (refractive index = 1.5) have one convex end of radius of curvature 10 cm. They are placed with the curved surfaces at a distance d as shown in the figure, with their axes (shown by the dashed line) aligned. When a point source of light P is placed inside rod S_1 on its axis at a distance of 50 cm from the curved face, the light rays emanating from it are found to be parallel to the axis inside S_2 . The distance d is **(2015)**




- (A) 60 cm (B) 70 cm
 (C) 80 cm (D) 90 cm

Q.22 A monochromatic beam of light is incident at 60° on one face of an equilateral prism of refractive index n and emerges from the opposite face making an angle $\theta(n)$ with the normal (see the figure). For $n = \sqrt{3}$ the value of θ is 60° and $\frac{d\theta}{dn} = m$. The value of m is **(2015)**



Paragraph 1: Light guidance in an optical fiber can be understood by considering a structure comprising of thin solid glass cylinder of refractive index n_1 surrounded by a medium of lower refractive index n_2 . The light guidance in the structure takes place due to successive total internal reflections at the interface of the media n_1 and n_2 as shown in the figure. All rays with the angle of incidence i less than a particular value i_m are confined in the medium of refractive index n_1 . The numerical aperture (NA) of the structure is defined as $\sin i_m$



Q.23 For two structures namely S_1 with $n_1 = \sqrt{45}/4$ and $n_2 = 3/2$ and S_2 with $n_1 = 8/5$ and $n_2 = 7/5$ and taking the refractive index of water to be $4/3$ and that of air to be 1, the correct option(s) is(are) **(2015)**


(A) NA of S_1 immersed in water is the same as that of S_2 immersed in a liquid of refractive index $\frac{16}{3\sqrt{15}}$

(B) NA of S_1 immersed in liquid of refractive index $\frac{6}{\sqrt{15}}$ is the same as that of S_2 immersed in water

(C) NA of S_1 placed in air is the same as that of S_2 immersed in liquid of refractive index $\frac{4}{\sqrt{15}}$.

(D) NA of S_1 placed in air is the same as that of S_2 placed in water.

Q.24 A parallel beam of light is incident from air at an angle α on the side PQ of a right angled triangular prism of refractive index $n = \sqrt{2}$. Light undergoes total internal reflection in the prism at the face PR when α has a minimum value of 45° . The angle θ of the prism is **(2016)**



(A) 15° (B) 22.5° (C) 30° (D) 45°

Q.25 A plano-convex lens is made of a material of refractive index n . When a small object is placed 30 cm away in front of the curved surface of the lens, an image of double the size of the object is produced. Due to reflection from the convex surface of the lens, another faint image is observed at a distance of 10 cm away from the lens. Which of the following statement(s) is(are) true? **(2016)**

(A) The refractive index of the lens is 2.5

(B) The radius of curvature of the convex surface is 45 cm

(C) The faint image is erect and real

(D) The focal length of the lens is 20 cm

MASTERJEE Essential Questions

JEE Main/Boards

Exercise 1

Q.11 Q.12 Q.24

Exercise 2

Q. 3 Q.5 Q.7
Q.8 Q.10 Q.18
Q.23 Q.27 Q.37

JEE Advanced/Boards

Exercise 1

Q.3 Q.4 Q.8
Q.13 Q.22 Q.23

Exercise 2

Q.6 Q.9 Q.11
Q.24 Q.25 Q.30
Q.31 Q.33

Answer Key

JEE Main/ Boards

Exercise 1

Q.1 $2.12 \times 10^8 \text{ ms}^{-1}$

Q.2 $u = -30 \text{ cm}$

Q.4 $\frac{3 \times 10^8}{\sqrt{2}} \text{ m/s}$

Q.5 The image is formed on the same side of object.

Q.9 (b) (i) When the refractive index of the medium increases, the resolving power increases.

(ii) When the wavelength of the radiation increases, the resolving power decreases.

Q.11 For convex lens +7.5 cm, for concave lens +48 cm.

Q.12 A virtual image of height 10 cm is formed at a distance of 25 cm from the lens on the same side of the object.

Q.14 The angle of deviation is decreased.

Q.16 (i) +5 D; (ii) 10 cm

Q.17 25 cm

Q.20 $f_{\text{eq}} = \infty$

Q.21 (ii) -10

Q.23 (a) 5

Q.24 2.14

Q.25 $f_o = 40 \text{ cm}; f_e = 5 \text{ cm}$

Exercise 2

Single Correct Choice Type

Q.1 A

Q.2 B

Q.3 A

Q.4 C

Q.5 B

Q.6 A

Q.7 D

Q.8 D

Q.9 A

Q.10 A

Q.11 C

Q.12 D

Q.13 B	Q.14 B	Q.15 B	Q.16 C	Q.17 D	Q.18 D
Q.19 C	Q.20 B	Q.21 C	Q.22 A		

Previous Years' Questions

Q.1 D	Q.2 A	Q.3 A → p, r; B → q, s, t; C → p, r, t; D → q, s			
Q.4 A,B,C	Q.5 6	Q.6 D	Q.7 C	Q.8 B	Q.9 D
Q.10 D	Q.11 B	Q.12 B	Q.13 D	Q.14 D	Q.15 D
Q.16 A	Q.17 D				

JEE Advanced/Boards

Exercise 1

Q.1 (i) 75°	(ii) 165°	(iii) 195°	(iv) 285°	(v) 315°	
Q.2 160 cm; 320 cm		Q.3 75 cm	Q.4 2 cm	Q.5 80 m/s	Q.7 16 ft
Q.8 (i) $\tan\theta = \frac{dy}{dx} = \cot i$		(ii) 1	(iii) $y = k^2(x/4)^4$	(iv) 4.0, 1;	
(v) It will become parallel to x-axis		Q.9 5 cm	Q.10 h=5.95 m	Q.11 Same	Q.12 1.625
Q.13 1000 m/s		Q.14 $\mu_1^2 + \mu_3^2 + \mu_5^2 = 2 + \mu_2^2 + \mu_4^2$		Q.15 9 m	
Q.16 (a) $\sin^{-1}\left(\frac{1}{5}\right)$	(b) air	Q.17 $\left(\frac{d}{R}\right)_{\max} = \frac{1}{2}$		Q.18 45°	Q.19 $\frac{\sqrt{43}}{5}$
Q.20 $\sqrt{3}$		Q.21 15 cm toward the combination			
Q.22 (a) 5000 W/m ²	(b) 0.02 W/m ²	(c) 0.214 cm	(d) 24.56 W/m ²		
Q.23 +60, +4/5		Q.24 5.9 cm, 10.9 cm			
Q.25 90 cm from the lens toward right					

Exercise 2

Single Correct Choice Type

Q.1 B	Q.2 A	Q.3 C	Q.4 D	Q.5 B	Q.6 D
Q.7 B	Q.8 A	Q.9 A			

Multiple Correct Choice Type

Q.10 B, D	Q.11 B, C	Q.12 B, C	Q.13 A, C	Q.14 A, B, D	Q.15 B, C
Q.16 B, D	Q.17 A, C	Q.18 A, D	Q.19 A, C	Q.20 B, D	

Assertion Reasoning Type

Q.21 C	Q.22 A	Q.23 D	Q.24 D	Q.25 D
---------------	---------------	---------------	---------------	---------------

Comprehension Type

Q.26 B	Q.27 D	Q.28 D	Q.29 B	Q.30 B	Q.31 D
Q.32 A	Q.33 A				

Match the Column

Q.34 A → p, r; B → q, s; C → p, q, r, s; D → p, q, r, s

Previous Years' Questions**Q.1** C, D**Q.2** B**Q.3** A → p, q, r, s; B → q; C → p, q, r, s; D → p, q, r, s**Q.4** A**Q.5** B**Q.6** C**Q.7** B**Q.8** B**Q.9** B**Q.10** 3**Q.11** B**Q.12** A, C, D**Q.13** A**Q.14** A**Q.15** C**Q.16** D**Q.17** A, C**Q.18** C**Q.19** B**Q.20** 7**Q.21** B**Q.22** 2**Q.23** A, C**Q.24** A**Q.25** A, D**Solutions****JEE Main/Boards****Exercise 1**

$$\text{Sol 1: } \mu = \frac{\sin\left(\frac{A + \delta_m}{2}\right)}{\sin\frac{A}{2}} = \frac{\sin\left(\frac{90}{2}\right)}{\sin\left(\frac{60}{2}\right)} = \frac{1}{\frac{\sqrt{2}}{2}} = \sqrt{2}$$

Here $A = 60^\circ$, $\delta_m = 30^\circ$.

$$\text{Now } \mu = \frac{c}{v} \Rightarrow v = \frac{c}{\mu} \Rightarrow v = \frac{3 \times 10^8}{\sqrt{2}} = 2.12 \times 10^8 \text{ ms}^{-1}$$

Sol 2: Lens Formula: $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$. Let object is placed at distance x from lens and image is found at distance y from lens. For real image v is positive $v = +y$, $u = -x$, $m = \frac{v}{u} = \frac{+y}{-x} = -2$ (For real image m is negative) $\Rightarrow y = 2x$ (i) $\frac{1}{v} - \frac{1}{u} = \frac{1}{y} - \frac{1}{-x} = \frac{1}{20}$ (f is positive for convex lens)

$$\Rightarrow \frac{1}{2x} + \frac{1}{x} = \frac{1}{20} \text{ (using (1)) } \Rightarrow \frac{3}{2x} = \frac{1}{20} \Rightarrow x = 30 \text{ cm}$$

$u = -30 \text{ cm}$

Sol 3: (i) Focal length of a concave mirror is independent of the medium and wavelength of light. So there will not be any change.

(ii) Focal length of a concave lens depends on the refractive index μ of the medium which in-turn depends upon the wavelength of light. μ decreases with increasing wavelength. So for red light μ will decrease.

As per lens maker's formula $\left[\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right) \right]$ as μ decrease, f increase.

Sol 4: Total internal reflection (TIR) takes place when light travels from denser medium towards rarer medium and at the interface the angle of incidence exceeds θ_c , the critical angle, and the incident beam is completely reflected at the boundary (interface). Critical angle

$$\theta_c = \sin^{-1} \left(\frac{\mu_{\text{Rarer}}}{\mu_{\text{Denser}}} \right)$$

$$45 = \sin^{-1} \left(\frac{1}{\mu} \right) \Rightarrow \frac{1}{\mu} = \frac{1}{\sqrt{2}} \Rightarrow \mu = \frac{c}{v} = \sqrt{2}$$

$$\Rightarrow v = \frac{c}{\sqrt{2}} = \frac{3 \times 10^8}{\sqrt{2}} = 2.12 \times 10^8 \text{ ms}^{-1}$$


Sol 5: Lens formula.

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f} \text{ (for concave lens } f \text{ is } (-) \text{ ve)}$$

$$\text{Now for } u = -f, \frac{1}{v} - \frac{1}{-f} = \frac{1}{-f}$$


$$\frac{1}{v} = -\frac{1}{f} - \frac{1}{f} = -\frac{2}{f} \Rightarrow v = -\frac{f}{2}$$

Image is virtual, diminished and on the same side as object.




The variation of angle of deviation δ with the angle of incidence i of the ray incident on the first refracting surface of the prism is shown in figure. For one angle of incidence it has a minimum value δ_{\min} . At this value the ray passes symmetrically through the prism.

Sol 7: For relaxed eye, intermediate image should lie at first focus of eye piece or $u_e = f_e$



Magnification $M_\infty = \frac{f_o}{f_e}$

Sol 8: Reflection from a spherical surface: Here $n_1 < n_2$. Ray leaves point O and focuses at point I. Snell's Law at point P $n_1 \sin\theta_1 = n_2 \sin\theta_2$ or $n_1 \theta_1 = n_2 \theta_2$ (For small angles)



From geometry of figure

$$\theta_1 = \alpha + \beta, \beta = \theta_2 + \gamma$$

Eliminating θ_1 and θ_2 we get $\beta = \frac{n_1}{n_2}(\alpha + \beta) + \gamma$

Or $n_1\alpha + n_2\gamma = (n_2 - n_1)\beta$ (A)

Now angle at C is $\beta = \frac{S}{R}$ ($S = \text{arc}(PM)$)

Also in paraxial approximation $\alpha = \frac{S}{u}$ and $\gamma = \frac{S}{v}$

Putting these expressions with proper signs, in eqn. A,

we get $\frac{n_1}{-u} + \frac{n_2}{v} = \frac{n_2 - n_1}{R_1}$

Lens Maker's Formula: $\frac{1}{f} = (n-1)\left(\frac{1}{R_1} - \frac{1}{R_2}\right)$

As wavelength of light increases, the refractive index n decreases and from the lens maker formula we see that, as n decreases, f increases.

Sol 9: (i) Magnifying power:


$m = \frac{v}{u} \left(\frac{D}{f_e}\right)$ for normal adjustment

$m = \frac{v}{u} \left(1 + \frac{D}{f_e}\right)$ for final image at D, least distance for clear vision.

(ii) Resolving Power: $R = \frac{1}{\Delta d} = \frac{2\mu \sin\theta}{\lambda}$


$\mu \rightarrow$ Refractive index of the medium between the object and the objective.

$\lambda \rightarrow$ Wavelength of light.



Here we see that

(i) as μ increases, R increases.



(ii) as λ increases, R decreases.

Sol 10: Refracting astronomical telescope: It consists of an objective lens of a large focal length (f_o) and large aperture, also an eye lens of small aperture and focal length.

(i) Magnification when final image is formed at D,

$$\Rightarrow m = -\frac{f_o}{f_e} \left(1 + \frac{f_e}{D}\right) \text{ and length of telescope,}$$

$$L = |f_o| + \frac{f_e D}{f_e + D}$$

Sol 11: (i) Lens formula

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

$$u = +12 \text{ cm, } f = +20 \text{ cm}$$

$$\Rightarrow \frac{1}{v} = \frac{1}{u} + \frac{1}{f} = \frac{1}{12} + \frac{1}{20} = \frac{1}{4} \left(\frac{1}{3} + \frac{1}{5}\right) = \frac{8}{4 \times 15}$$

$$\Rightarrow v = +7.5 \text{ cm}$$

$$(ii) f = -16 \text{ cm} \Rightarrow \frac{1}{v} = \frac{1}{12} - \frac{1}{16} = \frac{4-3}{48} = \frac{1}{48}$$

$$\Rightarrow v = +48 \text{ cm}$$

$$\text{Sol 12: } \frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$\text{Here } R_1 = +30 \text{ cm ; } R_2 = -30 \text{ cm ; } \mu = 1.6$$

$$\Rightarrow \frac{1}{f} = (0.6) \left(\frac{1}{30} \right) \times 2 = \frac{2}{50} = \frac{1}{25} \Rightarrow f = +25 \text{ cm}$$

$$u = 12.5 \text{ cm, } \frac{1}{v} = \frac{1}{f} + \frac{1}{u} = \frac{1}{25} - \frac{1}{12.5} = -\frac{1}{25}$$

$$\Rightarrow v = -25 \text{ cm, } m = \frac{h_2}{h_1} = \frac{v}{u} = \frac{-25}{-12.5} = 2$$

$$\Rightarrow h_2 = 2h_1 = 2 \times 5 \text{ cm} = 10 \text{ cm}$$

Image is virtual and erect, on the same side as the object.

Sol 13: Predominance of bluish colour in a clear sky is due to the phenomena of scattering of light in the atmosphere around earth. If size of the air particles are smaller than the wavelength, the scattering is proportional to $1/\lambda^4$. This is the Rayleigh's law of scattering. The light of short wavelengths are strongly scattered by the air molecules and reach the observer.

Among the shorter wavelengths, the colour blue is present in large proportion in sunlight.

Sol 14: Angle of minimum deviation δ_m and angle of

$$\text{prism } A \text{ are related as, } \mu = \frac{\sin\left(\frac{A + \delta_m}{2}\right)}{\sin\left(\frac{A}{2}\right)}$$


Glass prism of refractive index 1.5 is immersed in a liquid of refractive index 1.3 so the relative refractive index of the prism decreases. $\mu' = \frac{1.5}{1.3} = 1.15$

So as per above equation as A is constant for a prism, as μ decreases, δ_m also decreases.

Sol 15: [Refer question 7 solution]

Sol 16: Consider an object O placed at a distance u from a convex lens as shown in figure. Let its image I after two refractions from spherical surfaces of radii R_1 (positive) and R_2 (negative) be formed at a distance v from the lens. Let v_1 be the distance of image formed by refraction from the refracting surface of radius R_1 . This image acts as an object for the second surface. Using,

$$\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R}$$



$$\text{twice, we have } \frac{\mu_2}{v_1} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R} \quad \dots (i)$$

$$\text{and } \frac{\mu_1}{v} - \frac{\mu_2}{v_1} = \frac{\mu_1 - \mu_2}{-R_2} \quad \dots (ii)$$

Adding Eqs. (i) and (ii) and then simplifying, we get

$$\frac{1}{v} - \frac{1}{u} = \left(\frac{\mu_2}{\mu_1} - 1 \right) \left(\frac{1}{R_1} - \frac{1}{R_2} \right) \quad \dots (iii)$$

This expression relates the image distance v of the image formed by a thin lens to the object distance u and to the thin lens properties (index of refraction and radii of curvature). It is valid only for paraxial rays and

only when the lens thickness is much less than R_1 and R_2 . The focal length f of a thin lens is the image distance that corresponds to an object at infinity. So, putting $u = \infty$ and $v = f$ in the above equation, we have

$$\frac{1}{f} = \left(\frac{\mu_2}{\mu_1} - 1\right) \left(\frac{1}{R_1} - \frac{1}{R_2}\right) \dots (iv)$$

If the refractive index of the material of the lens is μ and it is placed in air, $\mu_2 = \mu$ and $\mu_1 = 1$ so that Eq. (iv) becomes

$$\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2}\right) \dots (v)$$

This is called the lens maker's formula because it can be used to determine the values of R_1 and R_2 that are needed for a given refractive index and a desired focal length f .

Combining Eqs. (iii) and (v), we get

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f} \dots (vi) \text{ which is known as the lens formula.}$$

(i) $P = P_1 + P_2 = 10D - 5D = 5D$

(ii) $f = \frac{1}{P} = \frac{1}{5D} = 0.2m = 20cm$; $m = +2 = \frac{v}{u}$
 $\Rightarrow v = 2u$

For virtual image m is positive

$$\frac{1}{v} = \frac{1}{f} + \frac{1}{u} \Rightarrow \frac{1}{2u} - \frac{1}{u} = \frac{1}{20}$$

$$\Rightarrow -\frac{1}{2u} = \frac{1}{20} \Rightarrow u = -10 \text{ cm}$$

Sol 17:

$$P = P_1 + P_2 = 6 - 2 = 4D \Rightarrow f = \frac{1}{P} = \frac{1}{4D} = 0.25m$$


$$\Rightarrow f = 25cm$$

Sol 18: The speed of light in vacuum is a universal constant denoted by c . When a light wave travels in a transparent material, the speed is decreased by a factor μ , called the refractive index of the material.

$$\mu = \frac{\text{speed of light in vacuum}}{\text{speed of light in the material}}$$

For graph refer figure of question 6.

Sol 19: (a) (i)



(ii) Reflector telescope advantages:


1. Reflector telescopes do not suffer from chromatic aberration because all wavelengths will reflect off the mirror in the same way.
2. Support for the objective mirror is all along the back side so they can be made very BIG.
3. Reflector telescopes are cheaper to make than refractors of the same size.
4. Because light is reflecting off the objective, rather than passing through it, only one side of the reflector telescope's objective needs to be perfect.

(b) $f_1 = +10 \text{ cm}$, $f_2 = -10 \text{ cm}$, $u = -30 \text{ cm}$.

$$\frac{1}{v_1} = \frac{1}{f_1} + \frac{1}{u_1} = \frac{1}{10} - \frac{1}{30} = \frac{2}{30} \Rightarrow v_1 = 15 \text{ cm}$$

So for the concave lens

$$u_2 = +(15 - 5) \text{ cm} = +10 \text{ cm}$$



$$\frac{1}{v_2} = \frac{1}{f_2} + \frac{1}{u_2} = \frac{1}{-10} + \frac{1}{10} = 0 \Rightarrow v_2 = \infty$$


Final image will be formed at infinity.

Sol 20: $\frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2} = \frac{1}{f} - \frac{1}{f} = 0 \Rightarrow F = \infty$

Sol 21: (i) Astronomical Telescope for normal adjustment

It consists of two converging lenses placed coaxially.

The one facing the distant object is called the objective and has a large aperture and a large focal length. The other is called the eyepiece, as the eye is placed close to it. It has a smaller aperture and a smaller focal length. The lenses are fixed in tubes. The eyepiece tube can slide within the objective tube so that the separation between the objective and the eyepiece may be changed.



When the telescope is directed towards a distant object PQ, the objective forms a real image of the object in its focal plane. If the point P is on the principal axis, the image point P' is at the second focus of the objective. The rays coming from Q are focused at Q'. The eyepiece forms a magnified virtual image P'Q'' of P'Q'. This image is finally seen by the eye. In normal adjustment, the position is so adjusted that the final image is formed at infinity. In such a case, the first image P'Q' is formed in the first focal plane of the eyepiece. The eye is least strained to focus this final image. The image can be brought closer by pushing the eyepiece closer to the first image. Maximum angular magnification is produced when the final image is formed at the near point.

$$(ii) m = -\frac{f_o}{f_e} = -\frac{p_e}{p_o} = -\frac{10D}{1D} = -10$$

Here the objective has large focal length and smaller Power.

(b) (i) Figure shows a simplified version of a compound microscope and the ray diagram for image formation. It consists of two converging lenses arranged coaxially. The one facing the object is called the objective and the one close to the eye is called the eyepiece or ocular.

The object is placed at a distance u_o from the objective which is slightly greater than its focal length f_o . A real and inverted image is formed at a distance v_o on the other side of the objective. This image works as the object for the eyepiece. For normal adjustment, the position of the eyepiece is so adjusted that the image formed by the objective falls in the focal plane of the

eyepiece. The final image is then formed at infinity. It is erect with respect to the first image and hence, inverted with respect to the object. The eye is least strained in this adjustment as it has to focus the parallel rays coming to it. The position of the eyepiece can also be adjusted in such a way that the final virtual image is formed at the near point. The angular magnification is increased in this case. The ray diagram in figure refers to this case.

(ii) Magnifying power of a Compound microscope is

$$m = -\frac{v_o}{u_o} \left(\frac{D}{f_e} \right) \rightarrow \text{normal adjustment and}$$

$$m = -\frac{v_o}{u_o} \left(1 + \frac{D}{f_e} \right) \rightarrow \text{final image at } D.$$

Now for large magnification, m is to be large, so f_e should be small and u_o should be small. Now object is placed at a distance u_o from the objective which is slightly greater than its focal length f_o . So for u_o to be small, f_o should also be small.

Sol 22: Refer question 19.(a).(ii)

Sol 23: When two plane mirrors are placed at an angle θ to each other, the object is kept between them, then the numbers of images observed is $n = \frac{360^\circ}{\theta}$. If n is even then number of image is $(n-1)$.

$$\text{So here } \frac{360^\circ}{\theta} = \frac{360^\circ}{60} = 6 \Rightarrow n = 6 - 1 = 5$$

(b) At each reflection some of the light energy is lost due to absorption at the mirror surface. So the intensity of the reflected ray goes on decreasing at multiple reflections due to parallel mirrors.

Sol 24: For compound microscope $m = -\frac{v_o}{u_o} \left(\frac{D}{f_e} \right)$ (normal adjustment)

$$-20 = \left(-\frac{v_o}{u_o} \right) \left(\frac{25 \text{ cm}}{3 \text{ cm}} \right) \Rightarrow -\frac{v_o}{u_o} = \frac{-60}{25} = \frac{-12}{5} = -2.4$$

For final image at least distance:

$$m = -\frac{v_o}{u_o} \left(1 + \frac{D}{f_e} \right) \Rightarrow -20 = -\frac{v_o}{u_o} \left(1 + \frac{25}{3} \right)$$

$$\Rightarrow -\frac{v_o}{u_o} = \frac{-20}{28/3} = \frac{-5 \times 3}{7} \Rightarrow -\frac{v_o}{u_o} = -2.14$$

Sol 25: Astronomical telescope $m = \frac{-f_o}{f_e} \rightarrow$ normal adjustment

$$\Rightarrow -8 = -\frac{f_o}{f_e} \Rightarrow f_o = 8f_e$$

$$L = f_o + f_e = 45 \text{ cm} \Rightarrow 8f_e + f_e = 45 \text{ cm}$$


$$\Rightarrow f_e = 5 \text{ cm} \text{ and } f_o = 40 \text{ cm}$$

Exercise 2

Single Correct Choice Type

Sol 1: (A)

$$\delta_1 = 180^\circ - 2\theta; \delta_2 = 180^\circ - 2\theta' = 70^\circ + 90^\circ - \theta'$$




$$\Rightarrow \theta' = 20^\circ$$

$$\alpha = 180^\circ - 70^\circ - (90^\circ - \theta') = 180^\circ - 70^\circ - 70^\circ$$

$$\alpha = 40^\circ \Rightarrow \theta = 90^\circ - \alpha = 50^\circ$$

Sol 2: (B) With respect to mirror1 the object is going away from mirror. So first image will also more away w.r.t mirror 1 with same speed v . So with respect to object O the image speed is



$$v_{\text{image1}} = v_{\text{image1,mirror1}} + v_{\text{mirror1,obj}} = v + v = 2v$$

Now this image becomes object for mirror 2. With respect to mirror 2 the image is going away towards


left with speed $3v$. So its second image will more away towards right with speed $3v$ w.r.t mirror 2. Hence speed w.r.t O will be $3v+v=4v$

Hence for n th image $v_{\text{image}} = 2nv$

Sol 3: (A) $\tan \theta = \frac{h}{y} = \frac{3h}{10h + y} \lim_{x \rightarrow \infty}$

$$\Rightarrow 10h + y = 3y \Rightarrow y = 5h$$

For general case




$$\tan \theta = \frac{3h}{x + y} = \frac{h}{y} \Rightarrow 3y = x + y \Rightarrow 2y = x$$

$$\Rightarrow 2 \cdot \frac{dy}{dt} = \frac{dx}{dt} \Rightarrow \frac{dy}{dt} = \frac{1}{2} \cdot v$$

Sol 4: (C) A is head and E is feet of man. C is the eye. The mirror can be placed anywhere between the centre line BF (of AC) and DG (of CE) to get full image from head to feet.

So here $CE = 1.4$ m. So DE should be 0.7 m. But mirror is 0.8 m from ground so feet will not be visible. The upper edge of mirror is at height $(0.8 + 0.75)$ m equal to 1.55 m which is more than BE.



$$BE = BC + CE = \frac{AC}{2} + CE = \frac{0.1}{2} + 1.4$$

$$= 0.05 + 1.4 = 1.45 \text{ m}$$

So head will be visible.

Sol 5: (B) Rays from S going right from the normal will not reach the bottom horizontal mirror as they will hit the inclined mirror and get deviated. So C will see only the image formed by inclined mirror.

Sol 6: (A) Object O moves towards M_1 so image 1 due to M_1 will move towards left i.e. towards M_2 .

For M_2 we have formula for speed of image as

$$\frac{dv}{dt} = -\left(\frac{v^2}{u^2}\right)\frac{du}{dt}$$

So negative sign means final image

2 due to M_2 will move opposite to image 1 i.e. towards right.




Sol 7: (D) Intensity incident

$$= \frac{P\theta^2}{4 \times (\text{area on which light is incident})}$$

When the mirror is not present, light is reaching the screen up to height h. Maximum area on which light is incident = πh^2

$$\text{Intensity} = \frac{P\theta^2}{4\pi h^2} \text{ and}$$

$$\tan \theta = \frac{h}{60} \Rightarrow \frac{Ph^2}{4\pi h^2 \times (60)^2} = \frac{P}{4\pi \times (60)^2}$$




When the mirror is present

$$\text{Intensity} = \frac{P\theta_1^2}{4\pi h_1^2} \text{ and } \tan \theta = \frac{h}{20} \Rightarrow \frac{P}{4\pi \times (20)^2}$$


$$\text{Ratio} = \frac{\text{Intensity when mirror present}}{\text{Intensity when mirror object}}$$

$$= \frac{\frac{P}{4\pi \times (20)^2} + \frac{P}{4\pi \times (60)^2}}{\frac{P}{4\pi \times (20)^2}} = \frac{10}{1}$$

Sol 8: (D) As a result of water the apparent height of source will be beyond C, at C' . $OC' = R\mu$. So its image I' will not be formed at C but it will be formed between C and focus F of the mirror. But again in the return path of rays they will be again refracted at water to air boundary and final image I will be further shifted downwards towards O.




Sol 9: (A) The slab will cause a lateral shift in the incident rays as well as in the reflected rays from the circular mirror MM' . Now the angle of emergence θ_1 will be equal to the angle of incidence in case of a slab.



The rays reaching the edge M of the circular mirror after passing through the glass slab will be leaving the source O at a greater angle (θ_1) with the normal as compared to the angle (θ) when there is no slab. But due to symmetry of incident and reflected rays, the reflected rays from the edge M, after passing through the slab will reach the some point Q on the ground where they would have reached when there was no slab.

Here we have, $OQ=QP$, both without and with slab, between source O and mirror MM' .

Sol 10: (A)



$$90 - \theta \geq \theta_c = \sin^{-1} \left(\frac{6/5}{3/2} \right);$$

$$90 - \theta \geq \sin^{-1} \left(\frac{4}{5} \right);$$


$$90 - \theta \geq 53^\circ \Rightarrow \theta \leq 37^\circ$$

Sol 11: (C)

$$h = 4\text{m}; \theta_c = \sin^{-1} \left(\frac{1}{\mu} \right)$$

$$= \sin^{-1} \left(\frac{1}{5/3} \right) = \sin^{-1} \left(\frac{3}{5} \right) = 37^\circ$$


$$\tan \theta_c = \frac{r}{h} \Rightarrow r = h \tan 37^\circ = \frac{3h}{4} = \frac{3 \times 4\text{m}}{4}$$



$$\Rightarrow \text{diameter} = 2r = 6\text{m}$$

Sol 12: (D) $\theta_c = \sin^{-1} \frac{1}{\mu} = \sin^{-1} \left(\frac{3}{4} \right) = 48.59^\circ$

Solid angle subtended at source of light O by the circular area of radius r is



$$\Omega = 2\pi(1 - \cos \theta_c)$$

If total intensity is I then, intensity per unit solid angle is

$$\frac{I}{4\pi}. \text{ So intensity through the circular area is,}$$

$$I' = \frac{I}{4\pi} \cdot 2\pi(1 - \cos \theta_c)$$

$$\Rightarrow \frac{I'}{I} = \frac{(1 - \cos 48.59)}{2} \Rightarrow \frac{I'}{I} = 16.9\%$$

Sol 13: (B) $\frac{1}{v_1} = \frac{1}{f} + \frac{1}{(-u_1)} = \frac{1}{f} - \frac{1}{u_1} = \frac{u_1 - f}{fu_1}$

$$v_1 = \frac{f u_1}{u_1 - f} \text{ and } v_2 = \frac{f u_2}{u_2 - f}$$

$$\text{Now } m_1 = \frac{v}{u} = \frac{v_1}{-u_1} = \frac{-f}{u_1 - f}$$

$$\text{and } m_2 = \frac{-f}{u_2 - f}. \text{ also } |m_1| = |m_2|$$

Now m_1 is negative (real image) and m_2 is positive (virtual image). So we have,

$$\frac{f}{u_1 - f} = \frac{-f}{u_2 - f} \Rightarrow u_2 - f = -u_1 + f$$

$$\Rightarrow u_1 + u_2 = 2f \Rightarrow f = \frac{u_1 + u_2}{2}$$


Sol 14: (B) The image formed by first lens will lie at its second lens focus. This image will act as an object for the second lens. For the rays to become parallel after passing through the second lens, the object for second lens should lie on its first focus. Thus the distance between the two lenses will be equal to sum of their focal lengths.

$$D = f_1 + f_2 = 20\text{cm} + 10\text{cm} = 30\text{cm}$$

Sol 15: (B) Image formed by lens be at distance v_1 from

$$\text{lens. } \frac{1}{v_1} = \frac{1}{f_1} + \frac{1}{u_1} = \frac{1}{10} + \frac{1}{-15} = \frac{3-2}{30} = \frac{1}{30}$$

$v_1 = 30$ cm from lens.



For convex lens, $u_2 = +(30 - 10)$ cm = +20 cm


$v_2 = +20$ cm, because rays retrace their path after reflection.

$$\frac{1}{v_2} + \frac{1}{u_2} = \frac{1}{f_2} \Rightarrow \frac{1}{f_2} = \frac{1}{20} + \frac{1}{20} = \frac{1}{10} \Rightarrow f_2 = +10 \text{ cm}$$

Sol 16: (C) For first lens, convergent ray becomes parallel to principal axis after refraction.

So $f_1 = +5$ cm.

For second lens, ray parallel to principal axis becomes convergent and parallel to incident ray.



So focal length of second lens will be x as shown in figure.

$$\tan \theta = \frac{h}{5} = \frac{h}{x} \Rightarrow x = 5 \text{ cm} \Rightarrow f_2 = x = 5 \text{ cm}$$

Sol 17: (D) Let us work in the frame of reference attached to the lens. Lens formula: $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$

Differentiating w.r.t. time,

$$-v^{-2} \frac{dv}{dt} - (-u^{-2}) \frac{du}{dt} = 0; \text{ (f is constant)}$$

$$\Rightarrow \frac{dv}{dt} = \left(\frac{v^2}{u^2} \right) \frac{du}{dt}$$

Initially when $u=f$, $v \rightarrow \infty$ so speed image is very large and finally when $u \rightarrow \infty$, $v \rightarrow f$ and the speed of image is very low (nearly zero). With respect to lens, as object moves left, the image also moves left.

Speed of image with respect to object is

$$v_{I,O} = v_{I,L} + v_{L,O}$$

$$v_{I,O} = \left(\frac{dv}{dt} \right)_{\text{(towards left)}} + (-v)_{\text{(towards right)}}$$

Sol 18: (D) At first refracting surface we have, $\sin i_1 = \mu \sin r_1$. So as i_1 decreases, r_1 also decreases.

Now for prism $r_1 + r_2 = A$ (constant). So as r_1 decreases, r_2 increases. At the second refracting surface we have, $\mu \sin r_2 = \sin i_2$. So as r_2 increases, i_2 also increases. So out of all choices D is most appropriate as amount of increase in i_2 should be less than amount of decrease in i_1 .

Sol 19: (C) For prism with refracting angle A , we have

$$\mu = \frac{\sin \left(\frac{A + \delta_m}{2} \right)}{\sin \frac{A}{2}} \Rightarrow \sqrt{\frac{3}{2}} = \frac{\sin (90^\circ + \delta_m) / 2}{\sin 45^\circ}$$

$$\Rightarrow \sqrt{\frac{3}{2}} = \sin \left(\frac{90^\circ + \delta_m}{2} \right) \Rightarrow \frac{90^\circ + \delta_m}{2} = 60^\circ$$

$$\Rightarrow \delta_m = 120^\circ - 90^\circ = 30^\circ$$

Sol 20: (B) At min deviation $i_m = \frac{A + \delta_m}{2} = \frac{A + 38^\circ}{2}$


Now $\delta = (i_1 + i_2) - A$; Here $i_1 = 42^\circ$

and $i_2 = 62^\circ$, $\delta = 44^\circ$

$$\Rightarrow 44^\circ = (42^\circ + 62^\circ) - A \Rightarrow A = 104^\circ - 44^\circ = 60^\circ$$

$$\Rightarrow i_m = \frac{60^\circ + 38^\circ}{2} = 49^\circ$$

Sol 21: (C)



$$\delta = (\mu - 1)A; \mu = 1.5$$


$$\delta = 0.5 \times 5^\circ = 2.5^\circ \Rightarrow \delta = \frac{2.5}{180} \pi = \frac{5}{360} \pi = \frac{\pi}{72}$$

$$OP \approx OM \approx IM = 10 \text{ cm}; \Delta x = \delta \times (OM) = 10 \delta \text{ cm}$$

$$\Delta x = \frac{\pi}{72} \times 10 \text{ cm} = \frac{5\pi}{36} \text{ cm}$$

Sol 22: (A) At second refracting surface

$$\begin{aligned} \mu \sin r_2 &= 1 \sin 90 \\ \Rightarrow \sin r_2 &= \frac{1}{\mu} = \frac{1}{\sqrt{2}} \\ \Rightarrow r_2 &= 45^\circ \\ \Rightarrow r_1 &= A - r_2 \\ &= 60^\circ - 45^\circ = 15^\circ \end{aligned}$$



$$\begin{aligned} \Rightarrow \text{At first refracting surface, } \sin i_1 &= \mu \sin r_1 \\ \Rightarrow \sin i_1 &= \sqrt{2} \cdot \sin 15^\circ = \sqrt{2} \cdot \frac{\sqrt{3}-1}{2\sqrt{2}} = \frac{\sqrt{3}-1}{2} \\ \Rightarrow i_1 &= \sin^{-1} \left(\frac{\sqrt{3}-1}{2} \right) \end{aligned}$$

Previous Years' Questions

Sol 1: $\frac{1}{v} - \frac{1}{u} = \frac{1}{f} = \text{constant}$

Sol 2: (A) An experiment is performed to find the refractive index of glass using a travelling microscope. In this experiment, distances are measured by a vernier scale provided on the microscope.

Sol 3: (A) \rightarrow since $\mu_1 < \mu_2$, the ray of light will bend towards normal after first refraction.

(B) $\rightarrow \mu_1 > \mu_2$, the ray of light will bend away from the normal after first refraction.


(C) $\rightarrow \mu_2 = \mu_3$ means in second refraction there will be no change in the path of ray of light.

(D) \rightarrow Since $\mu_2 > \mu_3$, ray of light will bend away from the normal after second refraction.

Therefore the correct options are as under.

- (A) \rightarrow p, r
- (B) \rightarrow q, s, t
- (C) \rightarrow p, r, t
- (D) \rightarrow q, s

Sol 4: (A, B, C) Using Snell's law



$$\sin^{-1} \frac{1}{\sqrt{3}} < \sin^{-1} \frac{1}{\sqrt{2}}$$

Net deviation is 90°

Sol 5:

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

or $\frac{u}{v} - 1 = \frac{u}{f}$ or $\frac{u}{v} = \left(\frac{u+f}{f} \right)$

$$\therefore m = \frac{v}{u} = \left(\frac{f}{u+f} \right)$$

$$\frac{m_{25}}{m_{30}} = \frac{\left(\frac{20}{-25+20} \right)}{\left(\frac{20}{-50+20} \right)} = 6$$

\therefore Answer is 6.

Sol 6: (D) Case I: $u = -240 \text{ cm}$, $v = 12$, by lens formula

$$\frac{1}{f} = \frac{7}{80}$$

Case II: $v = 12 - \frac{1}{3} = \frac{35}{3}$

(normal shift) $= 1 - \frac{2}{3} = \frac{1}{3}$

$$f = \frac{7}{80}$$

$$u = 5.6$$

Sol 7: (C) $L.C = \frac{1}{60}$

Total Reading = $585 + \frac{9}{60} = 58.65$

Sol 8: (B) As intensity is maximum at axis.

$\therefore \mu$ will be maximum and speed will be minimum on the axis of the beam.

\therefore Beam will converge.

Sol 9: (D) For a parallel cylindrical beam, wave front will be planar.

Sol 10: (D) Case I: $u = -240$ cm, $v = 12$, by Lens formula

$$\frac{1}{f} = \frac{7}{80}$$

Case II: $v = 12 - \frac{1}{3} = \frac{35}{3}$

$$\left(\text{Normal shift} = 1 - \frac{2}{3} = \frac{1}{3} \right)$$

$$f = \frac{7}{80}$$


$$u = 5.6$$

Sol 11: (B) Self-explanatory

Sol 12: (B)

$$R^2 = d^2 + (R - t)^2$$

$$R^2 - d^2 = R^2 \left\{ 1 - \frac{t}{R} \right\}^2$$



$$1 - \frac{d^2}{R^2} = 1 - \frac{2t}{R}$$

$$R = \frac{(3)^2}{2 \times (0.3)} = \frac{90}{6} = 15 \text{ cm}$$

$$\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$\frac{1}{f} = \left(\frac{3}{2} - 1 \right) \left(\frac{1}{15} \right)$$

$$f = 30 \text{ cm}$$

Sol 13: (D) $\frac{f_m}{f} = \frac{(\mu - 1)}{\left(\frac{\mu}{\mu_m} - 1 \right)}$

$$\Rightarrow \frac{f_1}{f} = \frac{\left(\frac{3}{2} - 1 \right)}{\left(\frac{3/2}{4/3} - 1 \right)} = 4$$

$$\Rightarrow f_1 = 4f$$

$$\frac{f_2}{f} = \frac{\left(\frac{3}{2} - 1 \right)}{\left(\frac{3/2}{5/3} - 1 \right)} = -5$$

$$\Rightarrow f_2 < 0$$

Sol 14: (D) As frequency of visible light increases refractive index increases. With the increase of refractive index critical angle decreases. So that light having frequency greater than green will get total internal reflection and the light having frequency less than green will pass to air.

Sol 15: (D) At face AB,

$$\sin \theta = \mu \sin r$$

At face AC $r' < \theta_c$


$$A - r < \sin^{-1} \frac{1}{\mu}$$

$$\therefore r > A - \sin^{-1} \frac{1}{\mu}$$

$$\therefore \sin r > \sin \left(A - \sin^{-1} \frac{1}{\mu} \right)$$

$$\frac{\sin \theta}{\mu} > \sin \left(A - \sin^{-1} \frac{1}{\mu} \right)$$

$$\theta > \sin^{-1} \left[\mu \sin \left(A - \sin^{-1} \frac{1}{\mu} \right) \right]$$



Sol 16: (A) $\theta = 1.22 \frac{\lambda}{D}$

Minimum separation = $(25 \times 10^{-2}) \theta = 30 \mu\text{m}$

Sol 17: (D) $\delta = i + e - A \Rightarrow A = 74^\circ$

$$\mu = \frac{\sin\left(\frac{A + \delta_{\min}}{2}\right)}{\sin\left(\frac{A}{2}\right)} = \frac{5}{3} \sin\left(37^\circ + \frac{\delta_{\min}}{2}\right)$$

μ_{\max} can be $\frac{5}{3}$, so μ will be less than $\frac{5}{3}$

Since δ_{\min} will be less than 40° , so

$$\mu < \frac{5}{3} \sin 57^\circ < \frac{5}{3} \sin 60^\circ \Rightarrow \mu < 1.446$$


So the nearest possible value of μ should be 1.5

JEE Advanced/Boards

Exercise 1


Sol 1: $OB = OP \sin 15^\circ = \sqrt{2} \left(\frac{\sqrt{3}-1}{2\sqrt{2}} \right) = \frac{\sqrt{3}-1}{2}$

Number of image



$$n = \frac{360}{\theta} - 1 = \frac{360}{\theta} - 1 = 5$$

Sol 2:



By similar triangles

$\Delta BCD \sim \Delta BFE$; so $EF = 3 \times CD$

Because $BF = 3 \times BC$


$\Rightarrow EF = 120 \text{ cm} \Rightarrow EP = 160 \text{ cm}$

\therefore Minimum height of eye is 160 cm.

And similarly maximum height will be

$$E'P = 80 + 3 \times 80 = 320 \text{ cm}$$


Sol 3: By property of similar triangles,



$$\Delta MAB \sim \Delta MCD; \frac{x}{20 \text{ cm}} = \frac{300 \text{ cm}}{y}$$

$$y + 20 = 100 \text{ cm}; \Rightarrow y = 80 \text{ cm}$$

$$\Rightarrow x = \frac{20 \times 300 \text{ cm}}{80} = 75 \text{ cm}$$

Sol 4:


$$\frac{1}{v} + \frac{1}{u} = \frac{1}{f}; f = -20 \text{ cm}, u = -10 \text{ cm}$$


$$\frac{1}{v} - \frac{1}{10} = -\frac{1}{20} \Rightarrow \frac{1}{v} = \frac{1}{10} - \frac{1}{20} = \frac{2-1}{20} = \frac{1}{20} \Rightarrow v = 20$$

$$M = \frac{v}{u} = -\frac{20}{(-10)} = 2$$

Image will be erect with respect to the axis of each mirror. Distance between images is 2 cm.

Sol 5: OS = 15 m; OC = 20 m

Height of ball after 4 s is,



$$H = 15 + 20 \times 4 - \frac{1}{2} \times 10 \times 16 = 15 + 80 - 80 = 15 \text{ m}$$

 Speed of ball after 4 s is, $s_{\text{ball}} = 20 - 10 \times 4$

$$\Rightarrow s_{\text{ball}} = 20 - 40 = -20 \text{ ms}^{-1}$$

So ball is moving downward with speed of 20 ms^{-1} and is at height 15 m above the mirror.

$$\text{So } u = -15 \text{ m, and } f = \frac{20}{-2} \text{ m} = -10 \text{ m}$$

Mirror formula $\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$ gives

$$\frac{1}{v} = \frac{1}{-10} - \frac{1}{(-15)} = \frac{1}{15} - \frac{1}{-10} = \frac{2-3}{30} = \frac{-1}{30}$$

$$\Rightarrow v = -30 \text{ cm}$$

Relation between speed of image and speed of object for lens is

$$\frac{dv}{dt} = \left(\frac{v^2}{u^2} \right) \cdot \frac{du}{dt} \Rightarrow \frac{dv}{dt} = \left(\frac{(-30)^2}{(-15)^2} \right) \cdot 20 \text{ m s}^{-1} \text{ (downward)}$$

$$\frac{dv}{dt} = (4) \times 20 \text{ ms}^{-1} = 80 \text{ m s}^{-1} \text{ (downward)}$$


Sol 6: Refer theory

Sol 7: Apply Snell's law:

$$1 \times \sin \theta = \frac{4}{3} \sin(90 - \phi)$$

$$1 \times \frac{d}{\sqrt{30+d^2}} = \frac{4}{3} \times \frac{d \times 3}{\sqrt{1024+d^2}}$$

$$\sqrt{1024+d^2} = 4\sqrt{36+d^2}$$



On squaring both sides,

$$1024 + 9d^2 = 16(36 + d^2); 1024 + 9d^2 = 576 + 16d^2$$

$$7d^2 = 448; d^2 = \frac{448}{7} = 64$$


$$d = 8 \text{ feet, width} = 2d = 16 \text{ feet}$$

Sol 8: At any point by Snell's law

$$1 \sin 90 = n(y) \cdot \sin(90^\circ - \theta);$$

$$1 = (kg^{3/2} + 1)^{1/2} \cos \theta$$

Here angle of incidence at B(x, y) is $i = 90^\circ - \theta$



$$\cos \theta = \frac{1}{(ky^{3/2} + 1)^{1/2}}$$

$$\Rightarrow \tan \theta = \frac{dy}{dx} = \frac{1}{2} = \frac{\sqrt{ky^{3/2} + 1} - 1}{1} = \sqrt{ky^{3/2}}$$

(i) Now as

$$i = 90^\circ - \theta \Rightarrow \theta = 90^\circ - i \text{ So } \frac{dy}{dx} = \tan \theta = \cot i$$

(ii) Initial angle at air is 90° and $n=1$. At point B(x, y) angle of incidence is i .

So we have by Snell's law

$$1 \cdot \sin 90^\circ = n \sin i$$

$$\Rightarrow n \sin i = 1$$

(iii) Now, $\frac{dy}{dx} = \sqrt{k} y^{3/4}$ [From (i)]


$$\Rightarrow \int_0^y \frac{dy}{y^{3/4}} = \int_0^x \sqrt{k} dx \Rightarrow \frac{y^{+1/4}}{+1/4} = \sqrt{k} x$$

$$\Rightarrow y^{+1/4} = \frac{\sqrt{k} x}{4} \Rightarrow y = k^2 \left(\frac{x}{4} \right)^4 \quad \dots(ii)$$

(iv) For the point P, we have $y=1.0$ m $k=1$

$$\Rightarrow y = k^2 \left(\frac{x}{4} \right)^4 \text{ gives } 1 = \left(\frac{x}{4} \right)^4 \Rightarrow x = 4.0 \text{ m}$$

(v) At P, we have




$$\frac{dy}{dx} = \tan \theta = 1 \Rightarrow \theta = 45^\circ \text{ [From (i) put } y = 1]$$

$$n = (1+1)^{1/2} = \sqrt{2} \Rightarrow \frac{\sin 45^\circ}{\sin r} = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \sin r = 1 \Rightarrow r = 90^\circ$$

\Rightarrow Ray of light becomes parallel to x axis.

Sol 9: At first refracting surface the rays will pass undeviated. At the second refracting surface the rays are refracting from denser to rarer medium and hence suffer Total Internal Refraction if $i > \theta_c$



$$\theta_c = \sin^{-1} \left(\frac{1}{\mu} \right) \Rightarrow \theta_c = \sin^{-1} \left(\frac{\sqrt{3}}{2} \right) = 60^\circ$$

From right $\triangle BCD$, $\angle BCD = 90^\circ - \theta_c$

$$\text{From } \triangle ABD, h = R \sin \theta_c = R \sin 60^\circ = R \frac{\sqrt{3}}{2}$$

$$DC = h \cot (90^\circ - \theta_c) = h \cot 30^\circ = R \frac{\sqrt{3}}{2} \cdot \sqrt{3} = \frac{3R}{2}$$


$$AP = R, AD = R \cos \theta_c = \frac{R}{2}$$

$$AC = AD + DC = \frac{R}{2} + \frac{3R}{2} = 2R;$$

$$x = PC = AC - AP = 2R - R = R = 5 \text{ cm}$$

$$\Rightarrow x = 5 \text{ m}$$

Sol 10:




$$\frac{\sin i}{\sin r} = \mu \Rightarrow \frac{\sin 37^\circ}{\sin r} = \mu = \frac{4}{3} \Rightarrow \sin r = \frac{3}{\sqrt{h^2 + 9}} = \frac{3}{5}$$

$$\Rightarrow \frac{1}{\sqrt{h^2 + 9}} = \frac{3}{30} \Rightarrow 400 = 9(h^2 + 9) \Rightarrow h^2 = \frac{400}{9} - 9$$

$$\Rightarrow h = \sqrt{\frac{319}{9}} = 5.95 \text{ m}$$


Sol 11: $\frac{\sin i}{\sin r} = \mu$ All the ray in the beam are deviated by same angle so width of beam will not change after it goes over to air.



Sol 12: $\delta = (\mu - 1)A = 1^\circ 15' = 1.25^\circ = (\mu - 1) A$... (i)

Now for prism $r_1 + r_2 = A$. Here $r_1 = 0$,
 $r_2 = r$
 $\Rightarrow r = A$... (ii)

For reflection of reflected ray at first face of prism



$$\mu \sin 2r = \sin 6.5 = \frac{6.5}{180} \pi$$

$$\Rightarrow 2r\mu = \frac{6.5\pi}{180} \Rightarrow r\mu = \frac{6.5\pi}{360}$$
 ... (iii)

From (i), (ii) & (iii) we get

$$\frac{1.25 \pi}{180} = (\mu - 1)A = \mu A - A$$
 ... (iv)

and $\frac{6.5 \pi}{360} = \mu A$... (v)

Subtract (iv) from (v) to get

$$\frac{(6.5 - 2.5)\pi}{360} = A \Rightarrow A = \frac{\pi}{90} \text{ rad} = 2^\circ$$

and put value of A in (v) to get


$$\mu = \frac{6.5 \pi}{360} \times \frac{90}{\pi} \Rightarrow \mu = 1.625$$

Sol 13: For a given incident ray, if the mirror is rotated through an angle θ , then the reflected ray turns through an angle of 2θ . So if angular speed of mirror is ω then the angular speed by which the reflected ray is rotated is 2ω .

$$\omega_{\text{refl}} = 2 \times \omega = 2 \times \frac{9 \text{ rev}}{\pi \text{ sec}} = \frac{18}{\pi} \times 2\pi \frac{\text{rad}}{\text{sec}}$$

$$\omega_{\text{refl}} = 36 \text{ rad s}^{-1}$$

When



$i = 37^\circ$, we have

$$OB = \frac{OA}{\cos(90-i)}$$

$$= \frac{OA}{\sin i} = \frac{10}{\sin 37} = \frac{10}{3/5} \text{ m}$$


$$\Rightarrow OB = \frac{50}{3} \text{ m}$$

$$\Rightarrow \text{speed } v_{\text{refl}} = \omega_{\text{refl}} \times [OB]$$

$$\Rightarrow v_{\text{refl}} = 36 \times \frac{50}{3} \text{ m s}^{-1} = 600 \text{ m s}^{-1}$$

$$v_B = \frac{v_{\text{refl}}}{\cos(90-i)} = \frac{v_{\text{refl}}}{\sin i} = \frac{600}{3/5}$$

$$= 200 \times 5 = 1000 \text{ m/s.}$$



Sol 14: $1 \sin 90 = \mu_1 \sin r_1 \dots\dots(i)$

$\mu_1 \sin r_2 = \mu_2 \sin r_3 \dots\dots(ii)$

$\mu_2 \sin r_4 = \mu_3 \sin r_5 \dots\dots(iii)$

$\mu_3 \sin r_6 = \mu_4 \sin r_7 \dots\dots(iv)$

$\mu_4 \sin r_8 = \mu_5 \sin r_9 \dots\dots(v)$

$\mu_5 \sin r_{10} = 1 \sin 90 \dots\dots(vi)$

Also as all prisms are isosceles right angled prisms, we have

$r_1 + r_2 = r_3 + r_4 = r_5 + r_6 = r_7 + r_8 = r_9 + r_{10} = 90^\circ \dots\dots(viii)$

From equations (i), (ii) and (viii) we get,

$1 = \mu_1 \sin r_1 ; \mu_2 \sin r_3 = \mu_1 \cos r_1$

$$\Rightarrow \mu_1^2 = 1 + \mu_2^2 \sin^2 r_3 \dots\dots(A)$$

From (iii) $\mu_3 \sin r_5 = \mu_2 \cos r_3 \dots\dots(B)$

(A) and (B) gives

$\mu_1^2 + \mu_3^2 \sin^2 r_5 = 1 + \mu_2^2 \dots\dots(C)$

From (iv)

$\mu_3 \cos r_5 = \mu_4 \sin r_7 \dots\dots(D)$

(C) and (D) gives

$\mu_1^2 + \mu_3^2 = 1 + \mu_2^2 + \mu_4^2 \sin^2 r_7 \dots\dots(E)$

From (v) $\mu_5 \sin r_9 = \mu_4 \cos r_7 \dots\dots(F)$

(E) and (F) gives

$\mu_1^2 + \mu_3^2 + \mu_5^2 \sin^2 r_9 = 1 + \mu_2^2 + \mu_4^2 \dots\dots(G)$


From (vi) $\mu_5 \cos r_9 = 1 \dots\dots(H)$

(G) and (H) gives

$\Rightarrow \mu_1^2 + \mu_3^2 + \mu_5^2 = 2 + \mu_2^2 + \mu_4^2$

Sol 15: Snell's law at spherical surface for the first ray

$\mu \sin i_1 = \sin r_1$



$\Rightarrow 1.6 \left(\frac{h_1}{R} \right) = \frac{h_1}{\sqrt{h_1^2 + x_1^2}} \Rightarrow h_1^2 + x_1^2 = \frac{R^2}{2.56}$

$\Rightarrow 0.5^2 + x_2^2 = \frac{20^2}{2.56} \Rightarrow x_2 = 12.49 \text{ m}$

Similarly for second ray

$\Rightarrow 1.6 \left(\frac{h_2}{R} \right) = \frac{h_2}{\sqrt{h_2^2 + x_2^2}} \Rightarrow h_2^2 + x_2^2 = \frac{R^2}{2.56}$

$\Rightarrow x_2^2 = \frac{20^2}{2.56} - 12^2 \Rightarrow x_2 = 3.5 \text{ m}$

$\Rightarrow \Delta x_2 = x_1 - x_2 = 9 \text{ m}$


Sol 16: (a) For total internal reflection at the concrete-air interface we have critical angle

$\frac{\sin \theta_c}{\sin 90} = \frac{\mu_2}{\mu_1} = \frac{v_1}{v_2} = \frac{340 \text{ m/s}}{1700 \text{ m/s}}$

$\Rightarrow \sin \theta_c = \frac{1}{5} \Rightarrow \theta_c = \sin^{-1} \left(\frac{1}{5} \right)$

(b) The concrete is a rarer medium for sound because the speed of sound is higher in concrete, while air will be denser medium for sound as the speed of sound is lower in air. So for TIR, sound must travel in air which is denser medium in this case.

Sol 17: The outer most ray of the beam, ray 1, will be tangential to the circular surface of rod at point P and hence angle of incidence is 90° , hence greater than critical angle, and hence will travel tangentially at all points of the circular portion from P to P'.



The inner most ray of the beam, ray 2, will be incident on the inner circular surface at angle i .

From the geometry of figure we see that,

$$CQ = R; CA = R + d$$

$$R = (R + d) \cos(90 - i) \Rightarrow R = (R + d) \sin i$$

Here angle i will be the least of all angles of incidence of ray 2 during its path inside the circular rod. So, if i is greater than critical angle then ray 2 will suffer TIR at all point in circular rod.

$$\sin i > \frac{1}{\mu} \Rightarrow \frac{R}{R + d} > \frac{1}{\mu} \Rightarrow \frac{1}{1 + \frac{d}{R}} > \frac{1}{\mu} \Rightarrow 1 + \frac{d}{R} < \mu$$

$$\Rightarrow \frac{d}{R} < (\mu - 1) \Rightarrow \left(\frac{d}{R}\right)_{\max} = (\mu - 1) = 1.5 - 1$$

$$\Rightarrow \left(\frac{d}{R}\right)_{\max} = \frac{1}{2}$$

Sol 18: In a prime $r_1 + r_2 = A$;

Here $r_1 = r, r_2 = 0^\circ, A = 30^\circ \Rightarrow r = 30^\circ$ (i)

At first refracting surface

$$\frac{\sin i}{\sin r} = \mu = \sqrt{2} \Rightarrow \sin i = (\sin 30^\circ)\sqrt{2} = \sqrt{2} \left(\frac{1}{2}\right) = \frac{1}{\sqrt{2}}$$

$$\Rightarrow i = 45^\circ$$

Sol 19: Angle of deviation is gives as $\delta = (i_1 + i_2) - A$

Hence $i_2 - i_1 = 23^\circ, A = 60^\circ, \delta = 23^\circ$


$$\Rightarrow 23^\circ = i_1 + i_2 - 60$$

$$\Rightarrow i_1 + i_2 = 83^\circ \quad \dots (i)$$

$$\Rightarrow i_2 - i_1 = 23^\circ \quad \dots (ii)$$

From (i) and (ii) we get $i_1 = 30^\circ, i_2 = 53^\circ$

Snell's law at first refracting surface.




$$\sin i_1 = \mu \sin r_1 \Rightarrow \frac{1}{2} = \mu \sin r_1 \Rightarrow \sin r_1 = \frac{1}{2\mu}$$

Snell's law at second refracting surface.

$$\mu \sin r_2 = \sin i_2 \Rightarrow \sin r_2 = \frac{1}{\mu} \cdot \frac{4}{5} = \frac{4}{5\mu}$$

Now $r_1 + r_2 = 60^\circ$



$$\Rightarrow \sin(r_1 + r_2) = \frac{\sqrt{3}}{2} \Rightarrow \sin r_1 \cos r_2 + \cos r_1 \sin r_2 = \frac{\sqrt{3}}{2} \Rightarrow \mu$$

$$\Rightarrow \frac{1}{2\mu} \cdot \sqrt{1 - \frac{16}{25\mu^2}} + \sqrt{1 - \frac{1}{4\mu^2}} \cdot \frac{4}{5\mu} = \frac{\sqrt{3}}{2}$$

$$= \frac{\sqrt{43}}{5}$$

Sol 20: Deviation suffered by the transmitted ray is

$$\delta_1 = (60^\circ - r_1) + (60^\circ - r_1)$$

$$\delta_1 = 120 - 2r_1 \quad \dots (i)$$


Deviation suffered by the reflected ray after emerging out of sphere.

$$\delta_2 = (60^\circ - r_1) + (180^\circ - 2r_1) + (60^\circ - r_1)$$

$$\delta_2 = 300^\circ - 4r_1 \quad \dots (ii)$$

Now $\delta_1 = \frac{1}{3}\delta_2 \Rightarrow 3\delta_1 = \delta_2 \quad \dots (iii)$

From (i), (ii) and (iii) we get




$$3(120^\circ - 2r_1) = 300^\circ - 4r_1$$

$$\Rightarrow 360^\circ - 6r_1 = 300^\circ - 4r_1$$

$$\Rightarrow +2r_1 = +60^\circ \Rightarrow r_1 = 30^\circ \quad \dots (iv)$$

Snell's law of point P




$$\frac{\sin 60}{\sin r_1} = \mu \Rightarrow \mu = \frac{\sqrt{3}/2}{1/2} \Rightarrow \mu = \sqrt{3}$$

Sol 21: When air is filled between two similar glass pieces.

$$P_e = 2P_L + P_M$$

$$P_L = \frac{1}{f_L} \text{ where } \frac{1}{f_L} = \left(\frac{\mu_L - \mu_m}{\mu_m} \right) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$\Rightarrow \frac{1}{f_L} \left(\frac{1-1}{1} \right) \left(\frac{1}{R} - \frac{1}{-R} \right) \Rightarrow \frac{1}{f_L} = 0$$



Then $P_L = 0$ $P_M = -\frac{1}{f_M}$ where $f_M = -\frac{R}{2}$

(since mirror is concave)


Given that screen is placed at a distance 60 cm from the combination

$$m = -\frac{v}{u} \Rightarrow \text{Two times magnification means}$$

$$m = -2, \text{ then } v = 2u \Rightarrow 60 = 2u \Rightarrow v = 30 \text{ cm}$$

For equivalent combination $P_e = -\frac{1}{f_e} \Rightarrow f_e = -\frac{1}{P_e} = -\frac{R}{2}$


Apply mirror formula to the equivalent combination of mirror.



$$\frac{1}{v} + \frac{1}{u} = \frac{1}{f_e}$$

$$\Rightarrow \frac{1}{-60} + \frac{1}{-30} = \frac{1}{f_e} \Rightarrow -\frac{1}{60} - \frac{1}{30} = -\frac{2}{R} \left(\text{since } f_e = -\frac{R}{2} \right)$$

$$R = 40 \text{ cm}$$



Again if air between the glass pieces is replaced by water.

$$P_e = 2P_L + P_M$$

$$P_L = \frac{1}{f_L} \text{ where } \frac{1}{f_L} = \left(\frac{\mu_1 - 1}{1} \right) \left(\frac{1}{40} - \frac{1}{-40} \right)$$

$$\Rightarrow (\mu_w = 4/3) \Rightarrow P_L = \frac{1}{60} \Rightarrow P_M = -\frac{1}{f_M}$$

$$\text{Where } \frac{1}{f_M} = -\frac{R}{2} \text{ (for concave mirror)}$$

$$\frac{1}{f_M} = -\frac{R}{2} \text{ (for concave mirror)}$$

$$P_M = \frac{2}{R} = \frac{2}{40}$$

$$\Rightarrow P_e = 2P_L + P_M$$

$$2 \times \frac{1}{60} + \frac{2}{40} = \frac{5}{60} = \frac{1}{12}$$

$$-\frac{1}{f_e} \Rightarrow f_e = -12 \text{ cm}$$

$$\text{Again apply } \frac{1}{v} + \frac{1}{u} = \frac{1}{f_e}$$

$$\Rightarrow \frac{1}{-60} + \frac{1}{u} = \frac{1}{-12} \Rightarrow u = -15 \text{ cm}$$

When air is filled between the gap, object distance = 30 cm when water is filled between the gap, object distance = 15 cm, Then, object is displaced by 15 cm towards the combination.

Sol 22: (a) At surface light intensity

$$I = \frac{4.5 \pi W}{4\pi(1.5)^2 \times 10^{-4} \text{ m}^2} \Rightarrow I = \frac{1.125 \times 10^4}{2.25} \text{ Wm}^{-2}$$

$$= 5000 \text{ Wm}^{-2}$$

$$(b) I_p = \frac{4.5 \pi W}{4\pi(7.5)^2 \text{ m}^2} = 0.02 \text{ Wm}^{-2}$$

$$(c) u = -7.5 \text{ m} = -750 \text{ cm}$$

$$f = +30 \text{ cm}$$

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f} \Rightarrow \frac{1}{v} = \frac{1}{f} + \frac{1}{u} = \frac{1}{30} - \frac{1}{750} = \frac{14}{750} = \frac{7}{375}$$

$$\Rightarrow v = \frac{375}{7} \text{ cm,}$$

$$\text{Magnification, } m = \frac{v}{u} = \frac{375/7}{-750} \Rightarrow m = -\frac{1}{14}$$

$$\Rightarrow \text{diameter of image of bulb, } d_1 = \frac{3.0 \text{ cm}}{14} = 0.214 \text{ cm}$$

(d) Light intensity at image is the intensity focused by the lens.

$$I_{\text{image}} = \frac{0.02 \times \pi (7.5)^2 \times 10^{-4} \text{ W}}{4\pi \left(\frac{0.214}{2} \right)^2 \times 10^{-4} \text{ m}^2}$$

$$I_{\text{image}} = 24.56 \text{ Wm}^{-2}$$

Sol 23: Focal length of Plano - concave lens

$$\frac{1}{f_1} = (\mu_1 - 1) \left(\frac{1}{\infty} - \frac{1}{R} \right)$$

$$\frac{1}{f_1} = \left(\frac{3}{2} - 1 \right) \left(-\frac{1}{30} \right) = -\frac{1}{60} \Rightarrow f_1 = -60 \text{ cm}$$

Focal length of plano-convex lens

$$\frac{1}{f_2} = (\mu_2 - 1) \left(\frac{1}{\infty} - \frac{1}{-R} \right)$$

$$\frac{1}{f_1} = \left(\frac{5}{4} - 1 \right) \left(\frac{1}{30} \right) = \frac{1}{120} \Rightarrow f_2 = 120 \text{ cm}$$

(i) Plane surface of Plano-convex lens is silvered. So the equivalent focal length of the system,

$$\text{Power} = \frac{1}{-F} = \frac{1}{f_1} + \frac{1}{f_2} + \frac{1}{-f_m} + \frac{1}{f_2} + \frac{1}{f_1}$$

$$f_m = \text{focal length of planemirror} = \infty$$

$$\Rightarrow \frac{1}{-F} = \frac{2}{f_1} + \frac{2}{f_2} = -\frac{2}{60} + \frac{2}{120} = \frac{-1}{60}$$

$$\Rightarrow +F = +60 \text{ cm}$$

Focal length = 60 cm

The equivalent system behaves as a convex mirror.


(ii) Mirror formula $\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$. Hence $u = -15 \text{ cm}$, $f = +60 \text{ cm}$

$$\Rightarrow \frac{1}{v} = \frac{1}{60} - \frac{1}{-15} = \frac{1}{60} + \frac{4}{60} = \frac{5}{60}$$

$$\Rightarrow v = +12 \text{ cm}$$

$$\text{Magnification, } m = -\frac{v}{u} = \frac{-12}{-15} \Rightarrow m = \frac{4}{5}$$

Sol 24: Image from L_1 :



$$\frac{1}{v_1} = \frac{1}{f} + \frac{1}{u} = \frac{1}{15} - \frac{1}{20} = \frac{4-3}{60} = \frac{1}{60}$$

$$\Rightarrow v_1 = 60 \text{ cm from } L_1$$

For lens L_2 :

$$u_2 = +40 \text{ cm}, \frac{1}{v_2} = \frac{1}{15} + \frac{1}{40} = \frac{8+3}{120}$$

$$\Rightarrow v_2 = +\frac{120}{11} \text{ cm from } L_2$$

This final image should lie at the centre of curvature of convex mirror, so $MC = R = 5 \text{ cm}$

$$\text{So, } P_2M = P_2C_2 - MC = \frac{120}{11} - 5 = \frac{120-55}{11} \text{ cm}$$

$$\Rightarrow x = \frac{65}{11} \text{ cm} = 5.91 \text{ cm}$$

Sol 25: For equi-convex lens radius $R = f = 30 \text{ cm}$

Refraction at surface I: Air to glass

$$\frac{3/2}{v_1} - \frac{1}{(-90)} = \frac{3}{2} - \frac{1}{+30} \quad \dots (i)$$

Refraction at surface II: Glass to water

$$\frac{4/3}{v_2} - \frac{3/2}{v_1} = \frac{4}{3} - \frac{3}{2} \quad \dots (ii)$$

Add (i) and (ii)

$$\frac{4}{3v_2} + \frac{1}{90} = \frac{3}{2} - \frac{4}{3} + \frac{3}{2}$$

$$\Rightarrow \frac{4}{3v_2} = \frac{2}{90} - \frac{1}{90} \Rightarrow v_2 = +120 \text{ cm}$$

For mirror, image of lens acts as object.

For mirror $u_3 = + (120-80) = 40 \text{ cm}$ (right from mirror)

So $v_3 = -40 \text{ cm}$ (left from mirror)

Refraction from surface II after reflection from mirror.

$u_4 = -40 \text{ cm}$ (left from surface II)

$$\frac{3/2}{v_5} - \frac{4/3}{-40} = \frac{3}{2} - \frac{4}{3} \quad \dots (iii)$$

Refraction at surface I: Glass to air

$$\frac{1}{v_6} - \frac{3/2}{v_5} = \frac{1}{-30} \quad \dots (iv)$$

Add (iii) and (iv)

$$\frac{1}{v_6} + \frac{1}{30} = \frac{3}{2} - \frac{4}{3} - \frac{1}{-30} + \frac{3}{2}$$

$$\Rightarrow \frac{1}{v_6} = \frac{2}{90} - \frac{1}{30} = \frac{-1}{90} \Rightarrow v_6 = -90 \text{ cm}$$


from lens (right from lens)

So final image is 90 cm right of lens.

Exercise 2

Single Correct Choice Type


Sol 1: (B) Distance of image due to plane mirror from object will be 60 cm. So $OO' = 60 \text{ cm}$. So distance of image from convex mirror $PO' = [OO' - OP] = 10 \text{ cm}$



\Rightarrow Mirror formula $\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$ will give focal length of mirror by putting $u = -50 \text{ cm}$, $v = +10 \text{ cm}$,

$$\text{So, } \frac{1}{f} = \frac{1}{10} - \frac{1}{50} = \frac{4}{50} = \frac{2}{25} \Rightarrow f = \frac{25}{2}$$

Radius of curvature $R = 2f = 25 \text{ cm}$



Sol 2: (A) Mirror formula

$$\frac{1}{v} + \frac{1}{u} = \frac{1}{f} \Rightarrow \frac{1}{v} = \frac{1}{f} - \frac{1}{u} = \frac{u-f}{fu} \Rightarrow v = \frac{fu}{u-f}$$

At near end $|u| > |f|$. u and f both are negative.

So v is negative. At far end we have $u = -\infty$. So mirror

formula gives $\frac{1}{v_\infty} = \frac{1}{f} - \frac{1}{-\infty} = \frac{1}{f} \Rightarrow v_\infty = f$.

For near point $|v_{\text{near}}| > |f|$.

Image length

$$|v_{\text{near}}| - |v_{\infty}| = \frac{fu}{u-f} - f = \frac{fu - uf + f^2}{u-f}$$

$$\Rightarrow \Delta l = \frac{f^2}{u-f}$$

Sol 3: (C) Velocity of image

$$\frac{dv}{dt} = -\left(\frac{v^2}{u^2}\right)\frac{du}{dt}; \frac{dv}{dt} = -\left(\frac{v^2}{20^2}\right)4 \text{ cm s}^{-1}$$

From mirror formula

$$\frac{1}{v} = \frac{1}{f} - \frac{1}{u} \Rightarrow \frac{1}{v} = \frac{1}{-12} - \frac{1}{-20} = \frac{1}{20} - \frac{1}{12} = \frac{3-5}{60} = -\frac{1}{30}$$

$$\Rightarrow v = -30 \text{ cm} \Rightarrow \frac{dv}{dt} = -\left(\frac{900}{400}\right)4 \text{ cm s}^{-1} = -9 \text{ cm s}^{-1}$$

Sol 4: (D) Area of mirror, $A_1 = \pi \frac{d^2}{4}$.


Area left after putting opaque,

$$A_2 = A_1 - \pi \frac{d^2}{16} = \pi \frac{d^2}{4} - \pi \frac{d^2}{16}$$


$$\Rightarrow A_2 = \pi \frac{d^2}{4} \left(1 - \frac{1}{4}\right) = \frac{3}{4} \pi \frac{d^2}{4} = \frac{3}{4} A_1$$

Focal length will not change and intensity become $\frac{3}{4}I$.

Sol 5: (B) Rays should fall normally on plane mirror. This will happen if rays become parallel to principal axis after passing through lens. So $OL = f = 30 \text{ cm}$.



Sol 6: (D) Lens formula $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$; $u = -30 \text{ cm}$,
 $f = 20 \text{ cm}$



$$\frac{1}{v} = \frac{1}{20} - \frac{1}{30} = \frac{3-2}{60} = \frac{1}{60} \Rightarrow v = 60$$

$$\text{Magnification } m = \frac{v}{u} \Rightarrow m = \frac{60}{-30} = -2 \Rightarrow h_2 = -2h_1$$

So $h_2 = -2 \times 0.5 \text{ cm} = -1 \text{ cm}$ (below axis)

So Image of P is 1.5 cm below XY.

Sol 7: (B) Distance between object and screen is D , displacement of lens is d , and so focal length of lens is

$$f = \frac{D^2 - d^2}{4D} = \frac{90^2 - 20^2}{4 \times 90} = 21.4 \text{ cm}$$

Sol 8: (A) Object size

$$O = \sqrt{I_1 \times I_2} = \sqrt{6 \text{ cm} \times 3 \text{ cm}}$$

$$O = 4.24 \text{ cm}$$

Sol 9: (A) $\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$

Let $u = +x$ (virtual) and $|f| = -f$ (concave mirror)

$$\Rightarrow \frac{1}{v} = -\frac{1}{|f|} - \frac{1}{x} = \frac{-x - |f|}{|f|x} \Rightarrow v = \frac{|f|x}{-x - |f|} \rightarrow (-)ve$$

So v is always negative when u is positive ($+x$).

Multiple Correct Choice Type


Sol 10: (B, D) Slope of reflecting surface at the desired point will be $\tan 45^\circ = 1$

$$\frac{dy}{dx} = 2 \cos\left(\frac{\pi x}{L}\right) = 1 \Rightarrow \frac{\pi x}{L} = \frac{\pi}{3} \Rightarrow x = \frac{L}{3}$$

$$\Rightarrow y\left(\frac{L}{3}\right) = \frac{2L}{\pi} \sin\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}L}{\pi}$$

Sol 11: (B, C) Length of object is $AB = (50-20) \text{ cm} = 30 \text{ cm}$. After first reflection from plane mirror AB is inverted to B_1A_1 with distance from convex mirror as shown in figure.

Image of A in convex mirror




$$\frac{1}{v_1} = \frac{1}{f} - \frac{1}{u_1} \Rightarrow \frac{1}{v_1} = \frac{1}{60} - \frac{1}{-60} = \frac{1}{30} \Rightarrow v_1 = +30 \text{ cm}$$

Image of B in convex mirror

$$\frac{1}{v_2} = \frac{1}{f} - \frac{1}{u_2} = \frac{1}{60} - \frac{1}{-90} = \frac{3+2}{180} = \frac{1}{36}$$

$$\Rightarrow v_2 = +36 \text{ cm} \Rightarrow A'B' = (36 - 30) \text{ cm} = 6 \text{ cm}$$

Second image $A'B'$ is virtual and $\left(\frac{1}{5}\right)^{\text{th}}$ of magnification w. r. t. AB and erect.



Now formula for speed of image for convex mirror is, $\frac{dv}{dt} = \frac{v^2}{u^2} \frac{du}{dt}$. As object moves towards mirror, the image also moves towards the mirror.

Sol 12: (B, C) $i + i' = 90^\circ$ from figure

At point P, $\mu_1 \sin i > \mu_2$... (i)

At Q, $\mu_1 \sin i' > \mu_3$ or $\mu_1 \cos i > \mu_3$... (ii)

Squaring and adding (1) and (2) to get

$$\mu_1^2 > \mu_2^2 + \mu_3^2 \Rightarrow \mu_1^2 - \mu_2^2 > \mu_3^2 \Rightarrow \mu_1^2 - \mu_3^2 > \mu_2^2$$


Sol 13: (A, C) Angle of deviation $\delta = (\mu - 1)A$.

Sol 14: (A, B, D) Angle of deviation $\delta = (i_1 + i_2) - A$. i_1 is angle of incidence and i_2 is angle of emergence and angle of incidence and emergence are interchangeable.


If δ_m is minimum deviation, $\frac{\mu_p}{\mu_s} = \frac{\sin\left(\frac{\delta_m + A}{2}\right)}{\sin\left(\frac{A}{2}\right)}$

So as μ_p increases, δ_m also increases.

Sol 15: (B, C) The minimum length of a plane mirror to see one's full height in it is $\frac{H}{2}$, where H is the height of man. The mirror can be placed anywhere between the centre line BF (of AC) and DG (of CE). Eye is at C .



Sol 16: (B, D) The distance PQ_1 and PQ_2 will not change as the mirror MM' moves with speed v perpendicular to its length.



Sol 17: (A, C) $\mu_2 > \mu_1$ Rays from real object will be deviated away from radius of curvature and hence will become more diverging. For virtual object the deviated rays may converge on the principle axis.

$\mu_1 > \mu_2$: For virtual object the deviated rays will converge on principle axis.

For real object the refracted ray will deviate towards radius of curvature and may converge on principle axis.

Sol 18: (A, D) The image formed by a convex mirror is always, virtual and erect. So convex mirror cannot form inverted image of OA . Option B and C are ruled out.

Sol 19: (A, C) $u = -40$ cm; $f = +20$ cm

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f} \Rightarrow \frac{1}{v} = \frac{1}{f} + \frac{1}{u} = \frac{1}{20} - \frac{1}{40} = \frac{1}{40}$$

$$\Rightarrow v = +40 \text{ cm, and magnification } m = \frac{v}{u} = \frac{40}{-40}$$

$\Rightarrow m = -1 \Rightarrow y$ and z coordinate's values will change their sign but magnitude will remain the same.

$$\Rightarrow y_{\text{image}} = -1, z_{\text{image}} = +1$$

Sol 20: (B, D) Convex mirror and concave lens for diminished, virtual, correct image of object.

Assertion Reasoning Type

Sol 21: (C) The wall does reflect light, but the reflection is irregular. Reflected rays are deviated in different directions.

Sol 22: (A) This phenomena is called spherical aberration. The rays close to the principal axis are focused at the geometrical focus F of the mirror as given by mirror formula. The rays farthest from the principal axis are focused at a point somewhat closer to mirror.

Sol 23: (D) For object in liquid. $d_{\text{apparent}} = \frac{d_{\text{actual}}}{\mu}$

For a slab: normal shift $\Delta x = t \left(1 - \frac{1}{\mu} \right)$ where t is thickness of slab.


Sol 24: (D) Two image will be formed one for each mirror.

Sol 25: (D) If a plane mirror is moved such that its perpendicular distance from the point object does not change, then the image will not move.

Comprehension Type


Sol 26: (B) Paraxial rays are focused at the geometrical focus F of the mirror. The marginal rays are focused at a point F' somewhat closer to the mirror.

Sol 27: (D)



$$f = R - \frac{R}{2} \sec 60^\circ = R - R = 0$$

Sol 28: (D) Deviation = $180^\circ - 2 \times 60^\circ = 60^\circ$. Ans (D)



$$\text{Sol 29: (B)} \quad f = R - \frac{R}{2} \sec 0^\circ = \frac{R}{2}$$

Sol 30: (B) Spherical aberration cannot be completely eliminated, but it can be minimized by allowing either paraxial or marginal rays to hit the mirror.


Sol 31: (D)

$$y_{\text{min}} = \frac{(2n-1)\lambda D}{2d} \quad (n = 1, 2, \dots)$$

$$\text{For } n = +3, y_{\text{min}} = \frac{5\lambda D}{2d}$$

Sol 32: (A) Shift in the fringe due to the glass slab is $\Delta y = (\mu - 1)t \frac{D}{d}$ where t is thickness of glass slab. Due to glass slab path of ray from S_2 gets increased by $(\mu - 1)t$.


Sol 33: (A) Path of rays 1 is more than path of ray 2 by a distance $d \sin \alpha$. Draw perpendicular S_2M from S_2 to ray 1.



$$\angle MS_2S_1 = \alpha \quad \text{and} \quad MS_1 = d \sin \alpha$$

This path difference is suffered before passing the slits S_1 & S_2 . After passing through the slits, path of ray from

S_2 is increased by $(\mu - 1)t$. For net path difference to be zero at point P we have,



$$d \sin \alpha = (\mu - 1)t$$

$$\Rightarrow \alpha = \sin^{-1} \left(\frac{(\mu - 1)t}{d} \right)$$

Match the Column

Sol 34: A \rightarrow p, r; B \rightarrow q, s; C \rightarrow p, q, r, s; D \rightarrow p, q, r, s

Previous Years' Questions

Sol 1: (C, D) $\frac{1}{f} = \frac{1}{v} + \frac{1}{u}$ (mirror formula)

$$F = -24 \text{ cm}$$

Sol 2: (B) Critical angle from region III to region IV

$$\sin \theta_c = \frac{n_0/8}{n_0/6} = \frac{3}{4}$$

Now applying Snell's law in region I and region III

$$n_0 \sin \theta = \frac{n_0}{6} \sin \theta_c$$

$$\text{or } \sin \theta = \frac{1}{6} \sin \theta_c = \frac{1}{6} \left(\frac{3}{4} \right) = \frac{1}{8}$$

$$\therefore \theta = \sin^{-1} \left(\frac{1}{8} \right)$$

Sol 3: (A, C, D) In case of concave mirror or convex lens image can be real, virtual, diminished, magnified or of same size.

(B) In case of convex mirror image is always virtual (for real object).

Sol 4: (A) At minimum deviation ($\delta = \delta_m$):

$$r_1 = r_2 = \frac{A}{2} = \frac{60^\circ}{2} = 30^\circ \text{ (For both colours)}$$

Sol 5: (B) $\lambda_{\text{cutoff}} = \frac{hc}{eV}$ (independent of atomic number)

Sol 6: (C) The refractive index n for meta-material is negative.

Hence $\frac{\sin \theta_1}{\sin \theta_2}$ is negative.

Thus if θ_1 is negative, θ_2 will be negative. So the current choice is C.

Sol 7: (B) $N = \frac{c}{v} = v = \frac{c}{|n|}$ which is choice B

Also frequency $\nu = \frac{v}{\lambda}$ since ν remains unchanged

$$\frac{v_{\text{air}}}{\lambda_{\text{air}}} = \frac{v_m}{\lambda_m}$$

$$\Rightarrow \lambda_m = \lambda_{\text{air}} \times \frac{v_m}{v_{\text{air}}} = \lambda_{\text{air}} \times \frac{v_m}{c} \times \frac{c}{v_{\text{air}}}$$

$$= \lambda_{\text{air}} \times \frac{n_{\text{air}}}{n_m} \quad \left(\because v = \frac{c}{n} \right)$$

$$\frac{\lambda_{\text{air}}}{|n|} \quad \left(\because n_m = |n| \text{ and } n_{\text{air}} = 1 \right)$$

So choice D is wrong

Sol 8: (B) $P_T = (1.5 - 1) \left(\frac{1}{14} - 0 \right) + (1.2 - 1) \left(0 - \frac{1}{-14} \right)$

$$= \frac{0.5}{14} + \frac{0.2}{14} = \frac{1}{20}$$

$$f = +20 \text{ cm}$$


$$\frac{1}{v} - \frac{1}{-40} = \frac{1}{20}$$

$$\frac{1}{v} = \frac{1}{20} - \frac{1}{40} = \frac{1}{40}$$

$$\therefore v = 40 \text{ cm}$$

Sol 9: (B) Object is placed at distance $2f$ from the lens. So first image I_1 will be formed at distance $2f$ on other side. This image I_1 will behave like a virtual object for mirror. The second image I_2 will be formed at distance 20 cm in front of the mirror, or at distance 10 cm to the left hand side of the lens.

Now applying lens formula



$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

$$\therefore \frac{1}{v} - \frac{1}{+10} = \frac{1}{+15}$$

or $v = 6 \text{ cm}$

Therefore, the final image is at distance 16 cm from the mirror. But, this image will be real.

This is because ray of light is travelling from right to left.

Sol 10: (3) For $v_1 = \frac{50}{7} \text{ m}$, $u_1 = -25 \text{ m}$

$$v_2 = \frac{25}{3} \text{ m}, u_2 = -50 \text{ m}$$

$$\text{Speed of object} = \frac{25}{30} \times \frac{18}{5} = 3 \text{ km/h.}$$

Sol 11: (B)

$$\text{For the combination } \frac{1}{f_{\text{eq}}} = \frac{(\mu_1 - 1)}{R} + \frac{(\mu_2 - 1)}{R}$$

$$f_{\text{eq}} = 20$$

Here $u = -40$, $f = 20$


$$v = 40$$

Sol 12: (A, C, D) From Snell's Law

$$n_1 \sin \theta_i = n(d) \sin \theta_d = n_2 \sin \theta_f$$

The deviation of ray in the slab will depend on $n(z)$

Hence, l will depend on $n(z)$ but not on n_2 .




Sol 13: (A) First Image I_1 from the lens will be formed at 75 cm to the right of the lens.

Taking the mirror to be straight, the image I_1 after reflection will be formed at 50 cm to the left of the mirror.

On rotation of mirror by 30° the final image is I_3 .


$$\text{So } x = 50 - 50 \cos 60^\circ = 25 \text{ cm.}$$

$$\text{and } y = 50 \sin 60^\circ = 25\sqrt{3} \text{ cm.}$$



Sol 14: (A) Let angle between the directions of incident ray and reflected ray be θ

$$\cos \theta = \frac{1}{2} (\hat{i} + \sqrt{3} \hat{j})$$




$$\cos \theta = \frac{1}{2} (\hat{i} + \sqrt{3} \hat{j}) \cdot \frac{1}{2} (\hat{i} + \sqrt{3} \hat{j})$$

$$\cos \theta = -\frac{1}{2}$$

$$\theta = 120^\circ$$

Sol 15: (C)



$$\mu = \frac{\lambda_a}{\lambda_m} = \frac{3}{2}$$

$$\Rightarrow \frac{1}{f} = \frac{\mu - 1}{R} = \frac{1}{2R}$$

$$\Rightarrow \frac{1}{f} = \frac{1}{v} - \frac{1}{u}$$

$$\Rightarrow \frac{1}{8} - \frac{1}{-24} = \frac{1}{2R}$$

$$\Rightarrow \frac{3+1}{24} = \frac{1}{2R}$$

$$\Rightarrow R = 3 \text{ m}$$

Sol 16: (D) P \rightarrow (2); Q \rightarrow (3); R \rightarrow (4); S \rightarrow (1)

P. $\mu_2 > \mu_1$ (towards normal)

$\mu_2 > \mu_3$ (away from normal)

Q. $\mu_1 = \mu_2$ (No change in path)

$\angle i = 0 \Rightarrow \angle r = 0$ on the block.

R. $\mu_1 > \mu_2$ (Away from the normal)

$\mu_2 > \mu_3$ (Away from the normal)

$$\mu_1 \times \frac{1}{\sqrt{2}} = \mu_2 \sin r \Rightarrow \sin r = \frac{\mu_1}{\sqrt{2} \mu_2}$$

Since $\sin r < 1 \Rightarrow \mu_1 < \sqrt{2} \mu_2$

S. For TIR:

$$45^\circ > C \Rightarrow \sin 45^\circ > \sin C \Rightarrow \frac{1}{\sqrt{2}} > \frac{\mu_2}{\mu_1} \Rightarrow \mu_1 > \sqrt{2} \mu_2$$

Sol 17: (A, C) For air to glass

$$\frac{1.5}{f_1} = \frac{1.4 - 1}{R} + \frac{1.5 - 1.4}{R}$$


$$\therefore f_1 = 3R$$

For glass to air.

$$\frac{1}{f_2} = \frac{1.4 - 1.5}{-R} + \frac{1 - 1.4}{-R}$$

$$\therefore f_2 = 2R$$


Sol 18: (C) $\tan \theta_c = \frac{r}{h} = \frac{5.77}{10} \approx \sqrt{3}$



$$\Rightarrow \sin \theta_c = \frac{\mu_\ell}{\mu_b}$$

$$\Rightarrow \mu_\ell = 2.72 \times \frac{1}{2} = 1.36$$

Sol 19: (B) $\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$



Use $\frac{1}{f_{eq}} = \frac{1}{f_1} + \frac{1}{f_2}$

(P) $\frac{1}{f_{eq}} = \frac{1}{R} + \frac{1}{R} = \frac{2}{R}; f_{eq} = \frac{R}{2}$

(Q) $\frac{1}{f_{eq}} = \frac{1}{2R} + \frac{1}{2R} = \frac{1}{R}; f_{eq} = R$

(R) $\frac{1}{f_{eq}} = -\frac{1}{2R} - \frac{1}{2R} = -\frac{1}{R}; f_{eq} = -R$

(S) $\frac{1}{f_{eq}} = \frac{1}{R} - \frac{1}{2R} = \frac{1}{2R}; f_{eq} = 2R$

Sol 20: Image by mirror is formed at 30 cm from mirror at its right and finally by the combination it is formed at 20 cm on right of the lens. So in air medium, magnification by lens is unity. In second medium,

$\mu = \frac{7}{6}$. focal length of the lens is given by,

$$\frac{1}{10} = \frac{(1.5-1)\left(\frac{1}{R_1} - \frac{1}{R_2}\right)}{\frac{1}{f} \left(\frac{1.5}{7/6} - 1\right)\left(\frac{1}{R_1} - \frac{1}{R_2}\right)} \Rightarrow f = \frac{35}{2} \text{ cm}$$

So in second medium, final image is formed at 140 cm to the right of the lens. Second medium does not change

the magnification by mirror. So $\left|\frac{M_2}{M_1}\right| = \left|\frac{M_{m_2} M_{\ell_2}}{M_{m_1} M_{\ell_1}}\right| = 7$

Sol 21: (B) For 1st refraction

$$\frac{1}{v} - \frac{1.5}{-50} = \frac{1-1.5}{-10}$$

$$\Rightarrow v = 50 \text{ cm}$$

For 2nd refraction

$$\frac{1.5}{\infty} - \frac{1}{-x} = \frac{1.5-1}{+10}$$

$$\Rightarrow x = 20 \text{ cm}$$

$$\Rightarrow d = 70 \text{ cm}$$

Sol 22: Snell's Law on 1st surface: $\frac{\sqrt{3}}{2} = n \sin r_1$

$$\sin r_1 = \frac{\sqrt{3}}{2n} \quad \dots(i)$$

$$\Rightarrow \cos r_1 = \sqrt{1 - \frac{3}{4n^2}} = \frac{\sqrt{4n^2 - 3}}{2n}$$

$$r_1 + r_2 = 60^\circ \quad \dots(ii)$$

Snell's Law on 2nd surface:

$$n \sin r_2 = \sin \theta$$

Using equation (i) and (ii)

$$n \sin(60^\circ - r_1) = \sin \theta$$

$$n \left[\frac{\sqrt{3}}{2} \cos r_1 - \frac{1}{2} \sin r_1 \right] = \sin \theta$$

$$\frac{d}{dn} \left[\frac{\sqrt{3}}{4} (\sqrt{4n^2 - 3} - 1) \right] = \cos \theta \frac{d\theta}{dn}$$

For $\theta = 60^\circ$ and $n = \sqrt{3}$

$$\Rightarrow \frac{d\theta}{dn} = 2$$

Sol 23: (A, C) $\theta \geq c$

$$\Rightarrow 90^\circ - r \geq c$$


$$\Rightarrow \sin(90^\circ - r) \geq \sin c$$

$$\Rightarrow \cos r \geq \sin c$$

$$\text{Using } \frac{\sin i}{\sin r} = \frac{n_1}{n_m} \text{ and } \sin c = \frac{n_2}{n_1}$$


$$\text{We get, } \sin^2 i_m = \frac{n_1^2 - n_2^2}{n_m^2}$$

Putting values, we get, correct options as A & C



Sol 24: (A) $i = \beta + \theta$

For $\alpha = 45^\circ$; by Snell's law,



$$1 \times \sin 45^\circ = \sqrt{2} \sin \beta$$

$$\Rightarrow \beta = 30^\circ$$

For TIR on face PR,

$$\beta + \theta = \theta_c = \sin^{-1} \left(\frac{1}{\sqrt{2}} \right) = 45^\circ$$

$$\Rightarrow \theta = 45^\circ - \beta = 15^\circ$$

Sol 25: (A, D) For refraction through lens,

$$\frac{1}{v} - \frac{1}{-30} = \frac{1}{f} \text{ and } -2 = \frac{v}{u}$$

$$\therefore v = -2u = 60 \text{ cm}$$

$$\therefore f = +20 \text{ cm}$$

For reflection

$$\frac{1}{10} + \frac{1}{-30} = \frac{2}{R} \Rightarrow R = 30 \text{ cm}$$

$$(n-1)\left(\frac{1}{R}\right) = \frac{1}{f} = \frac{1}{20}$$

$$\therefore n = \frac{5}{2}$$

The faint image is erect and virtual.

Empower your JEE Main & Advanced preparation with specially designed and structured study packages from PLANCESS

Planceess offers this must-have Study Package in Physics comprising six books to meet the complete curriculum needs of engineering aspirants. The books offer complete and systematic coverage of JEE (Main) and JEE (Advanced) syllabi and aims to provide firm foundation in learning and problem solving skills.

The theory offers engaging explanation of concepts with important points highlighted through Facts, Notes, Tips & Tricks, Misconceptions and Key Take Away. The application of the concepts is explained by means of illustrative examples and the problem-solving skills are further enhanced through solved examples of JEE (Main) and JEE (Advanced) levels. Extensive and classified assessment (JEE Main, JEE Advanced and Boards) along with the questions from previous years papers of the target examinations, offers the aspirants a competitive edge in preparation. This study package thus provides a single solution for all their preparation requirements.

Salient Features

- ▶ Concept description sprinkled with Planceess Concepts and Illustrative Examples
- ▶ Problem-Solving Strategy (Tactics)
- ▶ Formula Sheet/Points to Remember
- ▶ Solved Examples - JEE Main
- ▶ Solved Examples - JEE Advanced
- ▶ Exercises - JEE Main and Boards
- ▶ Previous Years' Questions - JEE Main
- ▶ Exercises - JEE Advanced and Boards
- ▶ Previous Years Questions - JEE Advanced
- ▶ Answer Key and Solutions (Complete)

Planceess EduSolutions Pvt. Ltd.

A005/A006 Boomerang
Chandivali Farm Road,
Chandivali, Andheri (East)
Mumbai - 400072, India
Tel: 022 6604 3405
www.planceess.com

