

* Equation of tangent to the curve $y = f(x)$ at $A(x_1, y_1)$ is $y - y_1 = \left(\frac{dy}{dx} \right)_{(x_1, y_1)} (x - x_1)$

* Equation of normal at (x_1, y_1) to the curve $y = f(x)$ is $(y - y_1) = \frac{-1}{\left(\frac{dy}{dx} \right)_{(x_1, y_1)}} (x - x_1)$

* Length of Tangent, Normal, Subtangent and Subnormal

$$\text{Tangent: } PT = MP \operatorname{cosec} \Psi = y \sqrt{1 + \cot^2 \Psi} = \left| \frac{y \sqrt{1 + \left(\frac{dy}{dx} \right)^2}}{\frac{dy}{dx}} \right|$$

$$\text{Subtangent: } TM = MP \cot \Psi = \left| \frac{y}{(dy/dx)} \right|$$

$$\text{Normal: } GP = MP \sec \Psi = y \sqrt{1 + \tan^2 \Psi} = \left| y \sqrt{1 + \left(\frac{dy}{dx} \right)^2} \right|$$

$$\text{Subnormal: } MG = MP \tan \Psi = \left| y \left(\frac{dy}{dx} \right) \right|$$

* Angle of Intersection of Two Curves

$$\tan \Psi = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|,$$

where m_1 and m_2 are the slopes of the tangents T_1 and T_2 at the intersection point (x_1, y_1) .




Figure 21.9




Figure 21.10

Solved Examples

JEE Main/Boards

Example 1: Show that the function $f(x) = |x|$ is continuous at $x = 0$, but not differentiable at $x = 0$.

Sol: Evaluate $f'(0^+)$ and $f'(0^-)$.

$$\text{We have } f(x) = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

$$\text{Since } \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^-} f(x) = 0 = f(0)$$

The function is continuous at $x = 0$

We also have

$$f'(0^+) = \lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x} = \lim_{x \rightarrow 0^+} \frac{x - 0}{x} = 1$$

$$f'(0^-) = \lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x} = \lim_{x \rightarrow 0^-} \frac{-(-x) - 0}{-x} = -1$$

Since, $f'(0^+) \neq f'(0^-)$, the function is not differentiable at $x = 0$

Example 2: Find the derivative of the function $f(x)$, defined by $f(x) = \sin x$ by 1st principle.

Sol: Use the first principle to find the derivative of the given function.

Let dy be the increment in y corresponding to an increment dx in x . We have

$$y = \sin x$$

$$y + dy = \sin(x + dx)$$

Subtracting, we get

$$dy = \sin(x + dx) - \sin x = 2 \cos(x + dx/2) \sin(dx/2)$$

Dividing by dx , we obtain

$$\frac{\delta y}{\delta x} = \frac{\cos(x + \delta x/2) \sin(\delta x/2)}{(\delta x/2)}$$

Taking limits on both side, we get

$$\frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} = \lim_{\delta x \rightarrow 0} \cos(x + dx/2) \lim_{\delta x \rightarrow 0} \frac{\sin(\delta x/2)}{(\delta x/2)}$$

$$= \cos x \cdot 1 = \cos x$$

Hence, we have $d/dx(\sin x) = \cos x$

Example 3: The derivative of $\log |x|$ is

Sol: Use the definition of the modulus to expand the given function. Then evaluate L.H.D. and R.H.D. at the critical point.

Let $y = \log |x|$ then

$$y = \begin{cases} \log x, & \text{when } x > 0 \\ \log(-x), & \text{when } x < 0 \end{cases}$$

$$\therefore \frac{dy}{dx} = \frac{1}{x} \text{ when } x > 0$$

$$\text{and } \frac{dy}{dx} = \frac{1}{-x} (-1) = \frac{1}{x} \text{ when } x < 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{x} \text{ when } x \neq 0$$

Example 4: If $y = \sqrt{\frac{1+x}{1-x}}$, then $\frac{dy}{dx}$ equals

Sol: Differentiate using u/v rule.

Differentiating w.r.t., we get

$$\frac{dy}{dx} = \frac{1}{2} \sqrt{\frac{1-x}{1+x}} \frac{(1-x)1 - (1+x)(-1)}{(1-x)^2}$$

$$\Rightarrow \sqrt{\frac{1-x}{1+x}} \frac{1}{(1-x)^2} \Rightarrow \frac{1}{\sqrt{1+x}} \frac{1}{(1-x)^{3/2}}$$

$$\Rightarrow \frac{1}{\sqrt{1-x^2}} \left(\frac{1}{1-x} \right)$$

Example 5: If $x^y = e^{x-y}$, then dy/dx equals -

Sol: Take logarithms on both sides and differentiate.

Taking log on both sides, we get

$$y \log x = x - y \Rightarrow y = \frac{x}{1 + \log x}$$

$$\therefore \frac{dy}{dx} = \frac{(1 + \log x) \cdot 1 - x(1/x)}{(1 + \log x)^2} = \frac{\log x}{(1 + \log x)^2}$$

Example 6: If $y = \cot^{-1} \sqrt{x^2 - 1} + \sec^{-1} x$, then dy/dx equals

Sol: Use substitution to simplify the terms and then differentiate.

$$\text{Put } x = \sec \theta; \cot^{-1} \sqrt{x^2 - 1} = \cot^{-1} \sqrt{\sec^2 \theta - 1}$$

$$= \cot^{-1}(\tan \theta) = \frac{\pi}{2} - \theta = \frac{\pi}{2} - \sec^{-1} x$$

$$\therefore y = \left(\frac{\pi}{2} - \sec^{-1} x \right) + \sec^{-1} x = \frac{\pi}{2} \Rightarrow \frac{dy}{dx} = 0$$

Example 7: If $x^2 e^y + 2xy e^x + 13 = 0$, then $\frac{dy}{dx}$ equals -

Sol: Use the formula for derivative of implicit function.

Using partial derivatives, we have

$$\Rightarrow \frac{dy}{dx} = \frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial y}}$$

$$\frac{dy}{dx} = - \frac{2xe^y + 2ye^x + 2xye^x}{x^2 e^y + 2xe^x}$$

$$= - \frac{2xe^{y-x} + 2y + 2xy}{x^2 e^{y-x} + 2x} = - \frac{2xe^{y-x} + 2y(x+1)}{x(xe^{y-x} + 2)}$$

Example 8: $\frac{d}{dx} \left[\tan^{-1} \left(\frac{\cos x}{1 + \sin x} \right) \right]$ equals -

Sol: Convert $\frac{\cos x}{1 + \sin x}$ in terms of \tan and proceed.

$$\therefore \frac{\cos x}{1 + \sin x} = \frac{\sin(\pi/2 - x)}{1 + \cos(\pi/2 - x)}$$

$$= \tan \left(\frac{\pi}{4} - \frac{x}{2} \right) \therefore \tan^{-1} \left(\frac{\cos x}{1 + \sin x} \right)$$

$$= \tan^{-1} \left(\tan \left(\frac{\pi}{4} - \frac{x}{2} \right) \right) = \frac{\pi}{4} - \frac{x}{2} \Rightarrow \text{Derivative} = -\frac{1}{2}$$

Example 9: If $y = \frac{\sec x - \tan x}{\sec x + \tan x}$, then $\frac{dy}{dt}$ equals

Sol: Simplify the R.H.S. and differentiate.

$$y = \frac{\sec x - \tan x}{\sec x + \tan x} \cdot \frac{\sec x - \tan x}{\sec x - \tan x}$$

$$y = (\sec x - \tan x)^2$$

$$\therefore \frac{dy}{dx} = 2(\sec x - \tan x)(\sec x \tan x - \sec^2 x)$$

$$\Rightarrow -2 \sec x (\sec x - \tan x)^2$$

Example 10: If $y = \frac{1}{(t+2)(t+1)}$, then $\frac{dy}{dx}$ equals

Sol: Use the partial fraction method to find the derivative of given f^n

$$y = \frac{1}{(t+2)(t+1)} = \frac{1}{t+1} - \frac{1}{t+2}$$

$$\Rightarrow \frac{dy}{dx} = -\frac{1}{(t+1)^2} + \frac{1}{(t+2)^2}$$

Example 11: If $x = \theta - \frac{1}{\theta}$ and $y = \theta + \frac{1}{\theta}$,

then $\frac{dy}{dx} = ?$

Sol: $x = \theta - \frac{1}{\theta} \Rightarrow \frac{dx}{d\theta} = 1 + \frac{1}{\theta^2}$

$$y = \theta + \frac{1}{\theta} \Rightarrow \frac{dy}{d\theta} = 1 - \frac{1}{\theta^2}$$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{1 - (1/\theta^2)}{1 + (1/\theta^2)} = \frac{\theta - (1/\theta)}{\theta + (1/\theta)} = \frac{x}{y}$$

Example 12: Derivative of $\sin^{-1} x$ w.r.t. $\cos^{-1} \sqrt{1-x^2}$ is -

Sol: Substitute $\sin \theta$ in place of x .

$$\text{Let } y = \sin^{-1} x \text{ and } z = \cos^{-1} \sqrt{1-x^2}$$

$$\text{Put } x = \sin \theta \Rightarrow z = \cos^{-1}(\cos \theta) = \theta$$

$$\therefore y = z \text{ and } \frac{dy}{dz} = 1$$

Example 13: Derivative of $\sec^{-1} \left(\frac{1}{2x^2+1} \right)$ w.r.t. $\sqrt{1+3x}$

at $x = \frac{-1}{3}$ is

Sol: Differentiate the two functions and divide.

$$\text{Let } y = \sec^{-1} \left(\frac{1}{2x^2+1} \right) \text{ and } z = \sqrt{1+3x}$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = (2x^2+1) \frac{1}{\sqrt{(1/(2x^2+1))^2 - 1}}$$

$$\left(\frac{-4x}{2x^2+1} \right) \frac{2}{3} \sqrt{1+3x} \therefore \left(\frac{dy}{dz} \right)_{x=-\frac{1}{3}} = 0$$

Example 14: Find $\frac{d}{dx} \left\{ \sin^2 \left(\cot^{-1} \sqrt{\frac{1+x}{1-x}} \right) \right\} =$

Sol: Use Substitution to simplify the inside the square root and then differentiate.

$$\text{Let } y = \sin^2 \left(\cot^{-1} \sqrt{\frac{1+x}{1-x}} \right).$$

$$\text{Put } x = \cos 2\theta.$$

$$\therefore y = \sin^2 \cot^{-1} \left\{ \left(\sqrt{\frac{1+\cos 2\theta}{1-\cos 2\theta}} \right) \right\} = \sin^2 \cot^{-1} (\cot \theta)$$

$$\therefore y = \sin^2 \theta = \frac{1-\cos 2\theta}{2} = \frac{1-x}{2} = \frac{1}{2} - \frac{x}{2}$$

$$\therefore \frac{dy}{dx} = -\frac{1}{2}$$

Example 15: Find the equation of the normal to the curve $y = x + \sin x \cos x$ at $x = \frac{\pi}{2}$.

Sol: Find a point on the curve slope of the normal at that point.

$$x = \frac{\pi}{2} \Rightarrow y = \frac{\pi}{2} + 0 = \frac{\pi}{2}, \text{ so the given point} = \left(\frac{\pi}{2}, \frac{\pi}{2} \right).$$

$$\text{Now from the given equation } \frac{dy}{dx} = 1 + \cos^2 x - \sin^2 x$$

$$\Rightarrow \left(\frac{dy}{dx} \right)_{\left(\frac{\pi}{2}, \frac{\pi}{2} \right)} = 1 + 0 - 1 = 0$$

\therefore required equation of the normal is

$$y - \frac{\pi}{2} = \frac{-1}{0} \left(x - \frac{\pi}{2} \right) \Rightarrow x - \frac{\pi}{2} = 0 \Rightarrow 2x = \pi$$

Example 16: Find the point on the curve $y = x^2 - 3x$ at which tangent is parallel to x-axis.

Sol: Differentiate the given equation and put it equal to zero and proceed.

Let the point at which tangent is parallel to x-axis be $P(x_1, y_1)$

Then it must be on curve i.e., $y_1 = x_1^3 - 3x_1$

Also differentiating w.r.t. x , we get, $\frac{dy}{dx} = 3x^2 - 3$

$$\Rightarrow \left(\frac{dy}{dx}\right)_{(x_1, y_1)} = 3x_1^2 - 3 \quad \dots (i)$$

since, the tangent is parallel to x -axis

$$\therefore \left(\frac{dy}{dx}\right)_{(x_1, y_1)} = 0 \Rightarrow 3x_1^2 - 3 = 0$$

$$\Rightarrow x_1 = \pm 1 \quad \dots (ii)$$

From (1) and (2); $y_1 = x_1^3 - 3x_1$

When $x_1 = 1$ when $x_1 = -1$

$$y_1 = 1 - 3 = -2, y_1 = -1 + 3 = 2$$

\therefore points at which tangent is parallel to x -axis are $(1, -2)$ and $(-1, 2)$.

Example 17: Find the equation of normal to the curve $x + y = x^y$, where it cuts x -axis.

Sol: Given curve is $x + y = x^y$... (i)

at x -axis $y = 0$,

$$\therefore x + 0 = x^0 \Rightarrow x = 1$$

\therefore Point is $A(1, 0)$

Now to differentiation $x + y = x^y$

take log on both sides

$$\Rightarrow \log(x + y) = y \log x$$

$$\therefore \frac{1}{x+y} \left\{ 1 + \frac{dy}{dx} \right\} = y \cdot \frac{1}{x} + (\log x) \frac{dy}{dx}$$

$$\text{Putting } x = 1, y = 0, \left\{ 1 + \frac{dy}{dx} \right\} = 0$$

$$\Rightarrow \left(\frac{dy}{dx}\right)_{(1,0)} = -1 \quad \therefore \text{slope of normal} = 1$$

$$\text{Equation of normal is, } \frac{y-0}{x-1} = 1 \Rightarrow y = x - 1$$

Example 18: If the curve $y^2 = 6x$, $9x^2 + by^2 = 16$, cut each other at right angles then the value of b is

Sol: Equate the product of $\frac{dy}{dx}$ from the two equations to -1 .

The intersection of the two curves is given by $9x^2 + 6bx = 16$... (i)

Differentiating $y^2 = 6x$, We have $\frac{dy}{dx} = \frac{3}{y}$

Differentiating $9x^2 + 6y^2 = 16$

$$\text{We have } \frac{dy}{dx} = -\frac{9x}{by}$$

For curves to intersect at right angles, we must have at the point of intersection.

$$\frac{3}{y} \left(-\frac{9x}{by} \right) = -1 \Rightarrow 27x = by^2.$$

Thus we must have

$$9x^2 + by^2 = 16 \Rightarrow 9x^2 + 27x - 16 = 0 \quad \dots (ii)$$

(i) and (ii) must be identical so $27 = 6b \Rightarrow b = 9/2$.

Example 19: If the tangent at $(1, 1)$ on $y^2 = x(2-x)^2$ meets the curve again at P , then P is

Sol: Solve the equation of the tangent with the equation of the curve.

$$2y \frac{dy}{dx} = (2-x)^2 - 2x(2-x)$$

$$= 3x^2 - 8x + 4. \text{ So } \left. \frac{dy}{dx} \right|_{(1,1)} = -\frac{1}{2}$$

An equation of tangent at $(1, 1)$ is $Y - 1 = (-1/2)(X - 1)$.

i.e. $Y = (-1/2)x + 3/2$. The intersection of this line with the given curve is given by $((-x/2) + 3/2)^2 = x(2-x)^2$

$$\Rightarrow x^2 - 6x + 9 = 16x + 4x^3 - 16x^2. \text{ So,}$$

$$4x^3 - 17x^2 + 22x - 9 = 0$$

$$\Rightarrow (x-1)(4x-9)(x+1) = 0$$

Thus $x = 1, 9/4, -1$. But $x = -1$ cannot lie on the given curve so required point is $(9/4, 3/8)$.

JEE Advanced/Boards

Example 1: Examine differentiability of $f(x)$ at

$$x = 0 \text{ for } f(x) = \begin{cases} \frac{1 - \cos x}{x \sin x}, & x \neq 0 \\ \frac{1}{2}, & x = 0 \end{cases}$$

Sol: Find the left and right hand derivative of the function $f(x)$ about the point $x = 0$.

First we obtain $L.f'(0)$

$$= \lim_{h \rightarrow 0} \left[\frac{f(-h) - f(0)}{-h} \right] = \lim_{h \rightarrow 0} \left[\left(-\frac{1}{h} \right) \left\{ \frac{1 - \cosh}{h \sinh} - \frac{1}{2} \right\} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{h \sinh + 2(1 - \cosh)}{2h^2 \sinh} \right]; \left(\text{in } \frac{0}{0} \text{ form} \right)$$

$$= \lim_{h \rightarrow 0} \left[\frac{h \left(h - \frac{h^3}{3!} + \frac{h^5}{5!} - \dots \right) - 2 \left(\frac{h^2}{2!} - \frac{h^4}{4!} + \frac{h^6}{6!} - \dots \right)}{2h^2 \left(h - \frac{h^3}{3!} + \dots \right)} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{h^4 \left\{ \left(\frac{1}{12} - \frac{1}{3!} \right) + \left(\frac{1}{5!} - \frac{2}{6!} \right) h^2 \right\}}{2h^3 \left(h - \frac{h^3}{3!} + \dots \right)} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{h \left\{ \left(\frac{1}{12} - \frac{1}{3!} \right) + \left(\frac{1}{5!} - \frac{2}{6!} \right) h + \dots \right\}}{2 \left(1 - \frac{h^2}{3!} + \dots \right)} \right] = 0 \text{ and}$$

$$Rf'(0) = \lim_{h \rightarrow 0} \left(\frac{f(0+h) - f(0)}{h} \right)$$

$$= \lim_{h \rightarrow 0} \left\{ \frac{1}{h} \left(\frac{1 - \cosh}{h \sinh} - \frac{1}{2} \right) \right\} = 0,$$

similarly as above i.e. $Lf'(0) = Rf'(0)$

$\Rightarrow f(x)$ is differentiable at $x = 0$

Example 2: Examine differentiability of the function $f(x)$

$= \sin^{-1}(\cos x)$ at $x = n\pi + \frac{\pi}{2}$, where $n \in \mathbb{I}$.

Sol: Similar to the previous example.

first, we obtain $Lf' \left(nx + \frac{\pi}{2} \right)$

$$= \lim_{h \rightarrow 0} \left(\frac{f(nx + (\pi/2) - h) - f(nx + (\pi/2))}{-h} \right)$$

$$= \lim_{h \rightarrow 0} \left[\frac{\sin^{-1} \left\{ \cos(nx + (\pi/2) - h) \right\} - \sin^{-1} \left\{ \cos(nx + (\pi/2)) \right\}}{-h} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{\sin^{-1} \left\{ (-1)^n \cos((\pi/2) - h) \right\} - \sin^{-1} \left\{ (-1)^n \cos(\pi/2) \right\}}{-h} \right]$$

$$= \lim_{h \rightarrow 0} \left[\frac{\sin^{-1} \left\{ \sin(-1)^n h \right\} - \sin^{-1} 0}{-h} \right]$$

$$= \lim_{h \rightarrow 0} \frac{(-1)^n \sin^{-1} \sin h - \sin^{-1} 0}{-h}$$

$$= \lim_{h \rightarrow 0} \left[\frac{(-1)^n h}{-h} \right] = (-1)^{n-1} Rf' \left(n\pi + \frac{\pi}{2} \right)$$

$$= \lim_{h \rightarrow 0} \left\{ \frac{f(nx + (\pi/2) + h) - f(nx + (\pi/2))}{h} \right\}$$

$$= \lim_{h \rightarrow 0} \left\{ \frac{\sin^{-1} \left\{ \cos(nx + (\pi/2) + h) \right\} - \sin^{-1} \left\{ \cos(nx + (\pi/2)) \right\}}{h} \right\}$$

$$= \lim_{h \rightarrow 0} \left\{ \frac{\sin^{-1} \left\{ (-1)^n \cos((\pi/2) + h) \right\} - \sin^{-1} \left\{ (-1)^n \cos(\pi/2) \right\}}{h} \right\}$$

$$= \lim_{h \rightarrow 0} \left[\frac{\sin^{-1} \{ (-1)^{n+1} \sinh \}}{h} \right] = \lim_{h \rightarrow 0} \left[\frac{\sin^{-1} \{ \sin(-1)^{n+1} h \}}{h} \right]$$

$$= \lim_{h \rightarrow 0} \frac{(-1)^{n+1} \sin^{-1} \sin h}{h} = \lim_{h \rightarrow 0} \frac{(-1)^{n+1} h}{h} = (-1)^{n+1}$$

(Which is equal to $(-1)^{n-1}$)

Thus we find $Lf' \left(nx + \frac{\pi}{2} \right) = Rf' \left(nx + \frac{\pi}{2} \right)$

$\therefore f(x)$ is differentiable at $(nx + \pi/2)$

Example 3: If $x\sqrt{1+y} + y\sqrt{1+x} = 0$, then $\frac{dy}{dx}$ equals

Sol: Simplify the equation given and then differentiate it.

We have

$$x\sqrt{1+y} + y\sqrt{1+x} = 0 \quad \dots (i)$$

$$\Rightarrow x\sqrt{1+y} = -y\sqrt{1+x}$$

On squaring both sides $x^2(1+y) = y^2(1+x)$

$$\Rightarrow x^2 - y^2 + x^2y - xy^2 = 0$$

$$\Rightarrow (x-y)(x+y+xy) = 0$$

$x-y \neq 0$ [For $y=x$ does not satisfy (1)]

$$\therefore x+y+xy = 0 \Rightarrow y = -\frac{x}{(1+x)}$$

$$\therefore \frac{dy}{dx} = - \left\{ \frac{(1+x) \cdot 1 - x \cdot 1}{(1+x)^2} \right\} = - \frac{1}{(1+x)^2}$$

Example 4: If $x^y y^x = 1$, then $\frac{dy}{dx}$ equals -

Sol: Use logarithms on both sides and then differentiate

Taking log on both sides, we have $y \log x + x \log y = 0$

Now using partial derivatives, we have

$$\frac{dy}{dx} = -\frac{y/x + \log y}{\log x + x/y} \Rightarrow -\frac{y(y + x \log y)}{x(x + y \log x)}$$

Example 5: If $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$ then $\frac{dy}{dx}$ equals –

Sol: Use substitution for x and y .

Putting $x = a \sin A$, $y = a \sin B$, then given relation becomes

$$\cos A + \cos B = a(\sin A - \sin B)$$

$$\Rightarrow 2a \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$= 2a \cos\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

$$\text{Divide and multiply by } \cos\left(\frac{A+B}{2}\right)$$

$$\Rightarrow \cot\left(\frac{A-B}{2}\right) = a \left[\because \cos\left(\frac{A+B}{2}\right) \neq 0 \right]$$

$$\Rightarrow A - B = 2 \cot^{-1}a \Rightarrow \sin^{-1}x - \sin^{-1}y = 2 \cot^{-1}a$$

$$\frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-y^2}} \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = \sqrt{\frac{1-y^2}{1-x^2}}$$

Example 6: If $x^2 + y^2 = t - \frac{1}{t}$, $x^4 + y^4 = t^2 + \frac{1}{t^2}$ then $\frac{dy}{dx}$ equals

Sol: Eliminate t from the first and the second equation and then find the derivative.

Squaring the first equation, we have

$$x^4 + y^4 + 2x^2y^2 = t^2 + \frac{1}{t^2} - 2$$

$$\Rightarrow t^2 + \frac{1}{t^2} + 2x^2y^2 = t^2 + \frac{1}{t^2} - 2 \quad (\text{from second equation})$$

$$\Rightarrow x^2y^2 = -1 \Rightarrow y^2 = -\frac{1}{x^2}$$

$$\therefore 2y \frac{dy}{dx} = \frac{2}{x^3} \Rightarrow \frac{dy}{dx} = \frac{1}{x^3y}$$

Example 7: The derivation of

$$\sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right) \text{ w.r.t. } \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) \text{ is}$$

Sol: Differentiation w.r.t another function.

Putting $x = \tan \theta$

$$y = \sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right) = \sin^{-1}\left(\frac{\tan \theta}{\sec \theta}\right) = \theta$$

$$\& z = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) = \cos^{-1}\left(\frac{1-\tan^2 \theta}{1+\tan^2 \theta}\right) = 2\theta$$

\therefore Required derivative = $\frac{1}{2}$

$$\Rightarrow \frac{dy}{dz} = \frac{\frac{dy}{d\theta}}{\frac{dz}{d\theta}} = \frac{\theta}{2\theta} = \frac{1}{2}$$

Example 8: If $y = \sin^{-1}(\sqrt{\sin x})$ then $\frac{dy}{dx}$ equals–

Sol: Differentiation of function

$$\begin{aligned} \frac{dy}{dx} &= \frac{1}{\sqrt{1-\sin x}} \cdot \frac{1}{2\sqrt{\sin x}} \cdot \cos x \\ &= \frac{\sqrt{1+\sin x}}{2\sqrt{\sin x}} = \frac{1}{2} \sqrt{1+\operatorname{cosec} x} \end{aligned}$$

Example 9: If $\sqrt{\frac{v}{\mu}} + \sqrt{\frac{\mu}{v}} = 6$ then $\frac{dv}{d\mu} = ?$

Sol: Square the given equation and proceed.

$$\sqrt{\frac{v}{\mu}} + \sqrt{\frac{\mu}{v}} = 6 \Rightarrow \frac{v}{\mu} + \frac{\mu}{v} + 2 = 36$$

$$\Rightarrow \mu^2 + v^2 = 34 \mu v$$

Differentiating both sides w.r.t. μ we have

$$2\mu + 2v \frac{dv}{d\mu} = 34 v + 34 \mu \frac{dv}{d\mu}$$

$$\Rightarrow 2[17\mu - v] \frac{dv}{d\mu} = 2[\mu - 17v] \therefore \frac{dv}{d\mu} = \frac{\mu - 17v}{17\mu - v}$$

Example 10: If $x = \exp. \tan^{-1}\left(\frac{y-x^2}{x^2}\right)$, then $\frac{dy}{dx}$ equals

Sol: Simplify the given equation and differentiate.

Taking log on both sides, we get

$$\log x = \tan^{-1}\left(\frac{y-x^2}{x^2}\right)$$

$$\Rightarrow \tan(\log x) = (y - x^2) / x^2 \Rightarrow y = x^2 + x^2 \tan(\log x)$$

$$\therefore \frac{dy}{dx} = 2x + 2x \tan(\log x) + x \sec^2(\log x)$$

$$\Rightarrow 2x [1 + \tan(\log x)] + x \sec^2(\log x)$$

Example 11: Find $\frac{d}{dx} \cos^{-1} \left(\frac{4x^3}{27} - x \right)$

Sol: Let $y = \cos^{-1} \left(\frac{4x^3}{27} - x \right) = \cos^{-1} \left(4 \left(\frac{x}{3} \right)^3 - 3 \left(\frac{x}{3} \right) \right)$

$$\frac{x}{3} = \cos \theta \Rightarrow \theta = \cos^{-1} \left(\frac{x}{3} \right)$$

$$\therefore y = \cos^{-1} (4 \cos^3 \theta - 3 \cos \theta) = \cos^{-1} (\cos 3\theta) = 3\theta$$

$$\therefore y = 3 \cos^{-1} \left(\frac{x}{3} \right)$$

$$\therefore \frac{dy}{dx} = 3 \cdot \frac{-1}{\sqrt{1 - (x^2/9)}} \cdot \frac{1}{3} = \frac{-3}{\sqrt{9 - x^2}}$$

Example 12: If $\cos^{-1} \left(\frac{x^2 - y^2}{x^2 + y^2} \right) = \log a$ then $\frac{dy}{dx} =$

Sol: Take cosine on both sides and then apply componendo and dividendo.

$$\cos^{-1} \left(\frac{x^2 - y^2}{x^2 + y^2} \right) = \log a$$

$$\Rightarrow \frac{x^2 - y^2}{x^2 + y^2} = \cos(\log a) = k \text{ (say)}$$

by componendo and dividends,

$$\therefore \frac{(x^2 - y^2) + (x^2 + y^2)}{(x^2 - y^2) - (x^2 + y^2)} = \frac{k+1}{k-1}$$

$$\Rightarrow \frac{2x^2}{-2y^2} = \frac{k+1}{k-1} \quad \therefore \frac{x}{y} = \sqrt{\frac{k+1}{k-1}}$$

Differentiating both sides w.r.t. 'x' we get

$$\frac{1}{y} - \frac{x}{y^2} \frac{dy}{dx} = 0 \quad \Rightarrow \frac{dy}{dx} = \frac{y}{x}$$

Example 13: If $y^2 = p(x)$ is a polynomial of degree 3,

then $2 \frac{d}{dx} \left(y^3 \frac{d^2 y}{dx^2} \right)$ is equal to

Sol: Find first order, second order and third order derivative of $p(x)$.

$$p'(x) = 2yy' \Rightarrow p''(x) = 2yy'' + 2y'^2 \Rightarrow p'''(x) = 2yy''' + 4y'y''$$

$$\text{Also } 2 \frac{d}{dx} \left(y^3 \frac{d^2 y}{dx^2} \right) = 2 \frac{d}{dx} (y^3 y'')$$

$$= 2[y^3 y''' + 3y'^2 y''] = y^2 [2yy''' + 6y'y''] = p(x) p'''(x)$$

Example 14: If the tangent at the point $P(at^2, at^3)$ on the curve $ay^2 = x^3$ intersects the curve again at the point Q, find the point Q.

Sol: Solve the equation of the tangent and the equation of the curve.

$$ay^2 = x^3 \Rightarrow 2ay \frac{dy}{dx} = 3x^2$$

$$\text{Slope of tangent at P is } \left(\frac{3x^2}{2ay} \right)_P = \frac{3a^2 t^4}{2a^2 t^3} = \frac{3}{2}t$$

Let Q be (at_1^2, at_1^3) . Slope of line

$$PQ = \frac{at_1^3 - at^3}{at_1^2 - at^2} = \frac{t_1^2 + tt_1 + t^2}{t_1 + t}$$

which must be the slope of tangent at P. Hence,

$$\frac{t_1^2 + tt_1 + t^2}{t_1 + t} = \frac{3t}{2} \Rightarrow 2t_1^2 - tt_1 - t^2 = 0$$

$$\Rightarrow (t_1 - t)(2t_1 + t) = 0 \Rightarrow t_1 = -\frac{t}{2}$$

$$\text{Thus, Q has coordinates } \left(\frac{at^2}{4}, -\frac{at^3}{8} \right)$$

Example 15: Show that the curves $ax^2 + by^2 = 1$ and $cx^2 + dy^2 = 1$ cut orthogonally if, $\frac{1}{a} - \frac{1}{b} = \frac{1}{c} - \frac{1}{d}$

Sol: Equate the product of $\frac{dy}{dx}$ from the two equations to -1.

Let the two curves cut each other at the point (x_1, y_1) ; then

$$ax_1^2 + by_1^2 = 1 \quad \dots (i)$$

$$\& cx_1^2 + dy_1^2 = 1 \quad \dots (ii)$$

From (i) and (ii), we get

$$= (a - c)x_1^2 + (b - d)y_1^2 = 0 \quad \dots (iii)$$

Slope of the tangent to the curve

$$ax^2 + by^2 = 1, \text{ at } (x_1, y_1) \text{ is given by, } \left[\frac{dy}{dx} \right]_{(x_1, y_1)} = -\frac{ax_1}{by_1}$$

Slope of the tangent to the curve

$$cx^2 + dy^2 = 1, \text{ at } (x_1, y_1) \text{ is given by,}$$

$$\left[\frac{dy}{dx} \right]_{(x_1, y_1)} = - \frac{cx_1}{dy_1}$$

If the two curves cut orthogonally, we must have,

$$\left(-\frac{ax_1}{by_1} \right) \left(-\frac{cx_1}{dy_1} \right)$$

$$\Rightarrow acx_1^2 + bdy_1^2 = 0 \quad \dots (iv)$$

From (iii) and (iv), we have

$$\frac{a-c}{ac} = \frac{b-d}{bd} \Rightarrow \frac{1}{a} - \frac{1}{b} = \frac{1}{c} - \frac{1}{d}$$

Example 16: Find the acute angle between the curves $y = |x^2 - 1|$ and $y = |x^2 - 3|$ at their points of intersection when $x > 0$.

Sol: Solve the two curves and find the slope for the two tangents. Proceed to find the angle between the two lines.

For the intersection of the given curves

$$|x^2 - 1| = |x^2 - 3| \Rightarrow (x^2 - 1)^2 = (x^2 - 3)^2$$

$$\Rightarrow (x^2 - 1)^2 - (x^2 - 3)^2 = 0$$

$$\Rightarrow [(x^2 - 1) - (x^2 - 3)] [(x^2 - 1) + (x^2 - 3)] = 0$$

$$\Rightarrow [2x^2 - 4] = 0 \Rightarrow 2x^2 = 4 \Rightarrow x = \pm\sqrt{2}$$

neglecting $x = -\sqrt{2}$ as $x > 0$

We have point of intersection as $x = \sqrt{2}$

Here $y = |x^2 - 1| = (x^2 - 1)$ in the neighbourhood of

$x = \sqrt{2}$ and $y = -(x^2 - 3)$ in the neighbourhood of $x = \sqrt{2}$

$$\Rightarrow \left(\frac{dy}{dx} \right)_{c_1} = 2x = 2\sqrt{2} \quad \text{and} \quad \left(\frac{dy}{dx} \right)_{c_2} = -2x = -2\sqrt{2}$$

Hence, if θ is angle between them,

$$\Rightarrow \tan \theta = \left| \frac{2\sqrt{2} - (-2\sqrt{2})}{1 + 2\sqrt{2}(-2\sqrt{2})} \right| = \left| \frac{4\sqrt{2}}{-7} \right| = \left(\frac{4\sqrt{2}}{7} \right)$$

$$\therefore \theta = \tan^{-1} \left(\frac{4\sqrt{2}}{7} \right)$$

Example 17: At what points on the curve $y = \frac{2}{3}x^3 + \frac{1}{2}x^2$, then tangent make equal angles with coordinate axes.

Sol.: Find dy/dx and equate it to ± 1 .

$$\text{Given curve is } y = \frac{2}{3}x^3 + \frac{1}{2}x^2 \quad \dots (i)$$

Differentiating both sides w.r.t. x , then $\frac{dy}{dx} = 2x^2 + x$

\therefore Tangents make equal angles with coordinate axes.

$$\therefore \frac{dy}{dx} = \pm 1 \text{ or } 2x^2 + x = \pm 1 \text{ or}$$

$$2x^2 + x + 1 \neq 0 \text{ and } 2x^2 + x - 1 = 0$$

$$\text{or } 2x^2 + 2x - x - 1 = 0$$

(If $2x^2 + x + 1 = 0$ then x is imaginary)

$$\text{or } (2x - 1)(x + 1) \therefore x = \frac{1}{2}, -1$$

$$\text{From (1), } x = \frac{1}{2}, y = \frac{2}{3} \cdot \frac{1}{8} + \frac{1}{2} \cdot \frac{1}{4} = \frac{5}{24}$$

$$\text{and for } x = -1, y = -\frac{2}{3} + \frac{1}{2} = -\frac{1}{6}$$

$$\text{hence point are } \left(\frac{1}{2}, \frac{5}{24} \right) \text{ and } \left(-1, -\frac{1}{6} \right)$$

Example 18: The side of the rectangle of the greatest area, that can be inscribed in the ellipse $x^2 + 2y^2 = 8$, are given by


$$(A) 4\sqrt{2}, 4 \quad (B) 4, 2\sqrt{2}$$

$$(C) 2, \sqrt{2} \quad (D) 2\sqrt{2}, 2$$

Sol: (B) Consider a point on the ellipse and write the expression for the area of the rectangle. Then find the maximum area using first and second order derivative.

$$\text{Any point on the ellipse } \frac{x^2}{8} + \frac{y^2}{4} = 1 \text{ is}$$

$$(2\sqrt{2} \cos \theta, 2 \sin \theta) \text{ [see figure]}$$



A = area of the inscribed rectangle

$$= 4(2\sqrt{2} \cos \theta)(2 \sin \theta) = 8\sqrt{2} \sin 2\theta$$

$$\frac{dA}{d\theta} = 16\sqrt{2} \cos 2\theta = 0 \Rightarrow \theta = \frac{\pi}{4}$$

$$\text{Also } \frac{d^2A}{d\theta^2} = -32\sqrt{2} \sin 2\theta < 0 \text{ for } \theta = \frac{\pi}{4}$$

Hence, the inscribed rectangle is of largest area if the

sides are $4\sqrt{2} \cos \frac{\pi}{4}$ and $4 \sin \left(\frac{\pi}{4} \right)$ i.e. 4 and $2\sqrt{2}$.

JEE Main/Boards

Exercise 1

Methods of Differentiation

Q.1 Find the derivative of $e^{\sqrt{x+3}}$, with respect to x .

Q.2 Differentiate, $\sin(\log x)$, with the respect to x .

Q.3 If $x = \sin\theta$, $y = -\tan\theta$, find dy/dx .

Q.4 Differentiate, $\cos^{-1}(\sqrt{x})$, with the respect to x .

Q.5 Differentiate, $e^{\tan^{-1}x}$, with the respect to x .

Q.6 Differentiate, $\sin\{\log(x^3 - 1)\}$, with the respect to x .

Q.7 Differentiate, $\cos x$, with the respect to e^x .

Q.8 Differentiate the following w.r.t., $x : \log_2(\sin x)$.

Q.9 Differentiate the following w.r.t., $x : y = 5^{\log(\sin x)}$.

Q.10 Find $\frac{dy}{dx}$, when $\sqrt{x} + \sqrt{y} = 5$ at $(4, 9)$.

Q.11 $y = \cot^{-1}\left(\frac{1+x}{1-x}\right)$

Q.12 $y = \sqrt{\frac{1+\tan x}{1-\tan x}}$

Q.13 $y = \sin\left[\sqrt{\cos\sqrt{x}}\right]$

Q.14 $y = \tan^{-1}\left(\frac{\sqrt{x} + \sqrt{a}}{1 - \sqrt{ax}}\right)$

Q.15 $y = (\sin x)^{\cos^{-1}x}$

Q.16 $y = \cos^{-1}\left[(2 \cos x + 3 \sin x)\sqrt{13}\right]$

Q.17 $y = \sec^{-1}\left(\frac{1+x^2}{1-x^2}\right)$

Q.18 $y = \sin^{-1}\left[\frac{\sqrt{1+x} + \sqrt{1-x}}{2}\right]$

Q.19 $y = \sin^{-1}\left[2ax\sqrt{1-a^2x^2}\right]$

Q.20 $y = \sqrt{a + \sqrt{a+x}}$

Q.21 $y = \tan^{-1}\left[\frac{ax-b}{a+bx}\right]$

Q.22 $y = \log\left(x + \sqrt{x^2 + a^2}\right)$

Q.23 $y = \log\left(\sin\sqrt{1+x^2}\right)$

Application of Derivatives

Q.1 Find the point on the curve $y = x^2 - 4x + 5$, where tangent to the curve is parallel to x -axis.

Q.2 If two curves cut orthogonally, then what can we say about the angle between tangents at the point of intersection of the curves.

Q.3 Find the slopes of tangent and normal to the curve $f(x) = 3x^2 - 5$ at $x = \frac{1}{2}$.

Q.4 If the tangent of the curve $y = f(x)$ at point (x, y) on the curve is parallel to y -axis, then what is the value of $\frac{dy}{dx}$.

Q.5 Find a point on the curve $y = 2x^2 - 6x - 4$ at which the tangent is parallel to x -axis.

Q.6 Find a point on the curve $y = x^2 - 4x - 32$ at which the tangent is parallel to x -axis.

Q.7 Find the equations of tangent and normal to the curve $y = \sqrt[3]{5-x}$ at $(-3, 2)$.

Q.8 Find equations of tangent to the curve $y = \sqrt{4x-3}$, if parallel to x -axis.

Q.9 Verify that the point (1, 1) is a point of intersection of the curves $x^2 = y$ and $x^3 + 6y = 7$ and show that these curves cut orthogonally at this point.

Q.10 Find the equation of tangent to the parabola $y^2 = 8x$ which is parallel to line $4x - y + 3 = 0$.

Q.11 Find the equation of tangent to the curve $y = -5x^2 + 6x + 7$ at the point $\left(\frac{1}{2}, \frac{35}{4}\right)$.

Q.12 Find the equation of tangent to the curve $xy = c^2$ at the point $\left(\frac{c}{k}, ck\right)$ on it.

Q.13 Prove that the tangents to the curve $y = x^3 + 6$ at the points $(-1, 5)$ and $(1, 7)$ are parallel.

Q.14 At what point on the curve $y = x^2$ does the tangent make an angle of 45° with x-axis?

Q.15 Find the point (s) on the curve $\frac{x^2}{9} + \frac{y^2}{4} = 1$ parallel to y-axis.

Q.16 Find the slope of the normal to the curve $x = \frac{1}{t}$ $y = 2t$ at $t = 2$.

Q.17 Show that equation of the tangent to the curve $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at (x_0, y_0) is $\frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} = 1$.

Q.18 Find the equation of the normal lines to the curve $y = 4x^3 - 3x + 5$ which are parallel to the line $9y + x + 3 = 0$.

Q.19 Find the equation of normal line to the curve $y(x - 2)(x - 3) - x = 7 = 0$ at the point where it meets x-axis.

Q.20 Find the equation of tangent to the curve $\sqrt{x} + \sqrt{y} = \sqrt{a}$ at the point (x_1, y_1) and show that the sum of its intercepts on axis is constant.

Q.21 Find the equation of the normals to the curve $3x^2 + y^2 = 8$ parallel to the line $x + 3y = 4$.

Q.22 Find the equation of the tangents to the curve $\sqrt{x} + \sqrt{y} = \sqrt{a}$ at the point $\left(\frac{a^2}{4}, \frac{a^2}{4}\right)$.

Q.23 Find the points on the curve $y = x^3 - 2x^2 - 2x$ at which the tangent lines are parallel to the line $y = 2x - 3$.

Q.24 Find the angle between the parabolas $y^2 = 4ax$ and $x^2 = 4by$ at their point of intersection other than the origin.

Exercise 2

Methods of Differentiation

Single Correct Choice Type

Q.1 If $y = f\left(\frac{3x+4}{5x+6}\right)$ & $f'(x) = \tan x^2$ then $\frac{dy}{dx} =$

(A) $\tan x^3$

(B) $-2 \tan \left[\frac{3x+4}{5x+6} \right]^2 \cdot \frac{1}{(5x+6)^2}$

(C) $f\left(\frac{3 \tan x^2 + 4}{5 \tan x^2 + 6}\right) \tan x^2$

(D) None

Q.2 Let g is the inverse function of f & $f'(x) = \frac{x^{10}}{(1+x^2)}$. If $g(2) = a$ then $g'(2)$ is equal to

(A) $\frac{5}{2^{10}}$ (B) $\frac{1+a^2}{a^{10}}$ (C) $\frac{a^{10}}{1+a^2}$ (D) $\frac{1+a^{10}}{a^2}$

Q.3 If $y = \frac{1}{1+x^{n-m}+x^{p-m}} + \frac{1}{1+x^{m-n}+x^{p-n}} + \frac{1}{1+x^{m-p}+x^{n-p}}$

Then $\frac{dy}{dx}$ at e^{mnp} is equal to :

(A) e^{mnp} (B) $e^{mn/p}$ (C) $e^{np/m}$ (D) None

Q.4 Let f is differentiable in $(0, 6)$ & $f'(4) = 5$ then $\lim_{x \rightarrow 2} \frac{f(4) - f(x^2)}{2-x} =$

(A) 5 (B) $5/4$ (C) 10 (D) 20

Q.5 Let $\ell = \lim_{x \rightarrow 0} x^m (\ln x)^n$ where $m, n \in \mathbb{N}$ then

(A) ℓ is independent of m and n

(B) ℓ is independent of m and depend on m

(C) ℓ is independent of n and depend on m

(D) ℓ is dependent on both m and n

Q.6 Let $f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2\sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$ then $\lim_{x \rightarrow 0} \frac{f'(x)}{x} =$

(A) 2 (B) -2 (C) -1 (D) 1

Q.7 Let $f(x) = \begin{vmatrix} \cos x & \sin x & \cos x \\ \cos 2x & \sin 2x & 2\cos 2x \\ \cos 3x & \sin 3x & 3\cos 3x \end{vmatrix}$ then $f'\left(\frac{\pi}{2}\right) =$

(A) 0 (B) -12 (C) 4 (D) 1

Q.8 If $y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$, then dy/dx at $x = \pi/2$ is

(A) $\frac{-8}{\pi^2 + 4}$ (B) $\frac{4}{\pi^2 + 4}$

(C) $\frac{8}{\pi^2 + 4}$ (D) Does not exists

Q.9 If $f(4) = g(4) = 2$; $f'(4) = 9$; $g'(4) = 6$ then

$\lim_{x \rightarrow 4} \frac{\sqrt{f(x)} - \sqrt{g(x)}}{\sqrt{x} - 2}$ is equal to :

(A) $3\sqrt{2}$ (B) $\frac{3}{\sqrt{2}}$ (C) 0 (D) None

Q.10 If $y = x + e^x$ then $\frac{d^2x}{dy^2}$ is :

(A) e^x (B) $-\frac{e^x}{(1+e^x)^3}$

(C) $-\frac{e^x}{(1+e^x)^2}$ (D) $\frac{-1}{(1+e^x)^3}$

Q.11 If f is twice differentiable such that $f''(x) = -f(x)$, $f'(x) = g(x)$ $h'(x) = [f(x)]^2 + [g(x)]^2$ and $h(0) = 2$, $h(1) = 4$, then the equation $y = h(x)$ represents :

- (A) A curve of degree 2
 (B) A curve passing through the origin
 (C) A straight line with slope 2
 (D) A straight line with y intercept equal to -2

Q.12 Let $f(x) = x + 3 \ln(x-2)$ & $g(x) = x + 5 \ln(x-1)$, then the set of x satisfying the inequality $f'(x) < g'(x)$ is

(A) $\left(2, \frac{7}{2}\right)$ (B) $(1, 2) \cup \left(-\frac{7}{2}, \infty\right)$

(C) $(2, \infty)$ (D) $\left(\frac{7}{2}, \infty\right)$

Q.13 Let $f(x) = \sin x$; $g(x) = x^2$ & $h(x) = \log_e x$ & $f(x) = h[g(f(x))]$ then $\frac{df(x)}{dx^2}$ is equal to :

(A) $2 \operatorname{cosec}^3 x$ (B) $2 \cos t(x^2) - 4x^2 \operatorname{cosec}^2(x^2)$

(C) $2x \cot x^2$ (D) $-2 \operatorname{cosec}^2 x$

Q.14 Let $f(x) = x^n$, n being a non-negative integer. The number of value of n for which $f'(p+q) = f' \frac{b}{ab+2ay}$ $(p) + f'(q)$ is valid for all $p, q > 0$ is :

(A) 0 (B) 1 (C) 2 (D) None of these

Q.15 If $f(x) = \prod_{n=1}^{100} (x-n)^{n(101-n)}$; then $\frac{f(101)}{f'(101)} =$

(A) 5050 (B) $\frac{1}{5050}$ (C) 10010 (D) $\frac{1}{10010}$

Q.16 Let $f(x) = \begin{cases} \frac{3x^2 + 2x - 1}{6x^2 - 5x + 1} & \text{for } x \neq \frac{1}{3} \\ -4 & \text{for } x = \frac{1}{3} \end{cases}$ then $f'\left(\frac{1}{3}\right)$

(A) is equal to -9 (B) is equal to -27

(C) is equal to 27 (D) does not exist

Q.17 Let $f(x)$ be a quadratic expression which is positive for all real x . If $g(x) = f(x) + f'(x) + f''(x)$, then for any real x , which one is correct.

(A) $g(x) < 0$ (B) $g(x) > 0$ (C) $g(x) = 0$ (D) $g(x) \geq 0$

Q.18 If $y = \frac{x^4 + 4}{x^2 - 2x + 2}$ then $\left. \frac{dy}{dx} \right|_{x=1/2}$ is :

(A) 3 (B) -1 (C) 4 (D) None

Q.19 A function f , defined for all positive real numbers, satisfies the equation $f(x^2) = x^3$ for every $x > 0$. Then the value of $f'(4)$

(A) 12 (B) 3 (C) $3/2$ (D) Cannot be determined

Q.20 If $x = \sin t$ and $y = \sin 3t$, then the value of 'K' for which $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + Ky = 0$ is

(A) 3 (B) 6 (C) 12 (D) 9

Q.21 If $x = \ln t$ & $y = t^2 - 1$ then $y''(1)$ at $t = 1$ is

- (A) 2 (B) 4 (C) 3 (D) None

Application of Derivatives

Single Correct Choice Type

Q.1 The angle at which the curve $y = ke^{kx}$ intersects the y-axis is

- (A) $\tan^{-1}k^2$ (B) $\cot^{-1}(k^2)$
(C) $\sec^{-1}(\sqrt{1+k^4})$ (D) None

Q.2 The angle between the tangent lines to the graph of the function $f(x) = \int_0^x (2t - 5) dt$ at the point where the graph cuts the x-axis is -

- (A) $\pi/6$ (B) $\pi/4$ (C) $\pi/3$ (D) $\pi/2$

Q.3 If a variable tangent to the curve $x^2y = c^3$ makes intercepts a, b on x and y axis respectively then the value of a^2b is

- (A) $27c^3$ (B) $\frac{4}{27}c^3$ (C) $\frac{27}{4}c^3$ (D) $\frac{4}{9}c^3$

Q.4 Consider the function $f(x) = \begin{cases} x \sin \frac{\pi}{x} & \text{for } x > 0 \\ 0 & \text{for } x = 0 \end{cases}$ then the number of points in $(0, 1)$ where the derivative

$f'(x)$ vanishes, is

- (A) 0 (B) 1 (C) 2 (D) infinite

Q.5 The tangent to the graph of the function $y = f(x)$ at the point with abscissa $x = a$ forms with the x-axis an angle of $\pi/3$ and at the point with abscissa $x = b$ at an angle of $\pi/4$, then the value of the integral, $\int_a^b f'(x) \cdot f''(x) dx$ is equal to

- (A) 1 (B) 0 (C) $-\sqrt{3}$ (D) -1

[assume $f''(x)$ to be continuous]

Q.6 Let C be the curve $y = x^3$ (where x takes all real values). The tangent at A meets the curve again at B . If the gradient at B is K times the gradient at A then K is equal to

- (A) 4 (B) 2 (C) -2 (D) $1/4$

Q.7 The subnormal at any point on the curve $xy^n = a^{n+1}$ is constant for :

- (A) $n = 0$ (B) $n = 1$
(C) $n = -2$ (D) No value of n

Q.8 Equation of the line through the point $(1/2, 2)$ and tangent to the parabola $y = \frac{-x^2}{2} + 2$ and secant to the curve $y = \sqrt{4 - x^2}$ is

- (A) $2x + 2y - 5 = 0$ (B) $2x + 2y - 3 = 0$
(C) $y - 2 = 0$ (D) None of these

Q.9 Two curves $C_1: y = x^2 - 3$ and $C_2: y = kx^2$, $k \in \mathbb{R}$ intersect each other at two different point. The tangent drawn to C_2 at one of the point of intersection $A \equiv (a, y_1)$, ($a > 0$) meets C_1 again at $B(1, y_2)$ ($y_1 \neq y_2$). The value of ' a ' is

- (A) 4 (B) 3 (C) 2 (D) 1

Q.10 Number of roots of the equation $x^2 \cdot e^{2-x} = 1$ is:

- (A) 2 (B) 4 (C) 6 (D) Zero

Q.11 The x-intercept of the tangent at any arbitrary point of the curve $\frac{a}{x^2} + \frac{b}{y^2} = 1$ is proportional to

- (A) Square of the abscissa of the point of tangency
(B) Square root of the abscissa of the point of tangency
(C) Cube of the abscissa of the point of tangency
(D) Cube root of the abscissa of the point of tangency

Q.12 The line which is parallel to x-axis and crosses the curve $y = \sqrt{x}$ at an angle of $\frac{\pi}{4}$ is

- (A) $y = -1/2$ (B) $x = 1/2$ (C) $y = 1/4$ (D) $y = 1/2$

Q.13 The lines tangent to the curves $y^3 - x^2y + 5y - 2x = 0$ and $x^4 - x^3y^2 + 5x + 2y = 0$ at the origin intersect at an angle θ equal to

- (A) $\pi/6$ (B) $\pi/4$ (C) $\pi/3$ (D) $\pi/2$

Q.14 Consider $f(x) = \int_0^x \left(t + \frac{1}{t} \right) dt$ and $g(x) = 'f'$ for $x \in \left[\frac{1}{2}, 3 \right]$

If P is a point on the curve $y = g(x)$ such that the tangent to this curve at P is parallel to a chord joining the points $\left(\frac{1}{2}, g\left(\frac{1}{2}\right)\right)$ and $(3, g(3))$ of the curve, then the coordinates of the point P

- (A) can't be found out (B) $\left(\frac{7}{4}, \frac{65}{28}\right)$
 (C) (1, 2) (D) $\left(\sqrt{\frac{3}{2}}, \frac{5}{\sqrt{6}}\right)$

Q.15 The co-ordinates of the point on the curve $9y^2 = x^3$ where the normal to the curve makes equal intercepts with the axes is

- (A) $\left(1, \frac{1}{3}\right)$ (B) $(3, \sqrt{3})$ (C) $\left(4, \frac{8}{3}\right)$ (D) $\left(\frac{6}{5}, \frac{2}{5}\sqrt{\frac{6}{5}}\right)$

Previous Years' Questions

Q.1 The normal to the curve $x = a(\cos \theta + \theta \sin \theta)$, $y = a(\sin \theta - \theta \cos \theta)$ at any / point " θ " is such that **(1983)**

- (A) It makes a constant angle with the x-axis
 (B) It passes through the origin
 (C) It is at a constant distance from the origin
 (D) None of the above

Q.2 The slope of tangent to a curve $y = f(x)$ at $[x, f(x)]$ is $2x + 1$. If the curve passes through the point (1, 2), then the area bounded by the curve, the x-axis and the line $x = 1$ is **(1995)**

- (A) 5/6 (B) 6/5 (C) 1/6 (D) 6

Q.3 If the normal to the curve $y = f(x)$ at the point (3, 4) makes an angle $\frac{3\pi}{4}$ with the positive x-axis, then $f'(3)$ is equal to **(2000)**

- (A) -1 (B) -3/4 (C) 4/3 (D) 1

Q.4 The point(s) on the curve $y^3 + 3x^2 = 12y$ where the tangent is vertical, is (are) **(2002)**

- (A) $\left(\pm \frac{4}{\sqrt{3}}, -2\right)$ (B) $\left(\pm \sqrt{\frac{11}{3}}, 0\right)$
 (C) (0, 0) (D) $\left(\pm \frac{4}{\sqrt{3}}, 2\right)$

Q.5 The equation of the common tangent to the curves $y^2 = 8x$ and $xy = -1$ is **(2002)**

- (A) $3y = 9x + 2$ (B) $y = 2x + 1$
 (C) $2y = x + 8$ (D) $y = x + 2$

Q.6 Tangents are drawn to the ellipse $x^2 + 2y^2 = 2$, then the locus of the mid point of the intercept made by the tangents between the coordinate axes is **(2004)**

- (A) $\frac{1}{2x^2} + \frac{1}{4y^2} = 1$ (B) $\frac{1}{4x^2} + \frac{1}{2y^2} = 1$
 (C) $\frac{x^2}{2} + \frac{y^2}{4} = 1$ (D) $\frac{x^2}{4} + \frac{y^2}{2} = 1$

Q.7 The angle between the tangent drawn from the point (1, 4) to the parabola $y^2 = 4x$ is **(2004)**

- (A) $\frac{\pi}{6}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{2}$

Q.8 The tangent at (1, 7) to the curve $x^2 - y - 6$ touches the circle $x^2 + y^2 + 16x + 12y + c = 0$ at **(2005)**

- (A) (6, 7) (B) (-6, 7) (C) (6, -7) (D) (-6, -7)

Q.9 The tangent to the curve $y = e^x$ drawn at the point (c, e^c) intersects the line joining the point $(c-1, e^{c-1})$ and $(c+1, e^{c+1})$ **(2007)**

- (A) On the left of $x = c$ (B) On the right of $x = c$
 (C) At no point (D) At all points

Q.10 Let $f(x) = \begin{cases} (x-1)\sin\left(\frac{1}{x-1}\right), & \text{if } x \neq 1 \\ 0, & \text{if } x = 1 \end{cases}$.

Then which one of the following is true ? **(2008)**

- (A) f is neither differentiable at $x = 0$ not at $x = 1$
 (B) f is differentiable at $x = 0$ and at $x = 1$
 (C) f is differentiable at $x = 0$ but not at $x = 1$
 (D) f is differentiable at $x = 1$ but not at $x = 0$

Q.11 The area of the plane region bounded by the curves $x + 2y^2 = 0$ and $x + 3y^2 = 1$ is equal to **(2008)**

- (A) $\frac{5}{3}$ (B) $\frac{1}{3}$ (C) $\frac{2}{3}$ (D) $\frac{4}{3}$

JEE Advanced/Boards

Exercise 1

Methods of Differentiation

Q.1 Let $y = x \sin kx$. Find the possible value of k for which the differential equation $\frac{d^2y}{dx^2} + y = 2k \cos kx$ holds true for all $x \in \mathbb{R}$.

Q.2 Find a polynomial function $f(x)$ such that $f(2x) = f(x) f''(x)$.

Q.3 Let f and g be two real-valued differentiable function on \mathbb{R} If $f'(x) = g(x)$ and $g'(x) = f(x)$ " $x \in \mathbb{R}$ and $f(3) = 5$, $f'(3) = 4$ then find the value of $(f^2(\pi) - g^2(\pi))$.

Q.4 Find the value of the expression $y^3 \frac{d^2y}{dx^2}$ on the ellipse $3x^2 + 4y^2 = 12$.

Q.5 The function $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $f(x^2) \cdot f''(x) = f'(x) \cdot f'(x^2)$ for all real x . Given that $f(1) = 1$ and $f''(1) = 8$, compute the value of $f'(1) + f''(1)$.

Q.6 If $2x = y^{\frac{1}{5}} + y^{-\frac{1}{5}}$ then $(x^2 - 1)$

$\frac{d^2y}{dx^2} + x \frac{dy}{dx} = ky$, then find the value of ' k '.

Q.7 If the dependent variable y is changed to ' z ' by the substitution $y = \tan z$ then the differential equation

$\frac{d^2y}{dx^2} = 1 + \frac{2(1+y)}{1+y^2} \left(\frac{dy}{dx}\right)^2$ is changed to $\frac{d^2z}{dz^2} = \cos^2 z + k \left(\frac{dz}{dx}\right)^2$, then find the value of k .

Q.8 Show that the substitution $z = \ln \left(\tan \frac{x}{2} \right)$ changes the equation

$$\frac{d^2y}{dx^2} + \cot x \frac{dy}{dx} + 4y \operatorname{cosec}^2 x = 0$$

$$\text{to } \left(\frac{d^2y}{dz^2} \right) + 4y = 0$$

Q.9 Let $f(x) = \frac{\sin x}{x}$ if $x \neq 0$ and $f(0) = 1$. Define the function $f'(x)$ for all x and find $f''(0)$ if it exist.

Q.10 Show that $R = \frac{[1 + (dy/dx)^2]^{3/2}}{d^2y/dx^2}$ can be reduced to the form

$$R^{2/3} = \frac{1}{(d^2y/dx^2)^{2/3}} + \frac{1}{(d^2x/dy^2)^{2/3}}$$

Q.11 Suppose f and g are two functions such that $f, g :$

$\mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \ln(x + \sqrt{1+x^2})$ then find the value of $xe^{g(x)}$

$$\left(f\left(\frac{1}{x}\right) \right)' + g'(x) \text{ at } x = 1.$$

Q.12 Let $f(x)$ be a derivative function at $x = 0$ & $f\left(\frac{x+y}{k}\right) = \frac{f(x)+f(y)}{k}$ ($k \in \mathbb{R}$, $k \neq 0, 2$). Show that $f(x)$ is either a zero or an odd linear function.

$$\text{Q.13 If } f(x) = \begin{vmatrix} (x-a)^4 & (x-a)^3 & 1 \\ (x-b)^4 & (x-b)^3 & 1 \\ (x-c)^4 & (x-c)^3 & 1 \end{vmatrix} \text{ then}$$

$$f'(x) = \lambda \cdot \begin{vmatrix} (x-a)^4 & (x-a)^2 & 1 \\ (x-b)^4 & (x-b)^2 & 1 \\ (x-c)^4 & (x-c)^2 & 1 \end{vmatrix}.$$

Find the value of λ .

Q.14 Let $P(x)$ be a polynomial of degree 4 such that $P(1) = P(3) = P(5) = P'(7) = 0$. If the real number $x \neq 1, 3, 5$ is such that $P(x) = 0$ can be expressed as $x = p/q$ where ' p ' and ' q ' are relatively prime, then find $(p+q)$.

Q.15 If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function such that $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$ for all $x \in \mathbb{R}$, then prove that $f(2) = f(1) - f(0)$.

$$\text{Q.16 If } f(x) = \begin{vmatrix} \cos(x+x^2) & \sin(x+x^2) & -\cos(x+x^2) \\ \sin(x-x^2) & \cos(x-x^2) & \sin(x-x^2) \\ \sin 2x & 0 & \sin 2x^2 \end{vmatrix}$$

then find $f'(x)$

$$\text{Q.17 Let } f(x) = \begin{vmatrix} a+x & b+x & c+x \\ \ell+x & m+x & n+x \\ p+x & q+x & r+x \end{vmatrix}. \text{ Show that } f''(x) = 0$$

and that $f(x) = f(0) + kx$ where k denotes the sum of all the co-factors of the elements in $f(0)$

Q.18 If $y = \tan^{-1} \frac{1}{x^2 + x + 1} + \tan^{-1} \frac{1}{x^2 + 3x + 3} + \tan^{-1}$

$$\frac{1}{x^2 + 5x + 7} + \tan^{-1} \frac{1}{x^2 + 7x + 13} = + \dots \text{ to } n \text{ terms.}$$

Find dy/dx , expressing your answer in 2 terms.

Q.19 If $Y = sX$ and $Z = tX$, where all the letter denotes the functions of x and suffixes denotes the differentiation

w.r.t. then prove that $\begin{vmatrix} X & Y & Z \\ X_1 & Y_1 & Z_1 \\ X_2 & Y_2 & Z_2 \end{vmatrix} = X^3 \begin{vmatrix} s_1 & t_1 \\ s_2 & t_2 \end{vmatrix}$

Q.20 If $y = \tan^{-1} \frac{u}{\sqrt{1-u^2}}$ &

$$x = \sec^{-1} \frac{1}{2u^2 - 1}, u \in \left(0, \frac{1}{2}\right) \cup \left(\frac{1}{\sqrt{2}}, 1\right)$$

prove that $2 \frac{dy}{dx} + 1 = 0$

Q.21 If $y = \tan^{-1} \frac{x}{1 + \sqrt{1-x^2}} + \sin \left(2 \tan^{-1} \sqrt{\frac{1-x}{1+x}} \right)$,

The find $\frac{dy}{dx}$ for $x \in (-1, 1)$

Q.22 If $y = \cot^{-1} \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}}$,

find $\frac{dy}{dx}$ if $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right)$.

Q.23 Prove that the second order derivative of a single valued function parametrically represented by $x = \phi(t)$ and $y = \Psi(t)$, $\alpha < t < \beta$ where $\phi(t)$ and $\Psi(t)$ are differentiable functions and $\phi'(t) \neq 0$ is given

$$\text{by } \frac{d^2y}{dx^2} = \frac{\left(\frac{dx}{dt}\right)\left(\frac{d^2y}{dt^2}\right) - \left(\frac{d^2x}{dt^2}\right)\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)^3}$$

Q.24 (a) If $y = y(x)$ and it follows the relation $e^{xy} + y \cos x = 2$, then find (i) $y'(0)$ and (ii) $y''(0)$.

(b) A twice differentiable function $f(x)$ is defined for all real numbers and satisfies the following conditions $f(0) = 2$; $f'(0) = -5$ and $f''(0) = 3$

The function $g(x)$ is defined by $g(x) = e^{ax} + f(x) \forall x \in \mathbb{R}$, where 'a' is any constant. If $g'(0) + g''(0) = 0$. Find the value(s) of 'a'.

Application of derivatives

Q.1 Find the equations of the tangents drawn to the curves $y^2 - 2x^3 - 4y + 8 = 0$ from the point $(1, 2)$.

Q.2 The tangent to $y = ax^2 + bx + \frac{7}{2}$ at $(1, 2)$ is parallel to the normal at the point $(-2, 2)$ on the curve $y = x^2 + 6x + 10$. Find the value of a and b .

Q.3 Find the point of intersection of the tangents drawn to the curve $x^2y = 1 - y$ at the points where it is intersected by the curve $xy = 1 - y$.

Q.4 Find the equation of the normal to the curve $y = (1+x)^y + \sin^{-1}(\sin^2 x)$ at $x = 0$.

Q.5 A function is defined parametrically by the equation

$$f(t) = x = \begin{cases} 2t + t^2 \sin \frac{1}{t} & \text{if } t \neq 0 \\ 0 & \text{if } t = 0 \end{cases} \text{ and}$$

$$g(t) = y = \begin{cases} \frac{1}{t} \sin t^2 & \text{if } t \neq 0 \\ 0 & \text{if } t = 0 \end{cases}$$

Find the equation of the tangent and normal at the point for $t = 0$ is exist.

Q.6 A line is tangent to the curve $f(x) = \frac{41x^3}{3}$ at the point

P in the first quadrant, and has a slope of 2009. This line intersects the y-axis at $(0, b)$. Find the value of 'b'.

Q.7 Find all the tangents to the curve $y = \cos(x + y)$, $-2\pi \leq x \leq 2\pi$, that are parallel to the line $x + 2y = 0$

Q.8 There is a point (p, q) on the graph of $f(x) = x^2$ and a point (r, s) on the graph of $g(x) = \frac{-8}{x}$ where $p > 0$ and

$r > 0$. If the line through (p, q) and (r, s) is also tangent to both the curves at these points respectively, then find the value of $(p + r)$

Q.9 (i) Use differentials to approximate the values of; (a) $\sqrt{36.6}$ and (b) $\sqrt[3]{26}$.

(ii) If the radius of a sphere is measured as 9 cm with an error of 0.03 cm, then find the approximate error in calculating its volume.

Q.10 The chord of the parabola $y = -a^2x^2 + 5ax - 4$ touches the curve $y = \frac{1}{1-x}$ at the point $x = 2$ and is bisected by that point. Find 'a'.

Q.11 Tangent at a point P_1 [other than (0, 0)] on the curve $y = x^3$ meets the curve again at P_2 . The tangent at P_2 meets the curve at P_3 & so on. Show that the abscissae of $P_1, P_2, P_3, \dots, P_n$ form a GP. Also find the ratio $\frac{\text{area}(P_1P_2P_3)}{\text{area}(P_2P_3P_4)}$.

Q.12 Determine a differentiable function $y = f(x)$ which satisfies $f'(x) = [f(x)]^2$ and $f(0) = -\frac{1}{2}$. Find also the equation of the tangent at the point where the curve crosses the y-axis.

Q.13 The curve $y = ax^3 + bx^2 + cx + 5$, touches the x-axis at $P(-2, 0)$ & cuts the y-axis at a point Q where its gradient is 3. Find a, b, c

Q.14 Find the gradient of the line passing through the point (2, 8) and touching the curve $y = x^2$.

Q.15 Let $f: \{0, \infty\} \rightarrow \mathbb{R}$ be a continuous, strictly increasing function such that $f^3(x) = \int_0^x f^2(t) dt$. If a normal is drawn to the curve $y = f(x)$ with gradient $-\frac{1}{2}$, then find the intercept made by it on the y-axis.

Q.16 The graph of a certain function f contains the point (0, 2) and has the property that for each number 'p' the line tangent to $y = f(x)$ at $(p, f(p))$ intersect the x-axis at $p + 2$. Find $f(x)$

Q.17 (a) Find the value of n so that the subnormal at any point on the curve $xy^n = a^{n+1}$ may be constant (b) Show that in the curve $y = a \ln(x^2 - a^2)$ sum of the length of tangent & subtangent varies as the product of the coordinates of the point of contact

(c) If the two curve $C_1: x = y^2$ and $C_2: xy = k$ cut at right angles find the value of k .

Q.18 Let the function $f: [-4, 4] \rightarrow [-1, 1]$ be defined implicitly by the equation $x = 5y - y^5 = 0$.

Find the area of triangle formed by tangent and normal to $f(x)$ at $x = 0$ and the line $y = 5$.

Q.19 The normal at the point $P\left(2, \frac{1}{2}\right)$ on the curve $xy = 1$, meets the curve again at Q. If m is the slope of the curve at Q, then find $|m|$.

Q.20 Let C be the curve $f(x) = \ln^2 x + 2 \ln x$ and $A(a, f(a))$, $B(b, f(b))$ where $(a < b)$ are the points of tangency of two tangents drawn from origin to the curve C.

(i) Find the value of the product ab .

(ii) Find the number of values of x satisfying the equation $5x f'(x) - x \ln 10 - 10 = 0$.

Q.21 A particle moves along the curve $6y = x^3 + 2$. Find the points on the curve at which the y coordinate is changing 8 times as fast as the x coordinate

Q.22 A man 1.5 m tall walks away from a lamp post 4.5 m high at the rate of 4 km/hr.

(i) How fast is the father end of the shadow moving on the pavement?

(ii) How fast is his shadow lengthening?

Q.23 A water tank has the shape of a right circular cone with its vertex down. Its altitude is 10 cm and the radius of the base is 15 cm. Water leaks out of the bottom at a constant rate of 1 cu. cm/sec. Water is poured into the tank at a constant rate of C cu. cm/sec. Compute C so that the water level will be rising at the rate of 4 cm/sec at the instant when the water is 2 cm deep.

Q.24 Water is dripping out from a conical funnel of semi vertical angle $\frac{\pi}{4}$, at the uniform rate of 2 cm³/sec

through a tiny hole at the vertex at the bottom. When the slant height of the water is 4 cm, find the rate of decrease of the slant height of the water.

Q.25 Sand is pouring from a pipe at the rate of 12 cc/sec. The falling sand forms a cone on the ground in such a way that the height of the cone is always $\frac{1}{6}$ th of the radius of the base. How fast is the height of the sand cone increasing when the height is 4 cm?

Q.26 A circular ink blot grows at the rate of 2 cm^2 per second. Find the rate at which the radius is increasing after $2 \frac{6}{11}$ seconds. Use $\pi = \frac{22}{7}$

Q.27 A variable $\triangle ABC$ in the xy plane has its orthocentre at vertex 'B', a fixed vertex 'A' at the origin and the third vertex 'C' restricted to lie on the parabola $y = 1 + \frac{7x^2}{36}$. The point B starts at the point (0, 1) at time $t = 0$ and moves upward along the y axis at a constant velocity of 2 cm/sec . How fast is the area of the triangle increasing when $t = \frac{7}{2}$ sec.

Q.28 At time $t > 0$, the volume of a sphere is increasing at a rate proportional to the reciprocal of its radius. At $t = 0$, the radius of the sphere is 1 unit and at $t = 15$ the radius is 2 units.

- (a) Find the radius of the sphere as a function of time t
 (b) At what time t will the volume of the sphere be 27 times its volume at $t = 0$

Q.29 Water is flowing out at the rate of $6 \text{ m}^3/\text{min}$ from a reservoir shaped like a hemispherical bowl of radius $R = 13 \text{ m}$. The volume of water in the hemispherical bowl is given by $V = \frac{\pi}{3} \cdot y^2 (3R - y)$ when the water is y meter deep. Find

- (a) At what rate is the water level changing when the water is 8 m deep?
 (b) At what rate is the radius of the water surface changing when the water is 8 m deep?

Exercise 2

Methods of Differentiation

Single Correct Choice Type

Q.1 If $y = \frac{x}{a+b} \cdot \frac{x}{b+a} \cdot \frac{x}{a+b} \cdot \frac{x}{b+a} \cdot \frac{x}{a+b} \cdot \frac{x}{b+a} \dots \infty$, then $\frac{dy}{dx}$

- (A) $\frac{a}{ab+2ay}$ (B) $\frac{b}{ab+2by}$
 (C) $\frac{a}{ab+2ay}$ (D) $\frac{b}{ab+2ay}$

Q.2 The function $f(x) = e^x + x$, being differentiable and one to one to one, has a differentiable inverse $f^{-1}(x)$. The value of $\frac{d}{dx}(f^{-1})$ at the point $f(\ln 2)$ is

- (A) $\frac{1}{\ln 2}$ (B) $\frac{1}{3}$ (C) $\frac{1}{4}$ (D) None

Q.3 $f'(x) = g(x)$ and $g'(x) = -f(x)$ for all real x and $f(5) = 2 = f'(5)$ then $f(10) + g^2(10)$ is-

- (A) 2 (B) 4 (C) 8 (D) None

Q.4 Differential coefficient of

$$\left(x^{\frac{\ell+m}{m-n}}\right)^{\frac{1}{n-\ell}} \cdot \left(x^{\frac{m+n}{n-\ell}}\right)^{\frac{1}{\ell-m}} \cdot \left(x^{\frac{n+\ell}{\ell-m}}\right)^{\frac{1}{m-n}} \text{ w.r.t. is}$$

- (A) 1 (B) 0 (C) -1 (D) $x^{\ell mn}$

Q.5 Let $f(x) = (x^x)^x$ and $g(x) = x^{(x^x)}$ then:

- (A) $f'(1) = 1$ and $g'(1) = 2$ (B) $f'(1) = 2$ and $g'(1) = 1$
 (C) $f'(1) = 1$ and $g'(1) = 0$ (D) $f'(1) = 1$ and $g'(1) = 1$

Q.6 If $\frac{1}{y^m} + y^{\frac{1}{m}} = 2x$, then the value of $\frac{(x^2-1)y'' + xy'}{y}$ is equal to value equal to

- (A) 4 m^2 (B) 2 m^2 (C) m^2 (D) $-\text{m}^2$

Q.7 If $y^2 = P(x)$, is a polynomial of degree 3, then $2\left(\frac{d}{dx}\right)\left(y^3 \cdot \frac{d^2y}{dx^2}\right)$ equals:

- (A) $P'''(x) + P'(x)$ (B) $P''(x) \cdot P'''(x)$
 (C) $P(x) \cdot P'''(x)$ (D) a constant

Q.8 Given $f(x) = \frac{-x^3}{3} + x^2 \sin 1.5a - x \sin a \cdot \sin 2a - 5$ are $\sin(a^2 - 8a + 17)$ then :

- (A) $f(x)$ is not defined at $x = \sin 8$
 (B) $f'(\sin 8) > 0$
 (C) $f'(x)$ is not defined at $x = \sin 8$
 (D) $f'(\sin 8) < 0$

Q.9 If $y = \frac{\cos 6x + 6 \cos 4x + 15 \cos 2x + 10}{\cos 5x + 5 \cos 3x + 10 \cos x}$, then $\frac{dy}{dx} =$

- (A) $2 \sin x + \cos x$ (B) $-2 \sin x$
 (C) $\cos 2x$ (D) $\sin 2x$

Q.10 A curve is parametrically represented by $y = R(1 - \cos \theta)$ & $x = R(\theta - \sin \theta)$, then $\frac{d^2y}{dx^2}$ at $\theta = \pi$ is -

- (A) $-\frac{1}{2R}$ (B) $\frac{1}{4R}$ (C) $\frac{1}{2R}$ (D) $-\frac{1}{4R}$

Q.11 If $f(x) = (1 + x)^n$ then the value of $f(0) + f'(0) + \dots + \frac{f^n(0)}{n!}$ is -

- (A) n (B) 2^n (C) 2^{n-1} (D) None

Q.12 If the function $y = e^{4x} + 2e^{-x}$ is a solution of the differential equation $\frac{d^3y}{dx^3} - 13\frac{dy}{dx} = K$, then the value of K

- (A) 4 (B) 6 (C) 9 (D) 12

Q.13 $x^4 + 3x^2y^2 + 7xy^3 + 4x^3y - 15y^4 = 0$, then $\frac{d^2y}{dx^2}$ at $(1, 1)$ is -

- (A) 2 (B) 1 (C) 7 (D) 0

Q.14 If $f(x) = e^{e^x}$. Let $g(x)$ be its inverse then $g'(x)$ at $x = 2$ is -

- (A) $\frac{\ln 2}{2}$ (B) $\frac{1}{2\ln 2}$ (C) $2\ln 2$ (D) e^2

Q.15 $y = \tan^{-1}\left(\frac{1-2\ln|x|}{1+2\ln|x|}\right) + \tan^{-1}\left(\frac{3+2\ln|x|}{1-6\ln|x|}\right)$, then $\frac{d^2y}{dx^2}$ equals

- (A) 2 (B) 1 (C) 0 (D) -1

Q.16 $\lim_{x \rightarrow 0^+} (x^{x^x} - x^x)$ equals -

- (A) 1 (B) -1 (C) 0 (D) None of these

Q.17 $\lim_{x \rightarrow 0} \{(\cot x)^x + (1 - \cos x)^{\csc x}\}$ is equal to -

- (A) 2 (B) +1
(C) 0 (D) None of these

Multiple Correct Choice Type

Q.18 Let $f(x) = \tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$, then

(A) $f(x) = 112 \tan^{-1}x \Delta x \in \mathbb{R} - \{0\}$

(B) $f'(x) = \frac{1}{2(1+x^2)} \zeta x \in \mathbb{R} - \{0\}$

(C) $f(x)$ is an odd function

(D) $f(x) + f(-x) = \pi$

Q.19 If $y = \tan x \tan 2x \tan 3x$ then $\frac{dy}{dx}$ has the value to:

(A) $3 \sec^2 3x \tan x \tan 2x + \sec^2 x \tan 2x \tan 3x + 2 \sec^2 2x \tan 3x \tan x$

(B) $2y (\operatorname{cosec} 2x + 2 \operatorname{cosec} 4x + 3 \operatorname{cosec} 6x)$

(C) $3 \sec^2 3x - 2 \sec^2 2x - \sec^2 x$

(D) $\sec^2 x + 2 \sec^2 2x + 3 \sec^2 3x$

Q.20 Let $y = \sqrt{x} + \sqrt[3]{x} + \sqrt[4]{x} + \dots \infty$ then $\frac{dy}{dx}$

- (A) $\frac{1}{2y-1}$ (B) $\frac{x}{x-2y}$ (C) $\frac{1}{\sqrt{1+4x}}$ (D) $\frac{y}{2x+y}$

Q.21 If $2^x + 2^y = 2^{x+y}$ then $\frac{dy}{dx}$ has the value equal to

- (A) $-\frac{2^y}{2^x}$ (B) $\frac{1}{1-2^x}$ (C) $1-2^y$ (D) $\frac{2^x(1-2^y)}{2^y(2^x-1)}$

Q.22 If $\sqrt{y+x} + \sqrt{y-x} = c$, then $\frac{dy}{dx}$ is equal to

- (A) $\frac{2x}{c^2}$ (B) $\frac{x}{y + \sqrt{y^2 - x^2}}$


- (C) $\frac{y - \sqrt{y^2 - x^2}}{x}$ (D) $\frac{c^2}{2y}$

Application of Derivatives

Single Correct Choice Type

Q.1 The line $y = -\frac{3}{2}x$ and $y = -\frac{2}{5}x$ intersect the curve $3x^2 + 4xy + 5y^2 - 4 = 0$ at the point P and O respectively. The tangent drawn to the curve at P and Q

(A) Intersect each other at angle of 45°



- (B) Are parallel to each other
 (C) Are perpendicular to each other
 (D) None of these

Q.2 A curve is represented by the equations $x = \sec^2 t$ and $y = \cot t$ where t is a parameter. If the tangent at the point P on the curve where $t = \pi/4$ meets the curve again at the point Q then $|PQ|$ is equal to

- (A) $\frac{5\sqrt{3}}{2}$ (B) $\frac{5\sqrt{5}}{2}$
 (C) $\frac{2\sqrt{5}}{3}$ (D) $\frac{3\sqrt{5}}{2}$

Q.3 Let $f(x) = \begin{cases} x^{35} & \text{if } x \leq 1 \\ -(x-2)^3 & \text{if } x > 1 \end{cases}$ then the number of critical points on the graph of the function is

- (A) 1 (B) 2 (C) 3 (D) 4

Q.4 At any two points of the curve represented parametrically by $x = a(2 \cos t - \cos 2t)$, $y = a(2 \sin t - \sin 2t)$ the tangents are parallel to the axis of x corresponding to the values of the parameter t differing from each other by

- (A) $2\pi/3$ (B) $3\pi/4$ (C) $\pi/2$ (D) $\pi/3$

Q.5 At the point P (a, a^n) on the graph of $y = x^n$ ($n \in \mathbb{N}$) in the first quadrant a normal is drawn. The normal intersects the y -axis at the point ($0, b$). If $\lim_{a \rightarrow 0} b = \frac{1}{2}$, then n equal

- (A) 1 (B) 3 (C) 2 (D) 4

Q.6 Let $f(x) = \begin{cases} -x^2 & \text{for } x > 0 \\ x^2 + 8 & \text{for } x \leq 0 \end{cases}$. Then x intercept of the line that is tangent to the graph of $f(x)$ is

- (A) zero (B) -1 (C) -2 (D) -4

Q.7 The ordinate of all points on the curve

$$y = \frac{1}{2\sin^2 x + 3\cos^2 x} \text{ where the tangent is horizontal, is -}$$

- (A) Always equal to $1/2$
 (B) Always equal to $1/3$
 (C) $1/2$ or $1/3$ according as n is an even or an odd integer
 (D) $1/2$ or $1/3$ according as n is an odd or an even integer

Q.8 The equation of the tangent to the curve $\left(\frac{x}{a}\right)^n + \left(\frac{y}{b}\right)^n = 2$ ($n \in \mathbb{N}$) at the point with abscissa equal to 'a' can be

- (A) $\left(\frac{x}{a}\right) + \left(\frac{y}{b}\right) = 2$ (B) $\left(\frac{x}{a}\right) - \left(\frac{y}{b}\right) = 2$
 (C) $\left(\frac{x}{a}\right) - \left(\frac{y}{b}\right) = 0$ (D) $\left(\frac{x}{a}\right) + \left(\frac{y}{b}\right) = 0$

Multiple Correct Choice Type

Q.9 If $\frac{x}{a} + \frac{y}{b} = 1$ is a tangent to the curve $x = Kt$, $y = \frac{1}{t}$, $K > 0$ then

- (A) $a > 0, b > 0$ (B) $a > 0, b < 0$
 (C) $a < 0, b > 0$ (D) $a < 0, b < 0$

Q.10 The abscissa of the point on the curve $\frac{dy}{dx} = a + x$, the tangent at which cuts off equal intercepts from the co-ordinate axes is ($a > 0$)

- (A) $\frac{a}{\sqrt{2}}$ (B) $-\frac{a}{\sqrt{2}}$ (C) $a\sqrt{2}$ (D) $-a\sqrt{2}$

Q.11 The parabola $y = x^2 + px + q$ cuts the straight line $y = 2x - 3$ at a point with abscissa 1. If the distance between the vertex of the parabola and the x -axis is least then

- (A) $p = 0$ & $q = -2$
 (B) $p = -2$ & $q = 0$
 (C) least distance between the parabola and x -axis is 2
 (D) least distance between the parabola and x -axis is 1

Q.12 The co-ordinates of the point(s) on the graph

$$\text{function, } f(x) = \frac{x^3}{3} - \frac{5x^2}{2} + 7x - 4 \text{ where the tangent}$$

drawn cut off intercepts from the co-ordinate axes which are equal in magnitude but opposite in sign, is

- (A) (2, 8/3) (B) (3, 7/2)
 (C) (1, 5/6) (D) None of these

Q.13 Equation of a tangent to the curve $y \cot x = y^3 \tan x$ at the point where the abscissa is $\frac{\pi}{4}$ is

- (A) $4x + 2y = \pi + 2$ (B) $4x - 2y = \pi + 2$
 (C) $x = 0$ (D) $y = 0$

Q.14 The angle made by the tangent of the curve $x = a(t + \sin t \cos t)$; $y = a(1 + \sin t)^2$ with the x-axis at any point on it is -

- (A) $\frac{1}{4}(\pi + 2t)$ (B) $\frac{1 - \sin t}{\cos t}$ (C) $\frac{1}{4}(2t - \pi)$ (D) $\frac{1 + \sin t}{\cos 2t}$

Q.15 Consider the curve represented parametrically by the equation $x = t^3 - 4t^2 - 3t$ and $y = 2t^2 + 3t - 5$ where $t \in \mathbb{R}$

If H denotes the number of point on the curve where the tangent is horizontal and V the number of point where the tangent is vertical then

- (A) $H = 2$ and $V = 1$ (B) $H = 1$ and $V = 2$
(C) $H = 2$ and $V = 2$ (D) $H = 1$ and $V = 1$

Previous Years' Questions

Q.1 If the line $ax + bx + c = 0$ is a normal to the curve $xy = 1$, then **(1986)**

- (A) $a > 0, b > 0$ (B) $a > 0, b < 0$
(C) $a < 0, b > 0$ (D) $a < 0, b < 0$

Q.2 On the ellipse $4x^2 + 9y^2 = 1$, the point at which the tangents are parallel to the line $8x = 9y$, are **(1999)**

- (A) $\left(\frac{2}{5}, \frac{1}{5}\right)$ (B) $\left(-\frac{2}{5}, \frac{1}{5}\right)$ (C) $\left(-\frac{2}{5}, -\frac{1}{5}\right)$ (D) $\left(\frac{2}{5}, -\frac{1}{5}\right)$

Q.3 Let C be the curve $y^3 - 3xy + 2 = 0$. If H is the set of points on the curve C where the tangent is horizontal and V is the set of points on the curve C where the tangent is vertical, then $H = \dots\dots\dots$ and $V = \dots\dots\dots$ **[1997]**

Q.4 A swimmer S is in the sea at a distance d km from the closest point A on a straight shore. The house of the swimmer is on the shore at a distance L km from A. He can swim at a speed of u km/h and walk at a speed of v km/h

($v > u$). At what point on the shore should he land so that he reaches his house in the shortest possible time? **(1983)**

Q.5 Find the coordinates of the point on the curve $y = \frac{x}{1+x^2}$, where the tangent to the curve has the greatest slope. **(1997)**

Q.6 Find all the tangents to the curve $y = \cos(x + y)$, $-2\pi \leq x \leq 2\pi$, that are parallel to the line $x + 2y = 0$ **(1997)**

Q.7 Find the point on the curve $4x^2 + a^2y^2 = 4a^2$, $4 < a^2 < 8$ that is farthest from the point $(0, -2)$. **(1987)**

Q.8 Three normals are drawn from the point $(c, 0)$ to the curve $y^2 = x$. Show that c must be greater than $\frac{1}{2}$. One normal is always the x-axis. Find c for which the other two normals are perpendicular to each other **(1991)**

Q.9 What normal to the curve $y = x^2$ forms the shortest chord? **(1992)**

Q.10 Find the equation of the normal to the curve $y = (1 + x)^y + \sin^{-1}(\sin^2 x)$ at $x = 0$ **(1993)**

Q.11 Tangent at a point P_1 {other than $(0, 0)$ } on the curve $y = x^3$ meets the curve again at P_2 . The tangent at P_2 meets the curve at P_3 , and so on. Show that the abscissae of $P_1, P_2, P_3, \dots < P_n$ form a GP. Also find the ratio. $[\text{Area}(dP_1P_2P_3)] / [\text{area}(dP_2P_3P_4)]$ **(1993)**

Q.12 A rectangular sheet of fixed perimeter with sides having their lengths in the ratio 8 : 15 is converted an open rectangular box by folding after removing square of equal area from all four corners. If the total area of removed squares is 100, the resulting box has maximum volume. Then the lengths of the sides of the rectangular sheet are **(2013)**

- (A) 24 (B) 32 (C) 45 (D) 60

Q.13 Match List I with List II and select the correct answer using the code given below the lists : **(2013)**

	List - I		List - II
P	$\left(\frac{1}{y^2} \left(\frac{\cos(\tan^{-1} y) + y \sin(\tan^{-1} y)}{\cot(\sin^{-1} y) + \tan(\sin^{-1} y)} \right) + y^4 \right)^{1/2}$	1.	$\frac{1}{2} \sqrt{\frac{5}{3}}$
Q	If $\cos x + \cos y + \cos z = 0 = \sin x + \sin y + \sin z$ then possible value of $\cos \frac{x-y}{2}$ is	2.	$\sqrt{2}$

R	If $\cos\left(\frac{\pi}{4} - x\right)\cos 2x + \sin x \sin 2x \sec x = \cos x \sin 2x \sec x + \cos\left(\frac{\pi}{4} + x\right)\cos 2x \cos 2x$ then possible value of $\sec x$ is	3.	$\frac{1}{2}$
S	If $\cot\left(\sin^{-1}\sqrt{1-x^2}\right) = \sin\left(\tan^{-1}(x\sqrt{6})\right)$, $x \neq 0$, then possible value of x is	4.	1

(A) $P \rightarrow 4; Q \rightarrow 3; R \rightarrow 1; S \rightarrow 2$ (B) $P \rightarrow 4; Q \rightarrow 3; R \rightarrow 2; S \rightarrow 1$ (C) $P \rightarrow 3; Q \rightarrow 4; R \rightarrow 2; S \rightarrow 1$ (D) $P \rightarrow 3; Q \rightarrow 4; R \rightarrow 1; S \rightarrow 2$

MASTERJEE Essential Questions

JEE Main/Boards

Exercise 1

Methods of Differentiation

Q.9 Q.12 Q.15 Q.23

Application of Derivatives

Q.7 Q.11 Q.15 Q.21

Exercise 2

Methods of Differentiation

Q.2 Q.11 Q.12 Q.15
Q.17 Q.20

Application of Derivatives

Q.5 Q.7 Q.8 Q.10

Previous Years' Questions

Q.2 Q.5 Q.7 Q.9

JEE Advanced/Boards

Exercise 1

Methods of Differentiation

Q.5 Q.8 Q.11 Q.12
Q.14 Q.18 Q.23

Application of Derivatives

Q.8 Q.11 Q.15 Q.17
Q.20 Q.23 Q.24 Q.27

Exercise 2

Methods of Differentiation

Q.1 Q.3 Q.5 Q.9
Q.11 Q.15 Q.20

Application of Derivatives

Q.2 Q.4 Q.7 Q.11

Previous Years' Questions

Q.4 Q.9 Q.11

Answer Key

JEE Main/Boards

Exercise 1

Methods of Differentiation

$$\text{Q.1 } \frac{1}{2\sqrt{x+3}} e^{\sqrt{x+3}}$$

$$\text{Q.4 } \frac{-1}{2\sqrt{x}\sqrt{1-x}}$$

$$\text{Q.7 } -e^{-x} \sin x$$

$$\text{Q.10 } -\frac{3}{2}$$

$$\text{Q.13 } \frac{-\cos\sqrt{\cos\sqrt{x}} \sin\sqrt{x}}{4\sqrt{x}\sqrt{\cos\sqrt{x}}}$$

$$\text{Q.15 } (\sin x)^{\cos^{-1}x} \left[\cos^{-1}x \cdot \cot x - \frac{\log(\sin x)}{\sqrt{1+x^2}} \right]$$

$$\text{Q.18 } \frac{dy}{dx} = -\frac{1}{2\sqrt{1-x^2}}$$

$$\text{Q.21 } \frac{1}{1+x^2}$$

$$\text{Q.2 } \frac{1}{x} \cos(\log x)$$

$$\text{Q.5 } \frac{me^{m \tan^{-1}x}}{1+x^2}$$

$$\text{Q.8 } \cot x \cdot \log_2 e$$

$$\text{Q.11 } \frac{-1}{1+x^2}$$

$$\text{Q.24 } \frac{1}{2\sqrt{x}} \left(\frac{1}{1+x^2} \right)$$

$$\text{Q.16 } 1$$

$$\text{Q.19 } \frac{-2}{\sqrt{1-a^2x^2}}$$

$$\text{Q.22 } \frac{1}{\sqrt{a^2+x^2}}$$

$$\text{Q.3 } -\sec^3 \theta$$

$$\text{Q.6 } \frac{3x^2 \cos\{\log(x^3-1)\}}{x^3-1}$$

$$\text{Q.9 } 5^{\ln \sin x} (\cot x) (\ln 5)$$

$$\text{Q.12 } \frac{\sec^2 x}{(1-\tan x)^2} \sqrt{\frac{1-\tan x}{1+\tan x}}$$

$$\text{Q.17 } \frac{2}{1+x^2}$$

$$\text{Q.20 } \frac{1}{4} \times \frac{1}{\sqrt{a+\sqrt{a+x}}} \times \frac{1}{\sqrt{a+x}}$$

$$\text{Q.23 } \frac{x \cot\left(\sqrt{1+x^2}\right)}{\sqrt{1+x^2}}$$

Application of Derivatives

$$\text{Q.1 } (2, 1)$$

$$\text{Q.4 } \frac{dy}{dx} \text{ is not defined}$$

$$\text{Q.7 } x + 12y - 21 = 0; 12x - y + 38 = 0$$

$$\text{Q.10 } 8x - 2y + 1 = 0$$

$$\text{Q.13 } 3$$

$$\text{Q.16 } 1/818.x + 9y - 55 = 0; x + 9y - 35 = 0$$

$$\text{Q.19 } 20x + y - 140 = 0$$

$$\text{Q.22 } 2x + 2y = a^2$$

$$\text{Q.2 } 90^\circ$$

$$\text{Q.5 } \left(\frac{3}{2}, \frac{-17}{2} \right)$$

$$\text{Q.8 } \sqrt{2}bx + \sqrt{2}ay - ab$$

$$\text{Q.11 } 4x - 4y + 33 = 0$$

$$\text{Q.14 } \left(\frac{1}{2}, \frac{1}{4} \right)$$

$$\text{Q.17 } -1$$

$$\text{Q.20 } \frac{x}{\sqrt{x_1}} + \frac{x}{\sqrt{y_1}} = \sqrt{a}$$

$$\text{Q.23 } (2, -4); \left(-\frac{2}{3}, \frac{4}{27} \right)$$

$$\text{Q.3 } -\frac{1}{3}$$

$$\text{Q.6 } (2 - 36)$$

$$\text{Q.9 } M_1 M_2 = -1$$

$$\text{Q.12 } k^2x + y - 2ck = 0$$

$$\text{Q.15 } (\pm 3, 0)$$

$$\text{Q.18 } 55$$

$$\text{Q.21 } x + 3y = 8; x + 3y = -8$$

$$\text{Q.24 } \tan^{-1} \frac{3a^{1/3}b^{1/3}}{2(a^{2/3} + b^{2/3})}$$

Exercise 2

Methods of Differentiation

Single Correct Choice Type

Q.1 B	Q.2 B	Q.3 D	Q.4 D	Q.5 A	Q.6 B
Q.7 C	Q.8 A	Q.9 A	Q.10 B	Q.11 C	Q.12 D
Q.13 D	Q.14 D	Q.15 B	Q.16 B	Q.17 C	Q.18 A
Q.19 B	Q.20 D	Q.21 B			

Application of Derivatives

Single Correct Choice Type

Q.1 B	Q.2 D	Q.3 C	Q.4 D	Q.5 D	Q.6 A
Q.7 C	Q.8 A	Q.9 D	Q.10 B	Q.11 C	Q.12 D
Q.13 D	Q.14 D	Q.15 C			

Previous Years' Questions

Q.1 C	Q.2 A	Q.3 D	Q.4 D	Q.5 D	Q.6 A
Q.7 C	Q.8 D	Q.9 A	Q.10 A	Q.11 D	

JEE Advanced/Boards

Exercise 1

Methods of Differentiation

Q.1 $k = 1, -1$ or 0	Q.2 $\frac{4x^3}{9}$	Q.3 9
Q.4 $\frac{-9}{4}$	Q.56	Q.6 25
Q.7 $k = 2$	Q.9 $f'(x) = \begin{cases} \frac{x \cos x - \sin x}{x^2} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}; f''(0) = -\frac{1}{3}$	
Q.11 Zero	Q.13 3	Q.14 100
Q.15 6	Q.16 $2(1 + 2x) \cdot \cos 2(x + x^2)$	Q.17 $f(0) + kx$
Q.18 $\frac{1}{1 + (x+n)^2} - \frac{1}{1+x^2}$	Q.19 $= X[S_1 t_2 X^2 - S_2 t_1 X^2] + X_3 \begin{vmatrix} S_1 & t_1 \\ S_2 & t_2 \end{vmatrix}$	
Q.20 0	Q.21 $\frac{1-2x}{2\sqrt{1-x^2}}$	Q.22 $\frac{1}{2}$ or $-\frac{1}{2}$
Q.23 L.H.S = R.H.S	Q.24 (a) (i) $y'(0) = -1$; (ii) $y''(0) = 2$; (b) $a = 1, -2$	

Application of Derivatives

Q.1 $2\sqrt{3}x - y = 2(\sqrt{3} - 1)$ or $2\sqrt{3}x + y = 2(\sqrt{3} + 1)$	Q.2 $a = 1, b = \frac{-5}{2}$
Q.3 $(0, 1)$	Q.4 $x = y - 1 = 0$
	Q.5 T : $x - 2y = 0$; N : $2x + y = 0$

$$\text{Q.6 } -\frac{82.7^3}{3}$$

$$\text{Q.7 } x + 2y = \pi/2 \text{ \& } x + 2y = -3\pi/2$$

$$\text{Q.8 } 5$$

$$\text{Q.9 (i) (a) } 6.05, \text{ (b) } \frac{80}{27}; \text{ (ii) } 9.72 \pi \text{ cm}^2$$

$$\text{Q.10 } a = 1$$

$$\text{Q.11 } 1/16$$

$$\text{Q.12 } -\frac{1}{x+2}; x - 4y = 2$$

$$\text{Q.13 } a = -1/2; b = -3/4; c = 3$$

$$\text{Q.14 } 3, 12$$

$$\text{Q.15 } 9$$

$$\text{Q.16 } 2e^{\frac{-x}{2}}$$

$$\text{Q.17 (a) } n = -2, \text{ (c) } \pm \frac{1}{2\sqrt{2}}$$

$$\text{Q.18 } 65$$

$$\text{Q.19 } 64$$

$$\text{Q.20 (i) } 1, \text{ (ii) } 2$$

$$\text{Q.21 (4, 11) \& (-4, -31/3)}$$

$$\text{Q.22 (i) } 6 \text{ km/h; (ii) } 2 \text{ km/hr}$$

$$\text{Q.23 } 1 + 36 \pi \text{ cu. cm / sec}$$

$$\text{Q.24 } \frac{\sqrt{2}}{4\pi} \text{ cm/sec}$$

$$\text{Q.25 } 1/48 \pi \text{ cm/s}$$

$$\text{Q.26 } \frac{1}{4} \text{ cm/sec}$$

$$\text{Q.27 } \frac{66}{7}$$

$$\text{Q.28 (a) } r = (1 + t)^{1/4}, \text{ (b) } t = 80$$

$$\text{Q.29 (a) } -\frac{1}{24\pi} \text{ m/min, (b) } -\frac{5}{288\pi} \text{ m/min}$$

Exercise 2

Methods of Differentiation

Single Correct Choice Type

$$\text{Q.1 D}$$

$$\text{Q.2 B}$$

$$\text{Q.3 C}$$

$$\text{Q.4 B}$$

$$\text{Q.5 D}$$

$$\text{Q.6 C}$$

$$\text{Q.7 C}$$

$$\text{Q.8 D}$$

$$\text{Q.9 B}$$

$$\text{Q.10 D}$$

$$\text{Q.11 B}$$

$$\text{Q.12 D}$$

$$\text{Q.13 C}$$

$$\text{Q.14 C}$$

$$\text{Q.15 C}$$

$$\text{Q.16 B}$$

$$\text{Q.17 B}$$

Multiple Correct Choice Type

$$\text{Q.18 B, C}$$

$$\text{Q.19 A, B, C}$$

$$\text{Q.20 ACD}$$

$$\text{Q.21 A, B, C, D}$$

$$\text{Q.22 A, B, C}$$

Application of Derivatives

Single Correct Choice Type

$$\text{Q.1 C}$$

$$\text{Q.2 D}$$

$$\text{Q.3 C}$$

$$\text{Q.4 A}$$

$$\text{Q.5 C}$$

$$\text{Q.6 B}$$

$$\text{Q.7 D}$$

$$\text{Q.8 A}$$

Multiple Correct Choice Type

$$\text{Q.9 A, D}$$

$$\text{Q.10 A, B}$$

$$\text{Q.11 B, D}$$

$$\text{Q.12 A, B}$$

$$\text{Q.13 A, B, D}$$

$$\text{Q.14 A, B}$$

$$\text{Q.15 B, D}$$

Previous Years' Questions

$$\text{Q.1 B, C}$$

$$\text{Q.2 B, D}$$

$$\text{Q.3 } H = \emptyset, V = \{1.1\}$$

$$\text{Q.4 } \frac{ud}{\sqrt{v^2 - u^2}}$$

$$\text{Q.5 (0, 0)}$$

$$\text{Q.6 } x + 2y = \frac{\pi}{2} \text{ and } x + 2y = -\frac{3\pi}{2}$$

$$\text{Q.7 (0, 2)}$$

$$\text{Q.8 } 3/4$$

$$\text{Q.9 } \sqrt{2}x - 2y + 2 = 0$$

$$\text{Q.10 } dy/dx = 1$$

$$\text{Q.11 } 1/16$$

$$\text{Q.12 A C}$$

$$\text{Q.13 B}$$

Solutions

JEE Main/Boards

Exercise 1

Methods of Differentiation

Sol 1: $y = e^{\sqrt{x+3}}$

$$\frac{dy}{dx} = \frac{de^{\sqrt{x+3}}}{d\sqrt{x+3}} \cdot \frac{d\sqrt{x+3}}{dx} \cdot \frac{d(x+3)}{dx} \quad [\text{Chain Rule}]$$

$$= e^{\sqrt{x+3}} \cdot \frac{1}{2\sqrt{x+3}} \cdot 1 = \frac{1}{2\sqrt{x+3}} e^{\sqrt{x+3}}$$

Sol 2: $y = \sin(\log x)$

$$\frac{dy}{dx} = \frac{d\sin(\log x)}{d(\log x)} \cdot \frac{d(\log x)}{dx} \quad [\text{Chain rule}]$$

$$= \cos(\log x) \left(\frac{1}{x} \right) = \frac{1}{x} \cos(\log x)$$

Sol 3: $y = -\tan \theta$, $x = \sin \theta$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\frac{d(-\tan \theta)}{d\theta}}{\frac{d(\sin \theta)}{d\theta}} = \frac{-\sec^2 \theta}{\cos \theta} = -\sec^3 \theta$$

Sol 4: $y = \cos^{-1} \sqrt{x}$

$$\frac{dy}{dx} = \frac{d\cos^{-1} \sqrt{x}}{d\sqrt{x}} \cdot \frac{d\sqrt{x}}{dx} = \frac{-1}{\sqrt{1-x}} \cdot \frac{1}{2\sqrt{x}} = \frac{-1}{2\sqrt{x-x^2}}$$

Sol 5: $y = e^{m \tan^{-1} x}$

$$\frac{dy}{dx} = \frac{de^{m \tan^{-1} x}}{d(m \tan^{-1} x)} \cdot \frac{d(m \tan^{-1} x)}{dx}$$

$$= e^{m \tan^{-1} x} \cdot \left(m \cdot \frac{1}{1+x^2} \right) = \frac{me^{m \tan^{-1} x}}{1+x^2}$$

Sol 6: $y = \sin\{\log(x^3 - 1)\}$

$$\frac{dy}{dx} = \frac{d\sin\{\log(x^3 - 1)\}}{d\{\log(x^3 - 1)\}} \times \frac{d\{\log(x^3 - 1)\}}{d\{x^3 - 1\}} \times \frac{d(x^3 - 1)}{dx}$$

$$= \cos\{\log(x^3 - 1)\} \cdot \frac{1}{x^3 - 1} \cdot 3x^2 = \frac{3x^2 \cos\{\log(x^3 - 1)\}}{x^3 - 1}$$

Sol 7: $\frac{d\cos x}{de^x} = \frac{d\cos x}{dx} \cdot \frac{1}{\frac{de^x}{dx}} = \frac{-\sin x}{e^x} = -(\sin x)e^{-x}$

Sol 8: $y = \log_2(\sin x)$

$$\frac{dy}{dx} = \frac{1}{\log 2} \cdot \frac{d\log(\sin x)}{d(\sin x)} \cdot \frac{d(\sin x)}{dx} = \frac{\cos x}{\sin x} \log_2 e = (\cot x) \log_2 e$$

Sol 9: $y = 5^{\log(\sin x)}$

$$\log y = (\log 5) \log(\sin x)$$

$$\frac{1}{y} \frac{dy}{dx} = (\log 5) \frac{d\log(\sin x)}{d(\sin x)} \cdot \frac{d(\sin x)}{dx} = (\log 5) \frac{\cos x}{\sin x}$$

$$\therefore \frac{dy}{dx} = 5^{\log(\sin x)} ((\cot x) \log 5)$$

Sol 10: $\sqrt{x} + \sqrt{y} = 5$

Differentiate w.r.t x

$$\frac{1}{2\sqrt{x}} + \frac{1}{2\sqrt{y}} \frac{dy}{dx} = 0 \quad \therefore \frac{dy}{dx} = -\frac{\sqrt{y}}{\sqrt{x}}$$

For (4, 9)

$$\frac{dy}{dx} = -\frac{\sqrt{9}}{\sqrt{4}} = -\frac{3}{2}$$

Sol 11: $y = \cot^{-1} \left(\frac{1+x}{1-x} \right) = \tan^{-1} \left(\frac{1-x}{1+x} \right)$

Take $x = \tan \theta$

$$\therefore y = \tan^{-1} \left(\frac{\tan \frac{\pi}{4} - \tan \theta}{1 + \tan \frac{\pi}{4} \tan \theta} \right)$$

$$= \tan^{-1} \left(\tan \left(\frac{\pi}{4} - \theta \right) \right) = \frac{\pi}{4} - \theta = \frac{\pi}{4} - \tan^{-1} x$$

$$\therefore \frac{dy}{dx} = \frac{d\tan^{-1} x}{dx} = -\frac{1}{1+x^2}$$

$$\text{Sol 12: } y = \sqrt{\frac{1+\tan x}{1-\tan x}} = \sqrt{\tan\left(x + \frac{\pi}{4}\right)}$$

$$\therefore \frac{dy}{dx} = \frac{d\sqrt{\tan\left(x + \frac{\pi}{4}\right)}}{d\tan\left(x + \frac{\pi}{4}\right)} \cdot \frac{d\tan\left(x + \frac{\pi}{4}\right)}{dx}$$

$$= \frac{1}{2\sqrt{\tan\left(x + \frac{\pi}{4}\right)}} \cdot \sec^2\left(x + \frac{\pi}{4}\right)$$

$$= \frac{1}{2} \left(\sqrt{\frac{1-\tan x}{1+\tan x}} \right) \left(1 + \tan^2\left(x + \frac{\pi}{4}\right) \right)$$

$$= \frac{1}{2} \left(\sqrt{\frac{1-\tan x}{1+\tan x}} \right) \left(\frac{1 + \tan^2 x}{(1 - \tan x)^2} \right)^2$$

$$= \frac{\sec^2 x}{(1 - \tan x)^2} \sqrt{\frac{1 - \tan x}{1 + \tan x}}$$

$$\text{Sol 13: } \frac{-\cos\sqrt{\cos}\sqrt{x} \sin\sqrt{x}}{4\sqrt{x}\sqrt{\cos}\sqrt{x}}$$

$$\text{Sol 14: } \frac{2}{x(\log x^2)\log(\log x^2)}$$

$$\text{Sol 15: } y = (\sin x)^{\cos^{-1} x}$$

Taking log both sides

$$\log y = \cos^{-1} x \log(\sin x)$$

Differentiate both side w.r.t. x

$$\frac{1}{y} \frac{dy}{dx} = \frac{d\cos^{-1} x \log(\sin x)}{dx}$$

$$= \cos^{-1} x \frac{d\log(\sin x)}{dx} + \log(\sin x) \frac{d\cos^{-1} x}{dx}$$

$$= \cot x \cos^{-1} x - \frac{\log(\sin x)}{\sqrt{1-x^2}}$$

$$\therefore \frac{dy}{dx} = (\sin x)^{\cos^{-1} x} \left[\cot x \cos^{-1} x - \frac{\log(\sin x)}{\sqrt{1-x^2}} \right]$$

$$\text{Sol 16: } y = \cos^{-1} \left[(2 \cos x + 3 \sin x) \sqrt{13} \right]$$

$$= \cos^{-1} \left[\frac{2}{\sqrt{13}} \cos x + \frac{3}{\sqrt{13}} \sin x \right]$$

$$\text{where } \frac{2}{\sqrt{13}} = \cos \theta \text{ and } \frac{3}{\sqrt{13}} = \sin \theta$$

$$= \cos^{-1} [\cos \theta \cos x + \sin \theta \sin x] = \cos^{-1} [\cos(x - \theta)]$$

$$\therefore \cos^{-1}(\cos x) = x \quad \therefore y = x - \theta$$

$$\Rightarrow \frac{dy}{dx} = 1$$

$$\text{Sol 17: } y = \sec^{-1} \left(\frac{1+x^2}{1-x^2} \right) = \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)$$

Take $x = \tan \theta$

$$\therefore y = \cos^{-1} \left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right) = \cos^{-1}(\cos 2\theta)$$

$$\Rightarrow y = 2\theta \quad \therefore y = 2\tan^{-1} x$$

$$\Rightarrow \frac{dy}{dx} = \frac{2}{1+x^2}$$

$$\text{Sol 18: } y = \sin^{-1} \left[\frac{\sqrt{1+x} + \sqrt{1-x}}{2} \right]$$

Put $x = \cos 2\theta$

$$\therefore y = \sin^{-1} \left[\frac{\sqrt{1+\cos 2\theta} + \sqrt{1-\cos 2\theta}}{2} \right]$$

$$= \sin^{-1} \left[\frac{\sqrt{2\cos^2 \theta} + \sqrt{2\sin^2 \theta}}{2} \right]$$

$$= \sin^{-1} \left[\frac{1}{\sqrt{2}} \cos \theta + \frac{1}{\sqrt{2}} \sin \theta \right]$$

$$= \sin^{-1} \left[\sin \left(\theta + \frac{\pi}{4} \right) \right] = \theta + \frac{\pi}{4} \quad \therefore y = \frac{1}{2} \cos^{-1} x + \frac{\pi}{4}$$

$$\frac{dy}{dx} = -\frac{1}{2\sqrt{1-x^2}}$$

$$\text{Sol 19: } y = \sin^{-1} \left[2ax\sqrt{1-a^2x^2} \right]$$

$$\text{Put } x = \frac{1}{a} \cos \theta \quad \therefore y = \sin^{-1} \left[2\cos \theta \sqrt{1-a^2 \left(\frac{\cos^2 \theta}{a^2} \right)} \right]$$

$$= \sin^{-1} \left[2\cos \theta \sqrt{1-\cos^2 \theta} \right]$$

$$= \sin^{-1} [2\cos \theta \sin \theta] = \sin^{-1} [\sin 2\theta] = 2\theta = 2\cos^{-1} ax$$

$$\therefore \frac{dy}{dx} = -2 \times \frac{1}{\sqrt{1-a^2x^2}} = \frac{-2}{\sqrt{1-a^2x^2}}$$

Sol 20: $y = \sqrt{a + \sqrt{a + x}}$

$$\frac{dy}{dx} = \frac{d\sqrt{a + \sqrt{a + x}}}{d(a + \sqrt{a + x})} \times \frac{d(a + \sqrt{a + x})}{dx}$$

$$= \frac{1}{2} \times \frac{1}{\sqrt{a + \sqrt{a + x}}} \times \left(\frac{1}{2\sqrt{a + x}} \right) \times 1$$

$$= \frac{1}{4} \times \frac{1}{\sqrt{a + \sqrt{a + x}}} \times \frac{1}{\sqrt{a + x}}$$

Sol 21: $y = \tan^{-1} \left[\frac{ax - b}{a + bx} \right] = \tan^{-1} \left[\frac{x - \frac{b}{a}}{1 + \frac{b}{a}x} \right]$

Let $\tan \alpha = \frac{b}{a}$, $x = \tan t$

$$\therefore y = \tan^{-1} \left[\frac{\tan t - \tan \alpha}{1 + \tan \alpha \tan t} \right] \text{ or } y = \tan^{-1} \tan(t - \alpha)$$

$= t - \alpha$, α is constant

$$\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} \text{ or } y = t - \alpha = \tan^{-1} x - \alpha$$

$$\therefore \frac{dy}{dx} = \frac{1}{1 + x^2}$$

Sol 22: $y = \log(x + \sqrt{x^2 + a^2})$

$$\begin{aligned} \frac{dy}{dx} &= \frac{d \log(x + \sqrt{x^2 + a^2})}{d(x + \sqrt{x^2 + a^2})} \cdot \frac{d(x + \sqrt{x^2 + a^2})}{dx} \\ &= \frac{1}{(x + \sqrt{x^2 + a^2})} \cdot \left[1 + \frac{d(\sqrt{x^2 + a^2})}{d(x^2 + a^2)} \cdot \frac{d(x^2 + a^2)}{dx} \right] \end{aligned}$$

$$= \frac{1}{(x + \sqrt{x^2 + a^2})} \left[1 + \frac{1}{2\sqrt{x^2 + a^2}} \times 2x \right]$$

$$= \frac{1}{(x + \sqrt{x^2 + a^2})} \cdot \left[\frac{\sqrt{x^2 + a^2} + x}{\sqrt{x^2 + a^2}} \right] = \frac{1}{\sqrt{x^2 + a^2}}$$

Sol 23: $y = \log(\sin \sqrt{1 + x^2})$

$$\frac{dy}{dx} = \frac{d \log(\sin \sqrt{1 + x^2})}{d(\sin \sqrt{1 + x^2})} \times \frac{d(\sin \sqrt{1 + x^2})}{d(\sqrt{1 + x^2})} \times \frac{d\sqrt{1 + x^2}}{dx}$$

$$\begin{aligned} &= \frac{1}{\sin(\sqrt{1 + x^2})} (\cos \sqrt{1 + x^2}) \times \frac{1}{2\sqrt{1 + x^2}} \times \frac{d(1 + x^2)}{dx} \\ &= \frac{x \cot(\sqrt{1 + x^2})}{\sqrt{1 + x^2}} \end{aligned}$$

Sol 24: $y = \tan^{-1} \left(\frac{\sqrt{x} + \sqrt{a}}{1 - \sqrt{a}\sqrt{x}} \right)$

Put $\sqrt{x} = \tan t$

$\sqrt{a} = \tan \alpha$

$$\therefore y = \tan^{-1} \left(\frac{\tan t + \tan \alpha}{1 - \tan t \tan \alpha} \right) = \tan^{-1} \tan(t + \alpha) = t + \alpha$$

$$y = \tan^{-1} \sqrt{x} + \alpha$$

$$\begin{aligned} \therefore \frac{dy}{dx} &= \frac{d(\tan^{-1} \sqrt{x})}{d\sqrt{x}} \cdot \frac{d\sqrt{x}}{dx} + \frac{d\alpha}{dx} \\ &= \frac{1}{1 + (\sqrt{x})^2} \cdot \frac{1}{2\sqrt{x}} = \frac{1}{2\sqrt{x}(1 + x^2)} \end{aligned}$$

Application of Derivatives

Sol 1: $y = x^2 - 4x + 5$

Tangent is parallel to x axis $\Rightarrow \frac{dy}{dx} = 0 = 2x - 4$

$$\Rightarrow x = 2; y = 4 - 8 + 5 = 1$$

Point A (2, 1)

Sol 2: If two curves cut orthogonally, then the tangents at point of intersection are perpendicular.

Sol 3: $f(x) = 3x^2 - 5$

Tangent at $\left[x = \frac{1}{2}, y = \left(\frac{3}{4} - 5 \right) \right]$

$$\left(\frac{dy}{dx} \right)_{x=1/2} = 6x = 3 = (m)_{\text{tangent}}$$

$$\left(-\frac{dx}{dy} \right) = -\frac{1}{3} = (m)_{\text{normal}}$$

Sol 4: $y = f(x)$, $\frac{dy}{dx} = \infty$ i.e. not defined

Sol 5: $y = 2x^2 - 6x - 4$

$$\frac{dy}{dx} = 4x - 6 = 0 \Rightarrow x = \frac{3}{2}$$

$$y = 2 \times \frac{9}{4} - 6 \times \frac{3}{2} - 4 \Rightarrow \frac{9}{2} - 9 - 4 = -\frac{17}{2}$$

$$\left(\frac{3}{2}, -\frac{17}{2}\right)$$

Sol 6: $y = x^2 - 4x - 32$

$$\frac{dy}{dx} = 2x - 4 = 0$$

$$x = 2; y = 4 - 8 - 32 = -36$$

$$\text{Point } (2, -36)$$

Sol 7: $y = (5 - x)^{1/3}$

$$\text{At } P(-3, 2)$$

$$y = (8)^{1/3} = 2$$

$$\left(\frac{dy}{dx}\right)_{x=-3} = \frac{1}{3}(5-x)^{-2/3} = \frac{-1}{3(8)^{2/3}} = \frac{-1}{12}$$

$$\frac{y-2}{x+3} = \frac{-1}{12} \Rightarrow 12y - 24 = -(x+3) \Rightarrow x + 12y = 21$$

$$\text{Equation of normal is } \frac{y-2}{x+3} = +12$$

$$\Rightarrow y - 12x = 38$$

Sol 8: $y = (4x - 3)^{1/2} \Rightarrow \frac{dy}{dx} = \frac{1}{2\sqrt{4x-3}} = \frac{1}{\sqrt{4x-3}} \neq 0$, it can never be parallel to x axis.

Sol 9: $x^2 = y$ & $x^3 + 6y = 7$

$$x^3 + 6x^2 = 7$$

$$(x^2 + 7x + 7)(x - 1) = 0$$

$$\Rightarrow x = 1$$

$$\text{Point of intersection } (1, 1)$$

$$P_1(x^2 = y)P_2(x^3 + 6y = 7)$$

$$\frac{dy}{dx} = 2x \quad 3x^2 + \frac{6dy}{dx} = 0$$

$$\left(\frac{dy}{dx}\right)_{x=1} = 2 \quad \left(\frac{dy}{dx}\right)_{x=1} = \frac{-1}{2}$$

$$M_1 M_2 = -1 \text{ i.e., tangents are orthogonal at } (1, 1)$$

Sol 10: $y^2 = 8x$

$$\Rightarrow 2y \frac{dy}{dx} = 8 \Rightarrow \frac{dy}{dx} = 4 = \frac{8}{2y}$$

$$\text{This give } y = 1; x = \frac{1}{8}$$

$$\Rightarrow \frac{y-1}{x-\frac{1}{8}} = 4 \Rightarrow \frac{y-1}{8x-1} = \frac{1}{2} \Rightarrow 2y-2 = 8x-1$$

$$\Rightarrow 8x - 2y + 1 = 0$$

Sol 11: $y = -5x^2 + 6x + 7$

$$y' = -10x + 6 \Rightarrow (y')_{x=\frac{1}{2}} = -5 + 6 = 1$$

$$y - \frac{35}{4} = x - \frac{1}{2}$$

$$4y - 35 = 4x - 2 \Rightarrow 4x - 4y + 33 = 0$$

Sol 12: $y = \frac{c^2}{x}$

$$\frac{dy}{dx} = \frac{-c^2}{x^2} \Rightarrow \left(\frac{dy}{dx}\right)_{\frac{c}{k}} = -k^2$$

$$\text{Equation: } -y - ck$$

$$= -k^2 \left(x - \frac{c}{k}\right) \Rightarrow y + k^2x = 2x$$

Sol 13: $y = x^3 + 6 \Rightarrow \frac{dy}{dx} = 3x^2$

$$(y')_{x=-1, y=5} = 3; (y')_{x=-1, y=7} = 3$$

So the tangents are parallel

Sol 14: $y = x^2 \Rightarrow \left(\frac{dy}{dx}\right) = 2x = 1$

$$x = \frac{1}{2}; y = \frac{1}{4}; P\left(\frac{1}{2}, \frac{1}{4}\right)$$

Sol 15: $\frac{x^2}{9} + \frac{y^2}{4} = 1$

$$\frac{2x}{9} + \frac{2y}{4} y' = 0 \Rightarrow y' = -\frac{4x}{9y}$$

This will be parallel to y axis if $y = 0$

$$x = \pm 3$$

$$P(+3, 0), (-3, 0)$$

Sol 16: $x = \frac{1}{t}; y = 2t \Rightarrow xy = 2$

$$y = \frac{2}{x} \Rightarrow y' = \frac{-2}{x^2} = -8$$

At $t = 2$ i.e. $x = \frac{1}{2}$

$$y' = -8$$

Slope of normal = $\frac{1}{8}$

Sol 17: $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$$\frac{2x}{a^2} + \frac{2yy'}{b^2} = 0 \Rightarrow y' = -\frac{b^2x}{a^2y}$$

At $(x_0, y_0); y' = -\frac{b^2x_0}{a^2y_0}$

$$\frac{y - y_0}{x - x_0} = -\frac{b^2x_0}{a^2y_0}$$

$$a^2yy_0 - a^2y_0^2 = b^2x_0^2 - b^2xy_0$$

$$xx_0b^2 + yy_0a^2 = a^2y_0^2 + b^2x_0^2 = a^2b^2$$

$$\Rightarrow \frac{xx_0}{a^2} + \frac{yy_0}{b^2} = 1$$

Sol 18: $y = 4x^3 - 3x + 5$

$$9y + x + 3 = 0 \text{ [Given]}$$

$$M_{\text{normal}} = \frac{-1}{9}$$

$$y' = 12x^2 - 3 = 9 \Rightarrow x = \pm 1; y = 6, 4$$

$$\frac{y-4}{x+1} = \frac{-1}{9}, \frac{y-6}{x-1} = \frac{-1}{9}$$

$$x + 9y = 35, x + 9y = 55$$

Sol 19: $y(x-2)(x-3) = x-7$

$$y = \frac{x-7}{(x-2)(x-3)} = 0 \Rightarrow x = 7$$

$$\frac{dy}{dx} = \frac{(x-2)(x-3) - (x-7)(2x-5)}{(x-2)^2(x-3)^2}$$

$$\left(\frac{-dx}{dy} \right)_{x=7} = \frac{(x-2)^2(x-3)^2}{(x-7)(2x-5) - (x-2)(x-3)} = \frac{5^2(4)^2}{0 - (5)(4)} = -20$$

$$\frac{y}{x-7} = -20 \Rightarrow 20x + y = 140$$

Sol 20: $\sqrt{x} + \sqrt{y} = \sqrt{a}$

$$\frac{1}{2\sqrt{x}} + \frac{1}{2\sqrt{y}} y' = 0 \Rightarrow y' = -\sqrt{\frac{y}{x}} = -\sqrt{\frac{y_1}{x_1}}$$

$$\Rightarrow \frac{y-y_1}{x-x_1} = -\sqrt{\frac{y_1}{x_1}} \Rightarrow y\sqrt{x_1} - y_1\sqrt{x_1} = -x\sqrt{y_1} + x_1\sqrt{y_1}$$

$$\Rightarrow x\sqrt{y_1} + y\sqrt{x_1} = y_1\sqrt{x_1} + x_1\sqrt{y_1}$$

$$\frac{x}{\sqrt{x_1}} + \frac{y}{\sqrt{y_1}} = \sqrt{a}$$

Sol 21: $3x^2 + y^2 = 8 \Rightarrow 6x + 2yy' = 0 \Rightarrow y' = \frac{-3x}{y}$

$$(m)_{\text{normal}} = \frac{y}{3x} = \frac{-1}{3}$$

$$y = -x$$

$$x = \pm \sqrt{2}, y = \mp \sqrt{2}$$

$$\frac{y-\sqrt{2}}{x+\sqrt{2}} = \frac{-1}{3} \Rightarrow 3y+x = 2\sqrt{2}$$

$$\frac{y+\sqrt{2}}{x-\sqrt{2}} = \frac{-1}{3} \Rightarrow 3y+x = -2\sqrt{2}$$

Sol 22: $\sqrt{x} + \sqrt{y} = \sqrt{a}$

$$\frac{1}{2\sqrt{x}} + \frac{y'}{2\sqrt{y}} = 0$$

$$y' = -\sqrt{\frac{y}{x}} = -\sqrt{1} = -1$$

$$\frac{y - \frac{a^2}{4}}{x - \frac{a^2}{4}} = 1 \Rightarrow y + x = \frac{a^2}{2}$$

Sol 23: $y = x^3 - 2x^2 - 2x$

$$y' = 3x^2 - 4x - 2 = 2$$

$$3x^2 - 6x + 2x - 4 = 0$$

$$(3x+2)(x-2) = 0$$

$$x = 2, \frac{-2}{3}; y = 8 - 8 - 4 = -4,$$

$$y = \frac{-8}{27} - \frac{2 \times 4}{9} + \frac{4}{3} = -\frac{8}{27} - \frac{8}{9} + \frac{4}{3}$$

$$\Rightarrow \frac{36 - 24 - 8}{27} = \frac{4}{27}$$

Therefore, the points are

$$(2, -4) \left(\frac{-2}{3}, \frac{4}{27} \right)$$

Sol 24: $x^2 = 4$ by

$$y^2 = 4axx^2 = 16b(a^2b)^{1/3} = 16b^{4/3}a^{2/3}$$

$$y^4 = 16a^2 4byx = 4(b^2a)^{1/3}$$

$$y^3 = 64a^2by' = \frac{\frac{x}{y}}{\frac{2b}{2a}}$$

$$y = 4(a^2b)^{1/3}y' = 2 \frac{\left(\frac{a}{b}\right)^{1/3}}{2a} = \frac{1}{2} \left(\frac{a}{b}\right)^{1/3}$$

$$\tan \theta = \frac{2\left(\frac{a}{b}\right)^{1/3} - \frac{1}{2}\left(\frac{a}{b}\right)^{1/3}}{1 + \left(\frac{a}{b}\right)^{2/3}} = \frac{\frac{3}{2}\left(\frac{a}{b}\right)^{1/3}}{1 - \left(\frac{a}{b}\right)^{2/3}}$$

$$= \frac{3a^{1/3}b^{1/3}}{2(b^{2/3} + a^{2/3})}$$

Exercise 2

Methods of Differentiation

Single Correct Choice Type

Sol 1: (B) $y = f\left(\frac{3x+4}{5x+6}\right)$ $f'(x) = \tan x^2$

$$\begin{aligned} \frac{dy}{dx} &= \frac{df\left(\frac{3x+4}{5x+6}\right)}{d\left(\frac{3x+4}{5x+6}\right)} \cdot \frac{d\left(\frac{3x+4}{5x+6}\right)}{dx} \\ &= \left[\tan\left(\frac{3x+4}{5x+6}\right) \right]^2 \frac{(5x+6)3 - (3x+4)5}{(5x+6)^2} \\ &= -2 \tan\left(\frac{3x+4}{5x+6}\right)^2 \cdot \frac{1}{(5x+6)^2} \end{aligned}$$

Sol 2: (B) $f'(x) =$ $g(x) = f^{-1}(x)$

$$\frac{x^{10}}{(1+x^2)}$$

$$\therefore f(g(x)) = x$$

$$\therefore f'(g(x))g'(x) = 1$$

$$\therefore g'(x) = \frac{1}{f'(g(x))}$$

$$\therefore g'(2) = \frac{1}{f'(g(2))} = \frac{1}{f'(a)} = \frac{1}{\frac{1}{a^{10}}} = \frac{1+a^2}{a^{10}}$$

$$\text{Sol 3: (D)} \quad y = \frac{1}{1+x^{n-m}+x^{p-m}} + \frac{1}{1+x^{m-n}+x^{p-n}} + \frac{1}{1+x^{m-p}+x^{n-p}}$$

$$= \frac{x^m}{x^p+x^m+x^n} + \frac{x^n}{x^n+x^m+x^p} + \frac{x^p}{x^p+x^m+x^n} = \frac{x^m+x^n+x^p}{x^m+x^n+x^p} = 1$$

$$\therefore \frac{dy}{dx} = 0$$

$$\begin{aligned} \text{Sol 4: (D)} \quad \lim_{x \rightarrow 2} \frac{f(4) - f(x^2)}{2-x} &= \lim_{x \rightarrow 2} \frac{\frac{d(f(4) - f(x^2))}{dx}}{\frac{d(2-x)}{dx}} \\ &= \lim_{x \rightarrow 2} \frac{f'(x^2)[2x]}{-1} = f'(2^2) \cdot 2 \cdot 2 = 4f'(4) = 20 \end{aligned}$$

Sol 5: (A) $\ell = \lim_{x \rightarrow 0} x^m (\ln x)^n \quad m, n \in \mathbb{N}$

$$\ell = \lim_{x \rightarrow 0} \frac{(\ln x)^n}{\left(\frac{1}{x}\right)^m} = \lim_{x \rightarrow 0} \frac{n(\ln x)^{n-1} \frac{1}{x}}{-m \left(\frac{1}{x}\right)^{m+1}}$$

(using L-Hospital rule)

$$= \lim_{x \rightarrow 0} \left(-\frac{n}{m} \frac{(\ln x)^{n-1}}{\left(\frac{1}{x}\right)^m} \right) = \lim_{x \rightarrow 0} \frac{-n(n-1)(n-2) \dots 1 \ln x}{m^{n-1} \left(\frac{1}{x}\right)^m}$$

$$= \lim_{x \rightarrow 0} \frac{-(n)! \frac{1}{x}}{m^n \left(\frac{1}{x}\right)^{m+1}} = \lim_{x \rightarrow 0} \frac{-(n)!}{m^n} (x)^m = 0$$

\therefore Independent of n and m

$$\text{Sol 6: (B)} \quad f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2 \sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$$

$$f'(x) = \begin{vmatrix} -\sin x & 1 & 0 \\ 2\sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix} + \begin{vmatrix} \cos x & x & 1 \\ 2\cos x & 2x & 2 \\ \tan x & x & 1 \end{vmatrix}$$

$$+ \begin{vmatrix} \cos x & x & 1 \\ 2\sin x & x^2 & 2x \\ \sec^2 x & 1 & 0 \end{vmatrix}$$

$$f'(x) = [x^2 \sin x (2x \tan x - 2 \sin x)] + x[2 \tan x - 2 \cos x] + [2x \cos x - 2x \tan x] + (-2x \cos x) + 2x^2 \sec^2 x + 2 \sin x - x^2 \sec^2 x$$

$$= x^2 \sin x + x^2 \sec^2 x + 2x \tan x - 2x \cos x$$

$$\lim_{x \rightarrow 0} \frac{f'(x)}{x} = \lim_{x \rightarrow 0} \frac{x^2 (\sin x + \sec^2 x) (2x (\tan x - \cos x))}{x}$$

$$= \lim_{x \rightarrow 0} x (\sin x + \sec^2 x) + 2 (\tan x - \cos x) = -2$$

$$\text{Sol 7: (C)} \quad f'(x) = \begin{vmatrix} -\sin x & +\cos x & -\sin x \\ \cos 2x & \sin 2x & 2\cos 2x \\ \cos 3x & \sin 3x & 3\cos 3x \end{vmatrix}$$

$$+ \begin{vmatrix} \cos x & \sin x & \cos x \\ -2\sin 2x & 2\cos 2x & -4\sin 2x \\ \cos 3x & \sin 3x & 3\cos 3x \end{vmatrix}$$

$$+ \begin{vmatrix} \cos x & \sin x & \cos x \\ \cos 2x & \sin 2x & 2\cos 2x \\ -3\sin 3x & +3\cos 3x & -9\sin 3x \end{vmatrix}$$

$$f'\left(\frac{\pi}{2}\right) = \begin{vmatrix} -1 & 0 & -1 \\ -1 & 0 & -2 \\ 0 & -1 & 0 \end{vmatrix} + \begin{vmatrix} 0 & 1 & 0 \\ 0 & -2 & 0 \\ 0 & -1 & 0 \end{vmatrix}$$

$$+ \begin{vmatrix} 0 & 1 & 0 \\ -1 & 0 & -2 \\ 3 & 0 & 9 \end{vmatrix} = (2 - 1) + (0) + (3) = 4$$

$$\text{Sol 8: (A)} \quad y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$$

$$\text{Put } x = \tan \theta$$

$$\therefore y = \sin^{-1} \frac{2 \tan \theta}{1 + \tan^2 \theta}$$

$$y = \sin^{-1} \sin 2\theta$$

$$-\frac{\pi}{2} \leq 2\theta \leq \frac{\pi}{2} \Rightarrow -\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$$

$$\therefore \text{ as } x = \frac{\pi}{2}$$

$$\therefore \theta = \tan^{-1} \frac{\pi}{2}$$

$$\therefore 2\theta = 2 \tan^{-1} \frac{\pi}{2} > \frac{\pi}{2}$$

$$\therefore y = \sin^{-1} \sin(\pi - 2\theta)$$

$$y = \pi - 2\theta$$

$$y = \pi - 2 \tan^{-1} x$$

$$\therefore \frac{dy}{dx} = \frac{-2}{1+x^2}$$

$$\therefore \frac{dy}{dx} \Big|_{x=\frac{\pi}{2}} = \frac{-8}{\pi^2 + 4^2}$$

$$\text{Sol 9: (A)} \quad \lim_{x \rightarrow 4} \frac{\sqrt{f(x)} - \sqrt{g(x)}}{\sqrt{x} - 2}$$

$$\frac{d[\sqrt{f(x)} - (\sqrt{g(x)})]}{dx}$$

$$= \lim_{x \rightarrow 4} \frac{\frac{1}{2\sqrt{f(x)}} f'(x) - \frac{1}{2\sqrt{g(x)}} g'(x)}{\frac{1}{2\sqrt{x}}}$$

$$= \frac{\frac{1}{2\sqrt{f(4)}} f'(4) - \frac{1}{2\sqrt{g(4)}} g'(4)}{\frac{1}{2\sqrt{4}}} = \frac{\frac{9}{\sqrt{2}} - \frac{6}{\sqrt{2}}}{\frac{1}{2}}$$

$$= 3\sqrt{2}$$

$$\text{Sol 10: (B)} \quad y = x + e^x \Rightarrow \frac{dy}{dx} = 1 + e^x$$

$$\Rightarrow \frac{dx}{dy} = \frac{1}{1+e^x}$$

$$\frac{d^2x}{dy^2} = \frac{d(1+e^x)^{-1}}{dx} \times \frac{1}{(1+e^x)}$$

$$= \frac{1}{(1+e^x)} \frac{d(1+e^x)^{-1}}{d(1+e^x)} \frac{d(1+e^x)}{dx}$$

$$= \frac{1}{(1+e^x)} \left(-\frac{1}{(1+e^x)^2} \right) e^x = \frac{-e^x}{(1+e^x)^3}$$

$$\text{Sol 11: (C)} \quad h'(x) = [f(x)]^2 + [g(x)]^2$$

$$h''(x) = 2f(x)f'(x) + 2g(x)g'(x)$$

$$\text{Also } f'(x) = g(x)$$

$$\therefore f''(x) = g'(x) = -f(x)$$

$$\therefore h''(x) = 2f(x)g(x) + 2g(x)(-f(x)) = 0$$

$$\therefore h(x) = ax + b$$

$$h(0) = b = 2; h(1) = a + b = 4$$

$$\therefore a = 2, b = 2$$

$\therefore h(x)$ is a straight line with slope 2 and y intercept 2

$$\text{Sol 12: (D)} f(x) = x + 3\ln(x-2)$$

$$F'(x) = 1 + \frac{3}{(x-2)}$$

$$g(x) = x + 5\ln(x-1)$$

$$g(x) = 1 + \frac{5}{(x-1)}$$

$$\therefore f'(x) < g'(x)$$

$$\Rightarrow \frac{3}{(x-2)} < \frac{5}{(x-1)} \Rightarrow \frac{3(x-1) - 5(x-2)}{(x-2)(x-1)} < 0$$

$$\therefore \frac{7-2x}{(x-2)(x-1)} < 0 \Rightarrow x \in (1, 2) \cup \left(\frac{7}{2}, \infty\right)$$

$$\text{Also } x-2 > 0 \text{ and } x-1 > 0$$

$$\therefore x > 2 \Rightarrow x \in \left(\frac{7}{2}, \infty\right)$$

$$\text{Sol 13: (D)} g(x) = x^2, f(x) = \sin x, h(x) = \log_e x$$

$$\therefore g(f(x)) = (\sin x)^2$$

$$h(g(f(x))) = \log_e (\sin x)^2 = F(x)$$

$$\therefore F(x) = 2\log \sin x$$

$$\therefore \frac{dF}{dx} = \frac{2}{\sin x} \cdot \cos x = 2\cot x$$

$$\therefore \frac{d^2t}{dx^2} = -2\operatorname{cosec}^2 x$$

$$\text{Sol 14: (D)} f(x) = x^n$$

$$f'(p+q) = n(p+q)^{n-1}$$

$$f'(p) = n(p)^{n-1}$$

$$f'(q) = n(q)^{n-1}$$

$$\text{for } f'(p+q) = f'(p) + f'(q)$$

$$(p+q)^{n-1} = [(p)^{n-1} + (q)^{n-1}] \quad n \neq 0$$

$$\Rightarrow \left(1 + \frac{q}{p}\right)^{n-1} - 1 = \left(\frac{q}{p}\right)^{n-1}$$

This condition satisfies of $n-1=1$

$$\Rightarrow n = 2$$

Also if $n = 0$

$$\therefore f(x) = 1$$

$$\therefore f'(p+q) = 0 = f'(p) + f'(q)$$

$$\therefore n = 0, 2 \text{ (two values)}$$

$$\text{Sol 15: (B)} f(x) = \prod_{n=1}^{100} (x-n)^{n(101-n)}$$

$$\ln f(x) = \sum_{n=1}^{100} n(101-n) \ln(x-n)$$

$$\Rightarrow \frac{1}{f(x)} f'(x) = \sum_{n=1}^{100} \frac{n(101-n)}{(x-n)}$$

$$\therefore \frac{f'(101)}{f(101)} = \sum_{n=1}^{100} \frac{n(101-n)}{(101-n)} = \sum_{n=1}^{100} n = \frac{100 \times 101}{2} = 5050$$

$$\therefore \frac{f(101)}{f'(101)} = \frac{1}{5050}$$

$$\text{Sol 16: (B)} f(x) \text{ is continuous and differentiable at}$$

$$x = \frac{1}{3}$$

$$f(x) = \frac{3x^2 + 2x - 1}{6x^2 - 5x + 1} = \frac{x+1}{2x-1}$$

$$\therefore f'(x) = \frac{(2x-1) - 2(x+1)}{(2x-1)^2} = \frac{-3}{(2x-1)^2}$$

$$f'\left(\frac{1}{3}\right) = \frac{-3}{\left(2\left(\frac{1}{3}\right) - 1\right)^2} = 27$$

$$\text{Sol 17: (C)} f(x) = ax^2 + bx + c$$

For $x \in \mathbb{R}$ $f(x)$ is always positive

$$\therefore a > 0 \text{ and } b^2 - 4ac < 0$$

$$\therefore g(x) = f(x) + f'(x) + f''(x)$$

$$= ax^2 + bx + c + 2ax + b + 2a$$

$$= ax^2 + (2a+b)x + (2a+b+c)$$

$$D = (2a+b)^2 - 4(2a+b+c)a$$

$$= 4a^2 + b^2 + 4ab - 8a^2 - 4ab - 4ac$$

$$= b^2 - 4a^2 - 4ac = (b^2 - 4ac) - 4a^2$$

$$Qb^2 - 4ac < 0$$

$$\therefore D < 0 \therefore g(x) = 0$$

$$\text{Sol 18: (A)} y = \frac{x^4 + 4}{x^2 - 2x + 2}$$

$$\frac{dy}{dx} = \frac{(x^2 - 2x + 2)(4x^3) - (x^4 + 4)(2x - 2)}{(x^2 - 2x + 2)^2}$$

$$\frac{dy}{dx} = \frac{4x^5 - 8x^4 + 8x^3 - 2x^5 + 2x^4 - 8x + 8}{(x^2 - 2x + 2)^2}$$

$$\frac{dy}{dx} = \frac{2x^5 + 8x^3 - 8x + 8 - 6x^4}{(x^2 - 2x + 2)^2}$$

$$\left. \frac{dy}{dx} \right|_{1/2} = \frac{2 \times \left(\frac{1}{2}\right)^5 + 8 \left(\frac{1}{2}\right)^3 - 8 \times \frac{1}{2} + 8 - 6 \times \frac{1}{16}}{\left(\frac{1}{4} - 1 + 2\right)^2}$$

$$= \frac{\frac{1}{16} + 1 - 4 + 8 - \frac{6}{16}}{\left(\frac{5}{4}\right)^2} = \frac{\frac{81}{16} - \frac{6}{16}}{\frac{25}{16}} = \frac{75/16}{25/16} = 3$$

Sol 19: (B) $f(x^2) = x^3$

$$\therefore f(x) = x^{3/2}$$

$$\therefore f'(x) = \frac{3}{2}x^{1/2}$$

$$f'(4) = \frac{3}{2} \times 4^{1/2} = 3$$

Sol 20: (D) $x = \sin t, y = \sin 3t$

$$\frac{dy}{dx} = \frac{3\cos 3t}{\cos t} = \frac{3(4\cos^3 t - 3\cos t)}{\cos t}$$

$$= 12\cos^2 t - 9$$

$$\frac{d^2y}{dx^2} = \frac{24\cos t(-\sin t)}{\cos^2 t} = -24\sin t$$

$$\therefore (1 - \sin^2 t)(-24\sin t) - (\sin t)(12\cos^2 t - 9) + k(\sin 3t) = 0$$

$$= -24\sin t + 24\sin^3 t - 12\sin t(1 - \sin^2 t) + 9\sin t + k(3\sin t - 4\sin^3 t) = 0$$

$$\Rightarrow (3k - 36)\sin t + (36 - 4k)\sin^3 t = 0$$

$$\therefore 3k - 27 = 0 \text{ and } 36 = 4k$$

$$\Rightarrow k = 9$$

Sol 21: (B) $x = \ln t, y = t^2 - 1$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2t}{1/t} = 2t^2$$

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dt}\left(\frac{dy}{dx}\right)}{\frac{dx}{dt}} = \frac{4t}{1/t} = 4t^2$$

$$\therefore y''(1) = 4$$

Application of Derivatives


Single Correct Choice Type

Sol 1: (B) $y = ke^{kx} \Rightarrow y' = k^2 e^{kx} \Rightarrow y = k, x = 0$

$$y' = k^2$$

$$\Rightarrow \tan \theta = \frac{1}{k^2} \Rightarrow \cot \theta = k^2$$

$$\Rightarrow \theta = \cot^{-1} k^2$$



Sol 2: (D) $f(x) = \int_2^x (2t - 5) dt = t^2 - 5t \Big|_2^x = x^2 - 5x - 4 + 10$

$$= x^2 - 5x + 6 = (x - 2)(x - 3)$$

$$f'(x) = 2x - 5$$

$$[x = 2, f(x) = 0] \quad f'(x) = -1$$

$$[x = 3, f(x) = 0] \quad f'(x) = 1$$

Angle between the 2 tangents is 90°

(as $m_1 m_2 = -1$)

Sol 3: (C) $x^2 y = c^3$


$$y = \frac{c^3}{x^2} \Rightarrow y' = \frac{-2c^3}{x^3}$$

$$y - \frac{c^3}{t^2} = -\frac{2c^3}{t^3}(x - t) \quad \left[x = t, y = \frac{c^3}{t^2} \right]$$

$$x \text{ intercept} = \frac{3t}{2} = y \text{ intercept}$$

$$= \frac{3c^3}{t^2} = b \Rightarrow a^2 b = \frac{9t^2}{4} \times \frac{3c^3}{t^2} = \frac{27c^3}{4} \quad [C]$$

Sol 4: (D) $f(x) = \begin{cases} x \sin\left(\frac{\pi}{x}\right) & x > 0 \\ 0 & x = 0 \end{cases}$



$$f'(x) = x \cos\left(\frac{\pi}{x}\right) \left(-\frac{\pi}{x^2}\right) + \sin\frac{\pi}{x} = 0 \quad \frac{\pi}{x} = \tan\left(\frac{\pi}{x}\right)$$

$$x \in [0, 1] \text{ infinite solution}$$

$$\text{Sol 5: (D)} \quad y = f(x) \Rightarrow \int_a^b f'(x) f''(x) dx$$

$$\Rightarrow I = f'(x) \int_a^b f''(x) dx - \int_a^b f''(x) f'(x) dx$$

$$\Rightarrow 2I = [f'(x)]^2$$

$$\Rightarrow 2I = [f'(b)]^2 - [f'(a)]^2$$

$$\Rightarrow 2I = -2 \Rightarrow I = -1$$

$$\text{Sol 6: (A)} \quad y = x^3$$

$$y' = 3x^2$$

$$3(x_B)^2 = k^3 (x_A)^2$$

$$\frac{x_B}{x_A} = \pm \sqrt{k}$$

$$\Rightarrow (y - t^3) = 3t^2(x - t) \text{ [at } x = t]$$

$$\Rightarrow x^3 - t^3 = 3t^2(x - t)$$


$$\Rightarrow x^2 + t^2 + xt = 3t^2$$

$$\Rightarrow (x_B)^2 + (x_B x_A) = 2(x_A)^2$$

$$\Rightarrow kx_A^2 \pm \sqrt{k}x_A^2 = 2x_A^2$$

$$\Rightarrow 2 - k = \pm \sqrt{k}$$

$$\Rightarrow k = 4$$



$$\text{Sol 7: (C)} \quad \text{Subnormal} = y \frac{dy}{dx}$$

$$xny^{n-1}y' + y^n = 0$$

$$y' = \frac{-y}{nx}$$

$$\left| y \frac{dy}{dx} \right| = \frac{y^2}{nx} \text{ it is constant for } \Rightarrow \frac{y^{2+n}}{a^{n+1}n}$$


$$\text{Constant for } n = -2$$

$$\text{Sol 8: (A)} \quad y = -\frac{x^2}{2} + 2$$

$$y' = -x$$

$$\Rightarrow \frac{y-2}{x-\frac{1}{2}} = m$$

$$\Rightarrow y-2 = m\left(x-\frac{1}{2}\right) \Rightarrow \frac{-x^2}{2} = \frac{2mx}{2} - \frac{m}{2}$$



$$\Rightarrow x^2 + 2mx - m = 0$$

$$\Rightarrow x = \frac{-2m \pm \sqrt{4m^2 + 4m}}{2} = 0$$

$$\text{For } D = 0$$

$$\Rightarrow 4m^2 + 4m = 0 \Rightarrow m = 0, -1$$

$$\Rightarrow y-2 = -x + \frac{1}{2}$$

$$\Rightarrow x+y = \frac{5}{2}$$

$$\text{Sol 9: (D)} \quad c_1y = x^2 - 3 \text{ and } c_2y = kx^2$$

$$\Rightarrow kx^2 = x^2 - 3$$

$$\Rightarrow x = \pm \sqrt{\frac{3}{1-k}} = a \Rightarrow y = \frac{3}{1-k} - 3 = \frac{3k}{1-k}$$

$$\Rightarrow \frac{y-y_1}{x-a} = 2ka \Rightarrow y-y_1 = 2ka(x-a)$$

$$\Rightarrow x^2 - 3 - y_1 = 2ka(x-a)$$

$$\Rightarrow -2 - y_1 = 2ka(1-a)$$

$$y_2 = -2$$

$$y_1 = ka^2$$


$$\Rightarrow -2 - ka^2 = 2ka - 2ka^2$$

$$\Rightarrow ka^2 - 2ka - 2 = 0$$

$$\Rightarrow k\left(\frac{3}{1-k}\right) - 2 = 2k\sqrt{\frac{3}{1-k}} \Rightarrow \frac{5k-2}{\sqrt{1-k}} = 2k\sqrt{3}$$

$$\Rightarrow 5k-2 = 2k\sqrt{3-3k}$$

$$k = \frac{2}{3}, a = 1$$



$$\text{Sol 10: (B)} \quad x^2 = e^{|x|-2}$$

$$\text{No. of roots are 4}$$

$$\text{Sol 11: (C)} \quad -\frac{a}{x^3} - \frac{b}{y^3} y' = 0 \text{ at any general point}$$

$$\left(t, t\sqrt{\frac{b}{t^2-a}}\right)$$

$$y' = -\frac{ay^3}{bx^3}$$

$$y - t\sqrt{\frac{b}{t^2-a}} = \frac{-at^3}{b} \left(\frac{b}{t^2-a}\right)^{3/2} (x-t)$$


$$\Rightarrow x \text{ intercept } \frac{t-b(t^2-a)+t}{ab} = \frac{bt^3}{ab} = \frac{t^3}{a} = [B]$$

Sol 12: (D) $m = 0$

$$y = c$$

$$y = \sqrt{x}$$

$$y^1 = \frac{1}{2\sqrt{x}} = 1; \quad x = \frac{1}{4}y = \frac{1}{2} = c$$

**Sol 13: (D)** $y^3 - x^2y + 5y - 2x = 0$

$$\Rightarrow x^4 - x^3y^2 + 5x + 2y = 0$$

$$\Rightarrow 3y^2y' - x^2y' - 2xy + 5y' - 2 = 0$$

$$\Rightarrow y' = \frac{2+2xy}{3y^2-x^2+5}, \quad y'(0,0) = \frac{2}{5}$$

$$\Rightarrow 4x^3 - 3x^2y^2 - 2x^3yy' + 5 + 2y' = 0$$

$$\Rightarrow y' = \frac{4x^3 - 3x^2y^2 + 5}{-2 + 2x^3y}, \quad y'(0,0) = -\frac{5}{2}$$

$$\Rightarrow \tan \theta = \frac{\frac{2}{5} + \frac{5}{2}}{1-1} = \infty \quad \therefore \theta = \frac{\pi}{2}$$

Sol 14: (D) $f(x) = \int_0^x \left(t + \frac{1}{t}\right) dt$

$$g(x) = f'(x) = x + \frac{1}{x}$$

$$m = \frac{g(3) - g\left(\frac{1}{2}\right)}{3 - \frac{1}{2}} = \frac{3 + \frac{1}{3} - \frac{1}{2} - 2}{\frac{5}{2}} = \frac{\frac{5}{2} - \frac{5}{3}}{\frac{5}{2}} = \frac{2}{6 \times 1} = \frac{1}{3}$$

$$y' = 1 - \frac{1}{x^2} = \frac{1}{3}$$

$$x = x = \sqrt{\frac{3}{2}} \quad y = \frac{5}{\sqrt{6}}$$

Sol 15: (C) $y = \frac{x^{3/2}}{3}$

$$18yy' = 3x^2 \Rightarrow y' = \frac{x^2}{6y}$$

$$m_{\text{normal}} = \left(\frac{-1}{y^1}\right) = \frac{-6y}{x^2} = \pm 1$$

$$6y = \pm x^2 \Rightarrow y = \frac{x^2}{6}$$

$$\Rightarrow 9 \times \frac{x^4}{36} = x^3 \Rightarrow x = 4 \text{ and } y = \frac{8}{3}$$

Previous Years' Questions

Sol 1: (C) Given, $x = a(\cos \theta + \theta \sin \theta)$ and

$$y = a(\sin \theta - \theta \cos \theta)$$

$$\therefore \frac{dx}{d\theta} = a(-\sin \theta + \sin \theta + \theta \cos \theta)$$

$$= a\theta \cos \theta \quad \text{and} \quad \frac{dy}{d\theta} = a(\cos \theta - \cos \theta + \theta \sin \theta)$$

$$\frac{dy}{d\theta} = a\theta \sin \theta \Rightarrow \frac{dy}{dx} = \tan \theta$$

Thus, equation of normal is

$$\frac{y - a(\sin \theta - \theta \cos \theta)}{x - a(\cos \theta + \theta \sin \theta)} = \frac{-\cos \theta}{\sin \theta}$$

$$\Rightarrow -x \cos \theta + a\theta \sin \theta \cos \theta + a \cos^2 \theta$$

$$= y \sin \theta + \theta a \sin \theta \cos \theta - a \sin^2 \theta$$


$$\Rightarrow x \cos \theta + y \sin \theta = a$$

Whose distance from origin is,

$$\frac{|0+0-a|}{\sqrt{\cos^2 \theta + \sin^2 \theta}} = a$$

Sol 2: (A) Given, $\frac{dy}{dx} = 2x + 1$

on integrating both sides



$$\int dy = \int (2x+1) dx$$

$$\Rightarrow y = x^2 + x + c \text{ which passes through } (1, 2)$$

$$\therefore 2 = 1 + 1 + c \Rightarrow c = 0$$

$$\therefore y = x^2 + x$$

Thus, the required area bounded by x axis, the curve and $x = 1$

$$= \int_0^1 (x^2 + x) dx = \left[\frac{x^3}{3} + \frac{x^2}{2} \right]_0^1$$

$$= \frac{1}{3} + \frac{1}{2} = \frac{5}{6} \text{ sq unit}$$

Sol 3: (D) Slope of tangent $y = f(x)$ is

$$\frac{dy}{dx} = f'(x)_{(3,4)}$$

Therefore, slope of normal

$$= -\frac{1}{f'(x)_{(3,4)}} = -\frac{1}{f'(3)}$$

$$\text{But } -\frac{1}{f'(3)} = \tan\left(\frac{3\pi}{4}\right) \text{ (given)}$$

$$\Rightarrow -\frac{1}{f'(3)} = \tan\left(\frac{\pi}{2} + \frac{\pi}{4}\right) = -1$$

$$f'(3) = 1$$

Sol 4: (D) Given $y^3 + 3x^2 = 12y$

$$\Rightarrow 3y^2 \frac{dy}{dx} + 6x = 12 \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{6x}{12-3y^2} \Rightarrow \frac{dx}{dy} = \frac{12-3y^2}{6x}$$

$$\text{For vertical tangent, } \frac{dx}{dy} = 0$$

$$\Rightarrow 12 - 3y^2 = 0 \Rightarrow y = \pm 2$$

On putting $y = 2$ in Eq. (i), we get $x = \pm \frac{4}{\sqrt{3}}$ and again

putting $y = -2$ in Eq. (i), we get $3x^2 = -16$, no real solution

$$\therefore \text{The required point } \left(\pm \frac{4}{\sqrt{3}}, 2\right)$$

Sol 5: (D) Tangent to the curve $y^2 = 8x$ is,

$$y = mx + \frac{2}{m}$$

So it must satisfy $xy = -1$

$$\Rightarrow x\left(mx + \frac{2}{m}\right) = -1$$

$$\Rightarrow mx^2 + \frac{2}{m}x + 1 = 0,$$


Since it has equal roots, therefore $D = 0$

$$\Rightarrow \frac{4}{m^2} - 4m = 0 \Rightarrow m^3 = 1 \Rightarrow m = 1$$

So equation of common tangent is $y = x + 2$

Sol 6: (A) Let the point be $P(\sqrt{2} \cos \theta, \sin \theta)$

$$\text{on } \frac{x^2}{2} + \frac{y^2}{1} = 1$$



Equation of tangent is,

$$\frac{x\sqrt{2}}{2} \cos \theta + y \sin \theta = 1$$

whose intercept on coordinate axes are $A(\sqrt{2} \sec \theta, 0)$ and $B(0, \csc \theta)$

\therefore Mid point of its intercept between axes $\left(\frac{\sqrt{2}}{2} \sec \theta, \frac{1}{2} \csc \theta\right) = (h, k)$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{2}h} \text{ and } \sin \theta = \frac{1}{2k}$$

Thus, locus of mid point M is

$$(\cos^2 \theta + \sin^2 \theta) = \frac{1}{2h^2} + \frac{1}{4k^2}$$

$$\Rightarrow \frac{1}{2x^2} + \frac{1}{4y^2} = 1, \text{ is required locus}$$

Sol 7: (C) We know, tangent to parabola $y^2 = 4ax$ is y

$$= mx + \frac{a}{m}$$

$$\therefore \text{Tangent to } y^2 = 4x \text{ is } y = mx + \frac{1}{m}$$

Since, tangent passes through $(1, 4)$

$$\therefore 4 = m + \frac{1}{m}$$

$$\Rightarrow m^2 - 4m + 1 = 0$$

(Whose roots are m_1 and m_2)

$$\therefore m_1 + m_2 = 4 \text{ and } m_1 m_2 = 1$$

$$\text{and } |m_1 - m_2| = \sqrt{(m_1 + m_2)^2 - 4m_1 m_2}$$

$$= \sqrt{12} = 2\sqrt{3}$$

Thus, angle between tangents

$$\tan \theta = \left| \frac{m_2 - m_1}{1 + m_1 m_2} \right| = \left| \frac{2\sqrt{3}}{1 + 1} \right| = \sqrt{3} \Rightarrow \theta = \frac{\pi}{3}$$

Sol 8: (D) The tangent at $(1, 7)$ to the parabola $x^2 = y - 6$ is

$$x(1) = \frac{1}{2}(y + 7) - 6$$

(replacing $x^2 \rightarrow x x_1$ and $2y \rightarrow y + y_1$)

$$\Rightarrow 2x = y + 7 - 12$$

$$\Rightarrow y = 2x + 5$$

Which is also tangent to the circle

$$x^2 + y^2 + 16x + 12y + c = 0$$

$$\text{i.e., } x^2 + (2x + 5)^2 + 16x$$

$$+ 12(2x + 5) + c = 0$$

must have equal roots i.e., $\alpha = \beta$

$$\Rightarrow 5x^2 + 60x + 85 + c = 0$$

$$\Rightarrow \alpha + \beta = -\frac{60}{5}$$


$$\Rightarrow a = -6 \Rightarrow x = -6$$

$$\text{and } y = 2x + 5 = -7$$

\therefore Point of contact is $(-6, -7)$

Sol 9: (A) Slope of the line joining the point

$(c-1, e^{c-1})$ and $(c+1, e^{c+1})$ is equal to $\frac{e^{c+1} - e^{c-1}}{2} > e^c$



\Rightarrow Tangent to the curve $y = e^x$ will intersect the given line to the left of the line $x = c$.

Alternate Solution

The equation of the tangent to the curve $y = e^x$ at (c, e^c) is

$$y - e^c = e^c(x - c) \quad \dots (i)$$

Equation of the line joining the given points is

$$y - e^{c-1} = \frac{e^c(e - e^{-1})}{2}[x - (c-1)] \quad \dots (ii)$$

Eliminating y from equation (i) and (ii), we get

$$[x - (c-1)][2 - (e - e^{-1})] = 2e^{-1}$$

$$\Rightarrow x - c = \frac{e + e^{-1} - 2}{2 - (e - e^{-1})} < 0 \Rightarrow x < c$$

Sol 10: (A)

$$f'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$$

$$\Rightarrow f'(1) = \lim_{h \rightarrow 0} \frac{(1+h-1)\sin\left(\frac{1}{1+h-1}\right) - 0}{h}$$


$$= \lim_{h \rightarrow 0} \frac{h}{h} \sin\left(\frac{1}{h}\right)$$

$$\Rightarrow f'(1) = \lim_{h \rightarrow 0} \sin\left(\frac{1}{h}\right)$$

$\therefore f$ is not differentiable at $x = 1$.

$$\text{Similarly, } f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h}$$

$$\Rightarrow f'(0) = \lim_{h \rightarrow 0} \frac{(h-1)\sin\left(\frac{1}{h-1}\right) - \sin(1)}{h}$$



$\Rightarrow f$ is also not differentiable at $x = 0$.

Sol 11: (D)

Solving the equation we get the points of intersection $(-2, 1)$ and $(-2, -1)$

The bounded region is shown as shaded region.

$$\text{The required area} = 2 \int_0^1 (1 - 3y^2) - (-2y^2)$$

$$= 2 \int_0^1 (1 - y^2) dy = 2 \left[y - \frac{y^3}{3} \right]_0^1 = 2 \times \frac{2}{3} = \frac{4}{3}$$

JEE Advanced/Boards

Exercise 1

Methods of Differentiation

Sol 1: $y = x \sin kx$

$$\frac{dy}{dx} = \sin kx + xk \cos kx$$

$$\frac{d^2y}{dx^2} = k \cos kx + k \cos kx - xk^2 \sin kx$$

$$\therefore \frac{d^2y}{dx^2} + y = 2k \cos kx - xk^2 \sin kx + x \sin kx = 2k \cos kx$$

$$\Rightarrow x \sin kx(1 - k^2) = 0$$

$$\therefore k = 1, -1 \text{ when } 1 - k^2 = 0$$

$$\text{and } k = 0 \text{ when } \sin kx = 0$$

Sol 2: Let $f(x) = ax^3 + bx^2 + cx + d$

$$f(2x) = f'(x)f''(x)$$

$$a(2x)^3 + b(2x)^2 + c(2x) + d$$

$$= (3ax^2 + 2bx + c)(6ax + 2b)$$

$$8ax^3 + 4bx^2 + 2cx + d$$

$$= 18a^2x^3 + (6ab + 12ab)x^2$$

$$+ 4b^2x + 6acx + 2bc$$

$$\therefore 8a = 18a^2 \Rightarrow a = 0, \frac{4}{9}$$

$$\therefore f(x) \text{ should be a cubic equation}$$

$$\therefore a \neq 0 \text{ and } a = \frac{4}{9}$$

$$\text{also } 18ab = 4b$$

$$b(18a - 4) = 0 \Rightarrow b \left(18 \times \frac{4}{9} - 4 \right) = 8b = 0 \Rightarrow b = 0$$

$$\text{and } 4b^2 + 6ac = 2c$$

$$\Rightarrow c(6a - 2) = 0 \Rightarrow c = 0$$

$$\therefore f(x) = \frac{4}{9}x^3 \text{ and } d = 2bc = 0$$

Sol 3: $f'(x) = g(x)$; $g'(x) = f(x)$

$$\text{Let } h(x) = f^2(x) - g^2(x)$$

$$h'(x) = 2f(x)f'(x) - 2g(x)g'(x)$$

$$= 2f(x)g(x) - 2g(x)f(x) = 0$$

$\therefore h(x)$ is a constant function whose value is constant for every value of x

$$\therefore h(3) = f^2(3) - g^2(3) = (5)^2 - [f'(3)]^2 = 5^2 - 4^2 = 9$$

$$\therefore f^2(\pi) - g^2(\pi) = 9$$

Sol 4: $3x^2 + 4y^2 = 12 \Rightarrow y^2 = 3 - \frac{3}{4}x^2$

Differentiating both sides, we get

$$2y \frac{dy}{dx} = -\frac{3}{2}x \Rightarrow \frac{dy}{dx} = -\frac{3x}{4y}$$

$$2 \left(\frac{dy}{dx} \right)^2 + 2y \frac{d^2y}{dx^2} = -\frac{3}{2}$$

$$\text{or } 2 \left(\frac{-3x}{4y} \right)^2 + 2y \frac{d^2y}{dx^2} = -\frac{3}{2}$$

$$\frac{18x^2}{16y^2} + 2y \frac{d^2y}{dx^2} = -\frac{3}{2}$$

$$18x^2 + 2y^3 \times 16 \frac{d^2y}{dx^2} = -24y^2$$

$$y^3 \frac{d^2y}{dx^2} = \frac{-24y^2 - 18x^2}{32}$$

$$= -(3x^2 + 4y^2) \times \frac{6}{32} = \frac{-12 \times 6}{32} = -\frac{9}{4}$$

Sol 5: $f(x^2)f''(x) = f'(x)f'(x^2)$... (i)

$$f(1) = 1, f'''(1) = 8$$

$$\text{Find } f'(1) + f''(1)$$

Differentiate the given equation

$$f(x^2)f'''(x) + f'(x^2)f''(x)2x = f''(x)f'(x^2) + f'(x)f''(x^2)2x \quad \dots \text{ (ii)}$$

Put $x = 1$ in equation (1)

$$\Rightarrow f(1)f''(1) = f'(1)f'(1) \Rightarrow f''(1) = [f'(1)]^2 \quad \dots \text{ (iii)}$$

Put $x = 1$ in equation (2)

$$f(1)f'''(1) + f'(1)f''(1) \times 2 = f''(1)f'(1) + f'(1)f''(1)2$$

$$\Rightarrow f''(1)f'(1) = 8 \quad \dots \text{ (iv)}$$

From equation (3) and (4)

$$[f'(1)]^3 = 8$$

$$\therefore f'(1) = 2 \text{ and } f''(1) = (2)^2 = 4$$

$$\therefore f'(1) + f''(1) = 2 + 4 = 6$$

Sol 6: $2x = y^{1/5} + y^{1/5}$

Take $y^{1/5} = t$

$$t^2 - 2xt + 1 = 0$$

$$\therefore t = \frac{2x + \sqrt{4x^2 - 4}}{2} = x + \sqrt{x^2 - 1}$$

$$\therefore y^{1/5} = x + \sqrt{x^2 - 1}$$

$$\frac{1}{5} y^{-4/5} \frac{dy}{dx} = 1 + \frac{x}{\sqrt{x^2 - 1}} = \frac{x + \sqrt{x^2 - 1}}{\sqrt{x^2 - 1}} = \frac{y^{1/5}}{\sqrt{x^2 - 1}}$$

$$\therefore \frac{dy}{dx} = \frac{5y}{\sqrt{x^2 - 1}}$$

$$\frac{d^2y}{dx^2} = \frac{5\sqrt{x^2 - 1} \frac{dy}{dx} - \frac{5yx}{\sqrt{x^2 - 1}}}{(x^2 - 1)}$$

$$= \frac{5 \left[(x^2 - 1) \frac{dy}{dx} - xy \right]}{(x^2 - 1)^{3/2}} (x^2 - 1) \frac{d^2y}{dx^2} + x \frac{dy}{dx}$$

$$= 5\sqrt{x^2 - 1} \frac{dy}{dx} - \frac{5yx}{\sqrt{x^2 - 1}} + \frac{5xy}{\sqrt{x^2 - 1}} = 25y$$

$$\therefore k = 25$$

Sol 7: $\frac{d^2y}{dx^2} = 1 + \frac{2(1+y)}{1+y^2} \left(\frac{dy}{dx} \right)^2$

$$y = \tan z$$

$$\frac{dy}{dx} = \sec^2 z \frac{dz}{dx} \Rightarrow \frac{dz}{dx} = \cos^2 z \frac{dy}{dx}$$

$$\therefore \frac{d^2z}{dx^2} = \cos^2 z \frac{d^2y}{dx^2} - 2\cos z \sin z \frac{dy}{dx} \cdot \frac{dz}{dx}$$

$$= \cos^2 z \left[1 + \frac{2(1 + \tan z)}{\sec^2 z} \left(\sec^4 z \left(\frac{dz}{dx} \right)^2 \right) \right]$$

$$- 2\cos z \sin z \sec^2 z \left(\frac{dz}{dx} \right)^2$$

$$= \cos^2 z + [2(1 + \tan z) - 2\tan z] \left(\frac{dz}{dx} \right)^2$$

$$= \cos^2 z + 2 \left(\frac{dz}{dx} \right)^2 \Rightarrow k = 2$$

Sol 8: $z = \ln \left(\tan \frac{x}{2} \right)$

$$\frac{dz}{dy} = \frac{1}{\tan \frac{x}{2}} \sec^2 \frac{x}{2} \times \frac{1}{2} \frac{dx}{dy} = \operatorname{cosec} x \frac{dx}{dy} \Rightarrow \sin x \frac{dy}{dx} = \frac{dy}{dz}$$

$$\frac{d^2y}{dz^2} = \left[\cos x \frac{dy}{dx} + \sin x \frac{d^2y}{dx^2} \right] \frac{dx}{dz}$$

$$\therefore \frac{d^2y}{dx^2} + \cot x \frac{dy}{dx} = \operatorname{cosec} x \frac{d^2y}{dz^2} \times \frac{dz}{dx} = \operatorname{cosec}^2 x \frac{d^2y}{dz^2}$$

$$\therefore \operatorname{cosec}^2 x \frac{d^2y}{dx^2} + 4y \operatorname{cosec}^2 x = 0$$

$$\Rightarrow \frac{d^2y}{dx^2} + 4y = 0$$

Sol 9: $f(x) = \frac{\sin x}{x}, x \neq 0, f(0) = 1$

$$f'(x) = \begin{cases} \frac{x \cos x - \sin x}{x^2}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

$$f''(0) = \lim_{h \rightarrow 0} \frac{\frac{h \cosh - \sinh}{h^2} - 0}{h}$$

$$= \lim_{h \rightarrow 0} \left(\frac{h \cosh - \sinh}{h^3} \right) \text{ (L-Hospital's rule)}$$

$$= \lim_{h \rightarrow 0} \left(\frac{\cosh - h \sinh - \cosh}{3h^2} \right) = \lim_{h \rightarrow 0} -\frac{1}{3} \left(\frac{\sinh}{h} \right) = -\frac{1}{3}$$

$$\therefore f''(0) = -\frac{1}{3}$$

Sol 10: $R = \frac{\left[1 + \left(\frac{dy}{dx} \right)^2 \right]^{3/2}}{\frac{d^2y}{dx^2}} \Rightarrow R^{2/3} = \frac{1}{\left(\frac{d^2y}{dx^2} \right)^{2/3}} + \frac{1}{\left(\frac{d^2x}{dy^2} \right)^{2/3}}$

$$\text{Let } \frac{dy}{dx} = t \Rightarrow \frac{dx}{dy} = \frac{1}{t}$$

$$\therefore R^{2/3} = \frac{1}{\left(\frac{dt}{dx} \right)^{2/3}} + \frac{1}{\left(\frac{d(1/t)}{dy} \right)^{2/3}}$$

$$= \frac{1}{\left(\frac{dt}{dx}\right)^{2/3}} + \frac{1}{\left[\frac{-1}{t^2} \left(\frac{dt}{dx}\right)\right]^{2/3}} = \frac{1}{\left(\frac{dt}{dx}\right)^{2/3}} + \frac{1}{\left[-\frac{1}{t^3} \frac{dt}{dx}\right]^{2/3}}$$

$$\therefore \frac{dy}{dx} = t = \frac{1}{\left(\frac{dt}{dx}\right)^{2/3}} \left[1 + \frac{1}{1/t^2}\right] = \frac{1}{\left(\frac{dt}{dx}\right)^{2/3}} [1 + t^2]$$

$$\therefore R^{2/3} = \frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]}{\left(\frac{d^2y}{dx^2}\right)^{2/3}} \Rightarrow R = \frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2}}{\left(\frac{d^2y}{dx^2}\right)}$$

Sol 11: $f(x) = \ln(1 + \sqrt{1+x^2})$

$$g(x) = \ln(x + \sqrt{1+x^2}) \Rightarrow e^{g(x)} = (x + \sqrt{1+x^2})$$

$$f\left(\frac{1}{x}\right) = \ln\left(1 + \sqrt{1 + \frac{1}{x^2}}\right) = \ln(x + \sqrt{1+x^2}) - \ln x$$

$$f'\left(\frac{1}{x}\right) = \frac{1}{\sqrt{1+x^2}} - \frac{1}{x}$$

$$\therefore xe^{g(x)} \left(f\left(\frac{1}{x}\right)' + g'(x) \right)$$

$$= x \left(x + \sqrt{1+x^2} \right) \left[\frac{x - \sqrt{1+x^2}}{x\sqrt{1+x^2}} \right] + \frac{1}{\sqrt{1+x^2}}$$

$$= \frac{x^2 - (1+x^2)}{\sqrt{1+x^2}} + \frac{1}{\sqrt{1+x^2}} = 0$$

\therefore For every x , given function is zero

Sol 12: $f\left(\frac{x+y}{k}\right) = \frac{f(x)+f(y)}{k}$

Put $x, y = 0$

$$f(0) = \frac{2f(0)}{k}$$

$$\Rightarrow f(0) \left[\frac{k-2}{k} \right] = 0 \quad \therefore k \neq 2, 0$$

$$\therefore f(0) = 0$$

Put $y = -x$

$$f(0) = \frac{f(x) + f(-x)}{k} = 0$$

$$\therefore f(x) = -f(-x) \text{ or } f(-x) = -f(x)$$

\therefore If above function is satisfying the given condition then the function should be odd or $f(x) = 0$

Sol 13: $f'(x) = 4 \begin{vmatrix} (x-a)^3 & (x-a)^3 & 1 \\ (x-b)^3 & (x-b)^3 & 1 \\ (x-c)^3 & (x-c)^3 & 1 \end{vmatrix}$

$$+ 3 \begin{vmatrix} (x-a)^4 & (x-a)^2 & 1 \\ (x-b)^4 & (x-b)^2 & 1 \\ (x-c)^4 & (x-c)^2 & 1 \end{vmatrix} + \begin{vmatrix} (x-a)^4 & (x-a)^3 & 0 \\ (x-b)^4 & (x-b)^3 & 0 \\ (x-c)^4 & (x-c)^3 & 0 \end{vmatrix}$$

$$\therefore f'(x) = 3 \begin{vmatrix} (x-a)^4 & (x-a)^2 & 1 \\ (x-b)^4 & (x-b)^2 & 1 \\ (x-c)^4 & (x-c)^2 & 1 \end{vmatrix}$$

$$\therefore \lambda = 3$$

Sol 14: $P(x) = ax^4 + bx^3 + cx^2 + dx + e$

$$P(1) = a + b + c + d + e = 0 \quad \dots (i)$$

$$P(3) = 81a + 27b + 9c + 3d + e = 0 \quad \dots (ii)$$

$$P(5) = 625a + 125b + 25c + 5d + e = 0 \quad \dots (iii)$$

$$P'(7) = 4a(7)^3 + 3b(7)^2 + 2c(7) + d = 0$$

$$= 1372a + 147b + 14c + d = 0 \quad \dots (iv)$$

$$(2) - (1) \Rightarrow 80a + 26b + 8c + 2d = 0$$

$$\Rightarrow 40a + 13b + 4c + d = 0 \quad \dots (v)$$

$$(3) - (1) \Rightarrow 624a + 124b + 24c + 4d = 0$$

$$\Rightarrow 156a + 31b + 6c + d = 0 \quad \dots (vi)$$

$$(6) - (5) \Rightarrow 116a + 18b + 2c = 0 \quad \dots (vii)$$

$$(4) - (6) \Rightarrow 1216a + 116b + 8c = 0$$

$$\Rightarrow 304a + 29b + 2c = 0 \quad \dots (viii)$$

$$(8) - (7) \Rightarrow 188a + 11b = 0$$

$$\therefore -\frac{b}{a} = \frac{188}{11}$$

$$\text{also } \frac{-b}{a} = 1 + 3 + 5 + x = \frac{188}{11} \text{ (sum of roots)}$$

$$\therefore x = \frac{188}{11} - 9 = \frac{89}{11}$$

$$\therefore \left(x - \frac{89}{11}\right) \text{ is a root of 4 degree polynomial}$$

$$\therefore p = 89 \text{ } q = 11$$

$$\therefore p + q = 100$$

Alternate:

Take $P(x) = (x-1)(x-3)(x-5)(qx-p)$

Now apply condition that $P'(7) = 0$

Sol 15: $f(x) = x^3 + x^2 f(1) + x f''(2) + f'''(3)$

T.P. $f(2) = f(1) - f(0)$

$f'(x) = 3x^2 + 2x f'(1) + f''(2)$

$f'(1) = 3 + 2f'(1) + f''(2)$

$\therefore f'(1) + f''(2) + 3 = 0$... (i)

$f''(x) = 6x + 2f'(1)$

$f''(2) = 12 + 2f'(1)$... (ii)

$f'''(x) = 6$

$\therefore f'''(3) = 6$... (iii)

$f(2) = 8 + 4f'(1) + 2f''(2) + f'''(3)$

From (1), (2) and (3)

$f'(1) = -5, f''(2) = 2$

$f'''(3) = 6$

$f(1) = 1 + f'(1) + f''(2) + f'''(3) = 1 - 5 + 2 + 6 = 4$

$f(2) = 8 - 20 + 4 + 6 = -2$

$f(0) = f'''(3) = 6$

$\therefore f(2) = f(1) - f(0)$ Hence proved

Sol 16: $f(x) = \sin 2x [\sin(x+x^2)\sin(x-x^2) + \cos(x+x^2)\cos(x-x^2)] + \sin 2x^2 [\cos(x+x^2)\cos(x-x^2) - \sin(x+x^2)\sin(x-x^2)]$

$= \sin 2x \cos(x+x^2+x^2-x) + \sin 2x^2 [\cos(x+x^2+x-x^2)]$

$= \sin 2x \cos 2x^2 + \sin 2x^2 \cos 2x = \sin(2x + 2x^2)$

$\therefore f'(x) = 2(2x+1)\cos 2(x^2+x)$

Sol 17: $f(0) = \begin{vmatrix} a & b & c \\ \ell & m & n \\ p & q & r \end{vmatrix}$

$F'(x) = \begin{vmatrix} 1 & b+x & c+x \\ 1 & m+x & n+x \\ 1 & q+x & r+x \end{vmatrix} + \begin{vmatrix} a+x & 1 & c+x \\ \ell+x & 1 & n+x \\ p+x & 1 & r+x \end{vmatrix}$

$+ \begin{vmatrix} a+x & b+x & 1 \\ \ell+x & m+x & 1 \\ p+x & q+x & 1 \end{vmatrix}$

$f'(x) = (m-b)(r-c) - (n-c)(q-b)$

$+ (-1)[(\ell-a)(r-c) - (n-c)(p-a)]$

$\therefore f''(x) = 0$

$f(x) = \begin{vmatrix} a+x & b+x & c+x \\ \ell-a & x-b & n-c \\ p-a & q-b & r-c \end{vmatrix}$

$(a+x)[(m-b)(r-c) - (q-b)(n-c)]$

$+ (b+x)[(\ell-a)(r-c) - (n-c)(p-a)]$

$+ (c+x)[(\ell-a)(q-b) - (m-b)(p-a)]$

$= a[(m-b)(r-c) - (q-b)(n-c)]$

$+ b[(\ell-a)(r-c) - (n-c)(p-a)]$

$+ c[(\ell-a)(q-b) - (m-b)(p-a)]$

$+ x[(m-b)(r-c) - (q-b)(n-c)]$

$+ \{(\ell-a)(r-c) - (n-c)(p-a) \}$

$+ \{(\ell-a)(a, 0) - (m-b)(p-a)\}$

$= f(0) + kx$

$k =$ sum of all the co-factor of elements of $f(0)$

Sol 18: $y = \sum \tan^{-1} \frac{1}{x^2 + (2n-1)x + \{(n)(n-1) + 1\}}$

$= \sum \tan^{-1} \frac{1}{(x+n)(x+n-1) + 1}$

$= \sum \tan^{-1} \frac{(x+n) - (x+n-1)}{(x+n)(x+n-1) + 1}$

Let $\tan \alpha = x + n$

$\tan \beta = x + n - 1$

$\therefore y = \sum \tan^{-1} \tan(\alpha - \beta) = S\alpha - S\beta$

$y = \sum_{n=1}^n \tan^{-1}(x+n) - \sum_{n=1}^n \tan^{-1}(x+n-1)$

$\therefore y = \tan^{-1}(x+n) - \tan^{-1}x$

$y' = \frac{1}{1+(x+n)^2} - \frac{1}{x^2+1}$

$= \frac{1+x^2-1-(x+n)^2}{(1+x^2)(1+(x+n)^2)} = \frac{x^2-(x+n)^2}{(1+x^2)(1+(x+n)^2)}$

Sol 19: $Y = SX$

$Z = tX$

$Y_1 = SX_1 + S_1X$

$Z_1 = tX_1 + t_1X$

$Y_2 = SX_2 + S_1X_1 + S_2X + S_1X_1$

$$= SX_2 + 2S_1X_1 + S_2X$$

$$Z_2 = tX_2 + 2t_1X_1 + t_2X = 5X_2 + 2S_1X_1 + S_2X$$

$$Z_2 = tX_2 + 2t_1X_1 + t_2X$$

$$\begin{vmatrix} X & Y & Z \\ X_1 & Y_1 & Z_1 \\ X_2 & Y_2 & Z_2 \end{vmatrix}$$

$$= \begin{vmatrix} X & SX & tX \\ X_1 & SX_1 + S_1X & tX_1 + t_1X \\ X_2 & SX_2 + 2S_1X_1 + S_2X & tX_2 + 2t_1X_1 + t_2X \end{vmatrix}$$

$$R_2 \rightarrow R_2 - SX_1, R_3 \rightarrow R_3 - tX_1$$

$$\begin{vmatrix} X & 0 & 0 \\ X_1 & S_1X & t_1X \\ X_2 & 2S_1X_1 + S_2X & 2t_1X_1 + t_2X \end{vmatrix}$$

$$= X[S_1t_2X^2 - S_2t_1X^2] + X_3 \begin{vmatrix} S_1 & t_1 \\ S_2 & t_2 \end{vmatrix}$$

$$\text{Sol 20: } y = \tan^{-1} \frac{x}{\sqrt{1-u^2}}, x = \sec^{-1} \frac{1}{2u^2 - 1}$$

$$\mu = \left(0, \frac{1}{\sqrt{2}}\right) \cup \left(\frac{1}{\sqrt{2}}, 1\right)$$

$$\text{To prove } 2 \frac{dy}{dx} + 1 = 0, \text{ take } u = \cos \theta$$

$$\therefore y = \tan^{-1} \tan \theta = \theta = \cos^{-1} u$$

$$x = \sec^{-1} \frac{1}{2\cos^2 \theta - 1} = \sec^{-1} \sec 2\theta = \pi - 2\theta = \pi - 2\cos^{-1} u$$

$$\therefore \frac{dy}{dx} = \frac{-d\cos^{-1} u}{2d\cos^{-1} u} = \frac{-1}{2}$$

$$\Rightarrow \frac{2dy}{dx} + 1 = 0$$

$$\text{Sol 21: } \sin \left(2 \tan^{-1} \left(\sqrt{\frac{1-x}{1+x}} \right) \right)$$

$$= \frac{2 \tan^{-1} \sqrt{\frac{1-x}{1+x}}}{1 + \left(\tan^{-1} \sqrt{\frac{1-x}{1+x}} \right)^2} = \frac{2 \sqrt{\frac{1-x}{1+x}}}{\frac{2}{(1+x)}} = \sqrt{1-x^2}$$

$$\therefore y = \tan^{-1} \left(\frac{x}{\sqrt{1-x^2} + 1} \right) + \sqrt{1-x^2}$$

$$\therefore \text{ Put } x = \sin \theta$$

$$\therefore \frac{x}{\left(\sqrt{1-x^2}\right)+1} = \frac{\sin \theta}{1+\cos \theta} = \tan \frac{\theta}{2}$$

$$\therefore y_1 = \tan^{-1} \left(\tan \frac{\theta}{2} \right) = \frac{\theta}{2} = \frac{1}{2} \sin^{-1} x$$

$$\therefore y = \frac{1}{2} \sin^{-1} x + \sqrt{1-x^2}$$

$$y' = \frac{1}{2\sqrt{1-x^2}} + \frac{1(-2x)}{2\sqrt{1-x^2}}$$

$$= \frac{1}{2\sqrt{1-x^2}} - \frac{x}{\sqrt{1-x^2}} = \frac{1-2x}{2\sqrt{1-x^2}}$$

$$\text{Sol 22: } y = \cot^{-1} \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}}$$

$$= \cot^{-1} \frac{(\sqrt{1+\sin x} + \sqrt{1-\sin x})^2}{1+\sin x - (1-\sin x)}$$

$$= \cot^{-1} \frac{(2 + 2\sqrt{1-\sin^2 x})}{2\sin x} = \cot^{-1} \left(\frac{1+\cos x}{\sin x} \right)$$

$$= \cos^{-1} \cot \frac{x}{2}, x \in \left(0, \frac{\pi}{2}\right) = \frac{x}{2}, x \in \left(0, \frac{\pi}{2}\right)$$

$$\frac{dy}{dx} = \frac{1}{2} \text{ or } \cot^{-1} \cot \left(\frac{\pi}{2} - \frac{x}{2} \right) = \frac{\pi}{2} - \frac{x}{2}, x \in \left(\frac{\pi}{2}, \pi \right)$$

$$\therefore \frac{dy}{dx} = -\frac{1}{2}$$

$$\text{Sol 23: } x = \phi(t), y = \psi(t)$$

$$\frac{dx}{dy} = \phi'(t) \frac{dy}{dt} = \psi'(t)$$

$$\frac{d^2x}{dt^2} = \phi''(t) \frac{d^2y}{dt^2} = \psi''(t)$$

$$\frac{dy}{dx} = \frac{\psi'(t)}{\phi'(t)} - \frac{d^2y}{dx^2} = \frac{d\left(\frac{dy}{dx}\right)}{\frac{dx}{dt}}$$

$$\frac{d^2y}{dx} = \frac{d\frac{\psi'(t)}{\phi'(t)}}{\frac{dx}{dt}} = \frac{\phi'(t)\psi''(t) - \psi'(t)\phi''(t)}{[\phi'(t)]^2\phi'(t)}$$

$$= \frac{\left(\frac{dx}{dt}\right)\left(\frac{d^2y}{dt^2}\right) - \left(\frac{d^2x}{dt^2}\right)\frac{dy}{dt}}{\left(\frac{dx}{dt}\right)^3}$$

Sol 24: (a) $e^{xy} + y \cos x = 2$

Differentiate the equation w.r.t. x

$$e^{xy} \left[y + x \frac{dy}{dx} \right] + \cos x \frac{dy}{dx} - y \sin x = 0$$

$$\frac{dy}{dx} (xe^{xy} + \cos x) = y \sin x - ye^{xy}$$

$$\therefore \frac{dy}{dx} = \frac{y \sin x - ye^{xy}}{xe^{xy} + \cos x}$$

$$\left. \frac{dy}{dx} \right|_{x=0} = -y$$

at $x = 0$; $y = 1$

$$(i) \quad \therefore \left. \frac{dy}{dx} \right|_{x=0} = -1$$

also

$$\frac{d^2y}{dx^2} (xe^{xy} + \cos x) + \frac{dy}{dx} (e^{xy} + xe^{xy} (y + x \frac{dy}{dx}) - \sin x)$$

$$= y \cos x + \frac{dy}{dx} \sin x - ye^{xy} \left(y + x \frac{dy}{dx} \right) - \frac{dy}{dx} e^{xy}$$

$$(ii) \quad \left. \frac{d^2y}{dx^2} \right|_{x=0} = \frac{y - y^2 - \frac{dy}{dx} - \frac{dy}{dx}(1)}{(0+1)}$$

$$= \frac{1 - 1 - (-1) - (-1)}{1} = 2$$

$$(b) g(x) = e^{ax} + f(x)$$

$$g'(x) = ae^{ax} + f'(x)$$

$$g''(x) = a^2e^{ax} + f''(x)$$

$$g'(x) + g''(x) = (a + a^2)e^{ax} + f'(x) + f''(x)$$

$$\therefore g'(0) + g''(0) = a + a^2 + f'(0) + f''(0) = 0$$

$$\Rightarrow a + a^2 - 5 + 3 = 0 \Rightarrow a^2 + a - 2 = 0 \Rightarrow a = 1, -2$$

Application of Derivatives

Sol 1: $y - 2 = m(x - 1)$

$$\text{curve } (y - 2)^2 = 2x^3 - 4$$

$$\text{tangent } 2(y - 2) \frac{dy}{dx} = 6x^2$$

$$\left. \frac{dy}{dx} \right|_{h,k} = \frac{3h^2}{k - 2}$$

$$\frac{k - 2}{h - 1} = \frac{3h^2}{k - 2}$$

$$(k - 2)^2 = 3h^2(h - 1)$$

$$\Rightarrow 2h^3 - 4 = 3h^3 - 3h^2$$

$$\Rightarrow 3h^2 - 4 = h^3 \Rightarrow h^3 - 3h^2 + 4 = 0$$

$$\Rightarrow (h + 1)(h^2 - 4h + 4) = 0$$

$$h = -1, 2$$

$$h = 2, \Rightarrow k = 2 \pm \sqrt{12}$$

Eq. of tangent

$$y - 2 = \pm \frac{\sqrt{12}}{1}(x - 1)$$

Sol 2: $y = ax^2 + bx + \frac{7}{2}$ at $(1, 2)$

Now at $(1, 2)$, we will get

$$2 = a + b + \frac{7}{2} \quad \dots\dots(i)$$

The tangent will be

$$y' = 2ax + b$$

$$y = x^2 + 6x + 10$$

$$y'_{(-2,2)} = 2(-2) + 6 = 2$$

$$m_{\text{normal}} = \frac{-1}{2}$$

$$2a(1) + b = \frac{-1}{2} \Rightarrow 4a + 2b = -1$$

$$\Rightarrow a + b = \frac{-3}{2}$$

$$\Rightarrow 2a = 2; a = 1$$

$$b = -\frac{5}{2}$$

Sol 3: $xy = 1 - y$

$$x^2y = xy \Rightarrow y(x^2 - x) = 0$$

$$y = 0 \mid x = 0 \mid x = 1$$

$$A(0, 1), B\left(1, \frac{1}{2}\right)$$

$$x^2 y' + 2xy = -y' \Rightarrow y' = -\frac{2xy}{x^2 + 1}$$

$$y'_{(0,1)} = 0, y'_{\left(1, \frac{1}{2}\right)} = \frac{-1}{2}$$

Equation of tangents

$$\frac{y - \frac{1}{2}}{x - 1} = \frac{-1}{2} \Rightarrow y = -\frac{x}{2} + 1$$

This gives $y = 1, x = 0$ or $(0, 1)$

Sol 4:

$$y' = y(1+x)^{y-1} \left[\begin{array}{l} \text{if } y = (+x)^y \\ \ln y = y \ln 1 + y \\ y' = yy' \ln(1+x) + \frac{y^2}{1+x} \end{array} \right]$$

$$y' = \frac{(1+x)^{2y}}{(1+x) \left[1 - (1+x)^y \ln(1+x) \right]} + \frac{2 \sin x \cos x}{\sqrt{1 + \sin^4 x}}$$

$$\text{at } x = 0, y' = 1 + 0 = 1$$

$$m_{\text{normal}} = -1$$

$$\text{Equation of normal} \Rightarrow \frac{y-1}{x-0} = -1 \Rightarrow x+y=1$$

Sol 5: $x = 2t + t^2 \sin\left(\frac{1}{t}\right), \quad t \neq 0$

$$= 0 \quad t = 0$$

$$y = \frac{\sin t^2}{t} \quad t \neq 0$$

$$= 0 \quad t = 0$$

$$x' = 2 + t^2 \cos\left(\frac{1}{t}\right) \left(\frac{-1}{t^2}\right) + 2t \sin\left(\frac{1}{t}\right)$$

$$= 2 - \cos \frac{1}{t} + 2t \sin\left[\frac{1}{t}\right]$$

$$y' = \frac{2t^2 \cos t^2 - \sin t^2}{t^2} = 2 \cos t^2 - \frac{\sin t^2}{t^2}$$

$$\frac{y'}{x'} = \frac{2t^2 \cos t^2 - \sin t^2}{t^2 \left(2 - \cos \frac{1}{t} + 2t \sin \frac{1}{t} \right)}$$

$$\text{Slope at } (t=0) = \frac{2t^2 \cos t^2 - \sin t^2}{t^2 \left[1 + 2t \sin \frac{1}{t} + 2 \sin^2 \frac{1}{2t} \right]}$$

Does not exist

Sol 6: $y' = 41x^2 = 2009$

$$x^2 = \frac{2009}{41} = t^2 = 49$$

$$\left(y - \frac{41t^3}{3} \right) = 2009(x - t)$$

$$b - \frac{41t^3}{3} = -2009t$$

$$b = \frac{41t^3}{3} - 2009t = \frac{t}{3} (41t^2 - 2009.3)$$

$$= 7 \times \left(\frac{41 \times 49}{3} - 2009 \right) = -9375.33$$

Sol 7: $y^2 = -\sin(x+y)[1+y']$

$$y^1 = -\frac{\sin(x+y)}{1+\sin(x+y)} = -\frac{1}{2}$$


$$\sin(x+y) = 1, \cos(x+y) = 0$$

$$\text{i.e. } y = 0$$

$$x = \frac{\pi}{2}, -\frac{3\pi}{2}$$

$$\frac{y}{x - \pi/2} = -\frac{1}{2} \& \frac{y}{x + \frac{3\pi}{2}} = -\frac{1}{2}$$

$$2y + x = \frac{\pi}{2} \quad 2y + x = -\frac{3\pi}{2}$$



Sol 8: $q = p^2 \quad P > 0$

$$s = -\frac{8}{r} \quad r > 0, s < 0$$

$$y - t^2 = 2t(x - t) - \text{tangent to curve (1) at } x = t$$

$$y + \frac{8}{z} = \frac{8}{z^2}(x - z) - \text{tangent to curve (2) at } x = z$$

$$\text{Both pass through } (p, q) \quad (r, s)$$

$$y = 2tx - t^2 \quad t = p$$

$$\text{Same tangent}$$

$$y = \frac{8x}{z^2} - \frac{16}{z} \quad z = r$$

$$tz^2 = 4t^2z = 16$$

$$t^2z^4 = 16$$

$$z = 1, t = 4$$

$$z + t = p + r = 5$$

Sol 9: (a) $y = \sqrt{36.6}$

$$y = \sqrt{x} \Rightarrow y' = \frac{1}{2\sqrt{x}}$$

$$x = 36, \Delta x = 0.6$$

$$\Rightarrow f(x + \Delta x) = f(x) + f'(x) \Delta x$$

$$\Rightarrow f(36.6) = f(36) + \frac{1}{2\sqrt{36}} \times 0.6 = 6 + \frac{0.6}{120} = 6.05$$

(b) $(26)^{1/3}$

$$y = (x)^{1/3}$$

$$y' = \frac{1}{3}x^{-2/3} \Rightarrow \Delta x = -1$$

$$f(26) = f(27) + \frac{1}{3} \times \frac{(-1)}{27^{2/3}} = 3 - \frac{1}{3 \times 9} = 3 - \frac{1}{27} = \frac{80}{27}$$

(ii) $r = 9 \pm 0.03$

$$v = \frac{4}{3}\pi r^3$$

$$\frac{dv}{v} = \frac{4\pi r^2 dr}{v} = \frac{3dr}{R} = \frac{3 \times 0.03}{9} = \frac{1}{100}$$

$$dv = \frac{4}{3}\pi r^3 \times \frac{1}{100} = \frac{4\pi}{300} \times 9 \times 9 \times 9 = 9.72 \pi$$

Sol 10: Mid point was (2, -1)

$$\frac{y+1}{x-2} = 1 \Rightarrow x - y = 3 \text{ (equation of tangent)}$$

$$x - 3 = -a^2x^2 + 5ax - 4$$

$$\Rightarrow a^2x^2 + x(1 - 5a) + 1 = 0$$

$$\alpha + \beta \left(\frac{1-5a}{-a^2} \right) = \frac{5a-1}{a^2}$$

$$\frac{\alpha + \beta}{2} = 2 = \frac{5a-1}{2a^2} \Rightarrow 4a^2 - 5a + 1 = 0$$


$$a = 1, + \frac{1}{4}$$

Sol 11: $P_1(\alpha, \alpha^3)$

$$y - \alpha^3 = 3\alpha^2(x - \alpha) \Rightarrow y - 3\alpha^2x - 2\alpha^3 = 0$$

$$x^3 - 3\alpha^2x + 2\alpha^3 = 0$$

$$x_1 + x_2 + x_3 = 0$$



$$x_1 = x_2 = \alpha$$

$$x_3 = 2\alpha \Rightarrow P_2(-2\alpha, -8\alpha^3)$$

$$(\alpha, -2\alpha, 4\alpha, \dots) \text{ forms a G.P.}$$

Tangent at P_2

$$\frac{y + 8\alpha^3}{x + 2\alpha} = 12\alpha^2$$

$$y = 12\alpha^2x + 16\alpha^3$$

$$\alpha_1 + \alpha_2 + \alpha_3 = 0$$

$$\alpha_1 + \alpha_2 = -2\alpha$$

$$\alpha_3 = 4\alpha \Rightarrow P_3(4\alpha, 64\alpha^3)$$

$$\text{Area of } P_1P_2P_3 = \frac{1}{2} \begin{vmatrix} \alpha & \alpha^3 & 1 \\ -2\alpha & -8\alpha^3 & 1 \\ 4\alpha & 64\alpha^3 & 1 \end{vmatrix}$$

$$\text{Area of } P_2P_3P_1 = \frac{1}{2} \begin{vmatrix} -2\alpha & -8\alpha^3 & 1 \\ 4\alpha & 64\alpha^3 & 1 \\ -8\alpha & -512\alpha^3 & 1 \end{vmatrix}$$

$$\begin{aligned} \frac{P_1P_2P_3}{P_2P_3P_4} &= \frac{\alpha(-72\alpha^3) - \alpha^3(-6\alpha) + 1(-128 + 32)\alpha^4}{-2\alpha(576) + 8\alpha^3(12\alpha) + 1(-2048 + 512)\alpha^4} \\ &= \frac{-72 + 6 - 96}{96 - 1536 - 1152} = \frac{162}{2592} = \frac{1}{16} \end{aligned}$$

Sol 12: $f'(x) = (fx)^2$

$$\int \frac{dy}{y^2} = \int dx - \frac{1}{y} = x + c$$

For $f(0) = \frac{-1}{2}$, We have $c = 2$

$$\Rightarrow y = \frac{-1}{x+2}$$

$$\frac{y + \frac{1}{2}}{x} = \frac{1}{4}$$


$$4y + 2 = x$$

Sol 13: $y = ax^3 + bx^2 + cx + 5$

$$y' = 3ax^2 + 2bx + c$$

$$y'(-2) = 12a - 4b + c = 0$$

$$-8a + 4b - 2c + 5 = 0$$



$$c = 3 [y^1 (x = 0) = 3]$$

$$12a - 4b + 3 = 0$$

$$-8a + 4b = 1$$

$$4a = -2 \Rightarrow a = -\frac{1}{2} \text{ and } b = \frac{-3}{4}$$

$$\text{Sol 14: } y - t^3 = 3t^2(x - t)$$

$$\Rightarrow 8 - t^3 = 3t^2(2 - t) \Rightarrow 8 - t^3 = 6t^2 - 3t^3$$

$$\Rightarrow 2t^3 - 6t^2 + 8 = 0$$

$$t = -1, 2, 2$$

$$m = 3t^2, (m)_{-1} = 3, (m)_2 = 12$$

$$\text{Sol 15: } f^3(x) = \int_0^x t f^2(t) dt$$

$$3f^2(x) = f'x = xf^2x$$

$$f^2x(3f'x - x) = 0$$

Either $f(x) = 0$ (not possible)

$$\text{or } f'x = \frac{x}{3} M_{\text{normal}} = \frac{-3}{x} - \frac{1}{2} \Rightarrow x = 6$$

$$7x = \frac{x^2}{6} + c$$

Equation of normal

$$y - 6 - c = \frac{-1}{2}(x - 6)$$

Intercept on y axis

$$y = 9 + c$$

$$\left(\frac{x^2}{6} + c\right)^3 = \int_0^x \left(\frac{x^2}{6} + c\right)^2 x = \left(\frac{x^4}{36} + c^2 + \frac{x^2}{3}\right)x$$

$$= \int \frac{x^5}{36} + c^2x + \frac{x^3}{3} \Big|_0^x$$

$$\frac{x^6}{6.36} + c^3 + \frac{3c^2x^2}{6} + \frac{3x^4c}{36} = \frac{x^6}{6.36} + \frac{c^2x^2}{2} + \frac{x^4}{12} = c = 0$$

Intercept is 9

$$\text{Sol 16: } [y - f(p)] = f'(p)(x - p)$$

$$-f(p) = f'(p)2$$

$$\Rightarrow f'p = -\frac{1}{2}f(p) \Rightarrow \frac{f'p}{fp} = -\frac{1}{2}$$

$$\ln fp = \frac{-x}{2} + c \Rightarrow fp = ce^{-x/2}$$

Passes through (0, 2)

$$f(0) = c = 2$$

$$f(p) = 2e^{-x/2}$$

Sol 17: (a) Similar to exercise (3) q.7

$$(b) y = a \ln(x^2 - a^2)$$

$$\left| \frac{y\sqrt{1+(y')^2}}{y'} \right| + \left| \frac{y}{y'} \right|$$

$$y' = \frac{2ax}{x^2 - a^2}$$

$$\sqrt{1+(y')^2} = \frac{x^2 + a^2}{x^2 - a^2}$$

$$\frac{y}{y'} = \frac{\ln(x^2 - a^2)(x^2 - a^2)}{(2x)}$$

$$\Rightarrow \frac{x^2 - a^2}{2x} \ln(x^2 - a^2) \left[\frac{x^2 + a^2}{x^2 - a^2} + 1 \right]$$

$$\Rightarrow x \ln(x^2 - a^2) = \frac{xy}{a} \text{ i.e. } \boxed{\alpha xy}$$

$$x = y^2, xy = k$$

$$y' = \frac{1}{2y} \quad y' = \frac{-k}{x^2}$$

$$y^3 = k \text{ intersection is } (k^{2/3}, k^{1/3})$$

$$y'_1 \times y'_2 = -1$$

$$\frac{-k}{2x^2y} = -1$$

$$k = 2x^2y = 2y^5$$

$$k = 2k^{5/3}$$

$$k^{-2/3} = 2 \Rightarrow k^{-2} = 8$$

$$k = \pm \frac{1}{2\sqrt{2}}$$



$$\text{Sol 18: } x + 5y - y^5 = 0(0, 0) (0, 5^{1/4})$$

$$1 + 5y' - 5y^4y' = 0$$

$$y' = \frac{1}{5y^4 - 5}$$

Equation of tangent

$$= y = \frac{-1x}{5}$$



Equation of normal $y = 5x$

Coordinates are $(0, 0)$ $(-25, 5)$, $(1, 5)$

$$\text{Area} = \frac{1}{2} \times 5 \times 26 = 65$$

Sol 19: $\frac{y - \frac{1}{2}}{x - 2} = 4$

$$y = 4x - \frac{15}{2} \Rightarrow \frac{1}{x} = 4x - \frac{15}{2}$$

$$\Rightarrow 8x^2 - 15x - 2 = 0 \Rightarrow 8x^2 - 16x + x - 2 = 0$$

$$\Rightarrow (8x + 1)(x - 2) = 0 \Rightarrow x = -\frac{1}{8}$$

$$y' = \frac{-1}{x^2} = -64$$

$$|y'| = 64$$

Sol 20: $f(x) = \ln^2 x + 2 \ln x$

$$y = m_1 x$$

$$y' = \frac{2 \ln x + 2}{x}$$

$$y - (\ln^2 t + 2 \ln t)$$

$$= \frac{2}{t} (\ln t + 1) [x - t]$$

It passes through $(0, 0)$

$$-\ln^2 t - 2 \ln t = -2(1 + \ln t)$$

$$\ln^2 t = 2$$

$$\ln t = \pm \sqrt{2}$$

$$t = e^{\sqrt{2}}, e^{-\sqrt{2}} \text{ ab} = 1$$

$$(ii) 5x \left[\frac{2 \ln x + 2}{x} \right] - x \ln 10 - 10 = 0$$

$$10 \ln x + 10 - x \ln 10 - 10 = 0$$

$$\boxed{10 \ln x = x \ln 10}$$

2 solution from graph

1 is $x = 10$

Sol 21: Given that $6y = x^3 + 2$ and also $dy = 8dx$


$6dy = 3x^2 dx$ (Differentiating the given equation)

$$\Rightarrow \frac{x^2}{2} = 8 \Rightarrow x = \pm 4$$

$$y = \pm \frac{64 + 2}{6} = \frac{-62}{6} \text{ or } 11$$

$$\text{Hence } (4, 11) \left(-4, \frac{-31}{3} \right)$$

Sol 22:




$$\frac{1.5}{y} = \frac{4.5}{y + x} \Rightarrow 3y = 1.5x$$

$$y = \frac{x}{2} \frac{dy}{dx} = \frac{1}{2} \frac{dx}{dt} \Rightarrow \frac{dy}{dt} = \frac{1}{2} \frac{dx}{dt} \Rightarrow \frac{dy}{dt} = 2$$

$$\frac{dx}{dt} + \frac{dy}{dt} = \frac{12}{2} = 6$$

hcc shadow is lightening at rate $\frac{dy}{dt}$ i.e. 2 km/hr

Sol 23: $\frac{dv}{dt} = -1 \text{ cm}^3 / \text{sec}$



$$\frac{dv}{dt} = c \text{ cm}^3 / \text{secc}, \tan \theta = \frac{3}{2} = \frac{r}{h}$$

$$\frac{dh}{dt} = 4 \frac{dr}{dt} = \frac{3}{2} \frac{dh}{dt} = 6$$

$$v = \frac{1}{3} \pi r^2 h$$

$$\frac{dv}{dt} = \frac{\pi}{3} r^2 \frac{dh}{dt} + \frac{\pi h}{3} (2r) \frac{dr}{dt}$$

$$c - 1 = \frac{\pi r^2 4}{3} + \frac{2\pi}{3} h r \frac{dr}{dt}$$

$$h = 2r = 3$$

$$c - 1 = \frac{\pi}{3} \times 9 \times 4 + \frac{2\pi}{3} \times 6.6 = 12\pi + 24\pi = 36\pi$$

$$c = 1 + 36\pi$$

Sol 24: $\frac{dv}{dt} = -2$

$$v = \frac{1}{3}\pi r^2 h$$


$$r = h = \frac{l}{\sqrt{2}}$$

$$v = \frac{\pi}{3} \frac{l^3}{2\sqrt{2}}$$

$$\frac{dv}{dt} = \frac{\pi}{6\sqrt{2}} 3l^2 \frac{dl}{dt}$$

$$-2 = \frac{\pi \times 16}{2\sqrt{2}} \frac{dl}{dt}$$

$$\frac{dl}{dt} = -\frac{1}{\pi 2\sqrt{2}} = \frac{-\sqrt{2}}{4\pi}$$



Sol 25: $h = \frac{1}{6}r$

$$\Rightarrow \int dv = \frac{1}{3}\pi r^2 h \Rightarrow v = \frac{1}{3}\pi 36h^3$$

$$\int dv = 12\pi h^3$$

$$\frac{dv}{dt} = 36\pi h^2 \frac{dh}{dt}$$

$$\Rightarrow 12 = 36\pi h^2 \times \frac{dh}{dt}$$

$$\Rightarrow \frac{1}{3\pi \times 16} = \frac{dh}{dt} = \frac{1}{48\pi} \text{ cm/sec}$$

Sol 26: $\frac{dA}{dt} = 2 \text{ cm}^2 / \text{sec}$

$$A = \pi r^2$$

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt} \Rightarrow 2 = 2\pi r \frac{dr}{dt}$$

$$\Rightarrow \frac{dr}{dt} = \frac{1}{\pi r} \Rightarrow r dr = \frac{dt}{\pi}$$

$$\Rightarrow \frac{r^2}{2} = \frac{1}{\pi} \frac{28}{\pi} \Rightarrow \frac{7.28}{11.11} = r^2$$

$$\Rightarrow r = \frac{14}{\pi}$$

$$\frac{dr}{dt} = \frac{7 \times 11}{22 \times 14} = \frac{1}{4} \text{ cm/sec}$$

Sol 27: A (0, 0)

$$C \left(t, 1 + \frac{7t^2}{36} \right)$$

B (0, 1) initially

Co-ordinate of B at time (0, 1 + 2t)

Co-ordinate of C at time

$$1 + \frac{7x^2}{36} = 1 + 2t$$

$$x = 6\sqrt{\frac{2t}{7}}$$


$$BC = x = 6\sqrt{\frac{2t}{7}}$$

$$AB = 1 + 2t$$

$$\text{Area} = \frac{1}{2}(AB)(BC) = 3(1 + 2t)\sqrt{\frac{2t}{7}} = 3\sqrt{\frac{2t}{7}} + 6t\sqrt{\frac{2t}{7}}$$

$$\frac{dA}{dt} = \frac{3}{\sqrt{7t}} + \frac{3}{2} \times 6\sqrt{\frac{2t}{7}}$$

$$\left. \frac{dA}{dt} \right|_{t=\frac{7}{2}} = 3\sqrt{2} + 9$$



Sol 28:

$$\frac{dv}{dt} = \frac{k}{r}$$

$$v = \frac{4}{3}\pi r^3$$

$$\frac{dv}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$4\pi r^2 \frac{dr}{dt} = \frac{k}{r} \Rightarrow \frac{dr}{dt} = \frac{k}{4\pi r^3}$$

$$\Rightarrow \pi r^4 \Big|_1^2 = k t \Big|_0^{15} \Rightarrow \pi 15 = 15k$$


$$\Rightarrow k = \pi$$

$$\Rightarrow \pi r^4 \Big|_1^t = \pi \Big|_0^t$$

$$\Rightarrow \pi(r^4 - 1)\pi t$$

$$\Rightarrow r = (1 + t)^{1/4}$$

$$\Rightarrow \frac{dv}{dt} = \frac{k}{(1 + t)^{1/4}}$$



$$\Rightarrow dv = \pi(1+t)^{-1/4} dt$$

$$\Rightarrow \int_v^{27v} v = \frac{4\pi}{3}(1+t)^{3/4} \Big|_0^t$$

$$\Rightarrow 26v = \frac{4\pi}{3} \left[(1+t)^{3/4} - 1 \right]$$

$$\Rightarrow v = \frac{4\pi}{3} \Rightarrow 27 = (1+t)^{3/4}$$

$$\Rightarrow (1+t)^{1/4} = 3 \Rightarrow t = 80 \text{ sec}$$

Sol 29: $\frac{dv}{dt} = 6m^3 / \text{min}$

$$v = \frac{\pi}{3} y^2 (3R - y)$$

$$\frac{dv}{dt} = \frac{\pi}{3} \left[6Ry \frac{dy}{dt} - 3y^2 \frac{dy}{dt} \right] - 6 = \pi [2Ry - y^2] y'$$

For $y = 8$, $y' = \frac{-6}{8\pi(2R - y)} = \frac{-6}{8\pi(18)} = -\frac{1}{24\pi} \text{ m/min}$

$$\tan \theta = \frac{r}{y}$$

$$\frac{dr}{dt} = \frac{dy}{dt} \tan \theta = \frac{-1 \times 5}{24\pi \times 12} = -\frac{5}{288\pi}$$

Exercise 2

Methods of Differentiation

Single Correct Choice Type

Sol 1: (D) $y = \frac{x}{a + \frac{x}{b+y}}$

$$\Rightarrow ay + \frac{xy}{b+y} = x$$

$$aby + ay^2 + xy = xb + xy$$

$$\therefore aby + ay^2 = xb$$

$$\Rightarrow ab \frac{dy}{dx} + 2ay \frac{dy}{dx} = b$$

$$\therefore \frac{dy}{dx} = \frac{b}{2ay + ab}$$

Sol 2: (B) $f(x) = e^x + x$

$$f'(x) = 1 + e^x$$

Also $f(f^{-1}(x)) = x$

$$f'(f^{-1}(x)) (f^{-1}(x))' = 1$$

$$\therefore (f^{-1}(x))' = \frac{1}{f'[f^{-1}(x)]}$$

$$\therefore f(\ln 2) = y$$

$$\therefore f^{-1}(y) = \ln 2$$

$$Qf'(1\ln 2) = 1 + e^{\ln 2} = 1 + 2 = 3$$

$$\therefore [f^{-1}(y)]' = \frac{1}{f'[f^{-1}(y)]} = \frac{1}{f'(\ln 2)} = \frac{1}{3}$$

Sol 3: (C) $y(x) = f^2(x) + g^2(x)$

$$y'(x) = 2f(x) g(x) + 2g(x) g'(x)$$

$$= 2f(x) g(x) - 2f(x) g(x) = 0$$

$$\therefore y(x) = a = f^2(x) + g^2(x)$$

$$y(5) = a = f^2(5) + g^2(5) = (2)^2 + (2)^2$$

$$\therefore a = 8$$

$$\therefore y(10) = f^2(10) + g^2(10) = 8$$

Sol 4: (B) $y(x) = x^{\left[\left(\frac{\ell+m}{m-n} \right) \frac{1}{n-\ell} + \left(\frac{m+n}{n-\ell} \right) \frac{1}{\ell-m} + \left(\frac{n+\ell}{\ell-m} \right) \frac{1}{m-n} \right]}$

$$= x^{\left(\frac{1}{n-\ell} \right) \left[\frac{\ell+m}{m-n} + \frac{m+n}{\ell-m} \right] + \left(\frac{n+\ell}{\ell-m} \right) \frac{1}{m-n}}$$

$$= x^{\left(\frac{1}{n-\ell} \right) \left[\frac{\ell^2 - m^2 + m^2 - n^2}{(m-n)(\ell-m)} \right] + \frac{(n+\ell)}{(\ell-m)} \times \frac{1}{(m-n)}}$$

$$= x^{\frac{-(\ell+n)}{(m-n)(\ell-m)} + \frac{(n+\ell)}{(\ell-m)(m-n)}} = x^0 = 1$$

$$\therefore \frac{dy}{dx} = 0$$

Sol 5: (D) $f(x) = (x^x)^x g(x) = x^{(x^x)}$

$$\log f(x) = x \log x^x = x^2 \log x$$

$$\frac{1}{f(x)} f'(x) = 2x \log x + x$$

$$f'(x) = (x^x)^x [2x \log x + x]$$

$$\log(x) = x^x \log x$$

$$\frac{1}{g(x)} g'(x) = \frac{x^x}{x} + \log x \frac{dx^x}{dx}$$

$$= x^{x-1} + (\log x) (x^x (\log x + 1))$$

$$\therefore g'(x) = x^{\left(x^x \right)} \left[x^{x-1} + x^x (\log x + 1) \log x \right]$$

$$f'(1) = [2.1 \cdot \log 1 + 1] (1^1)^1 = 1$$

$$g'(1) = (1)^{(1^1)} [1^{1-1} + 1^1(\log 1 + 1)\log 1] = 1$$

Sol 6: (C) $y^{1/m} + y^{-1/m} = 2x$

Let $y^{1/m} = a$

$$\therefore a + \frac{1}{a} = 2x \Rightarrow a^2 - 2ax + 1 = 0$$

$$\Rightarrow a = x + \sqrt{x^2 - 1}$$

$$y^{1/m} = x + \sqrt{x^2 - 1}$$

$$\therefore y = \left(x + \sqrt{x^2 - 1}\right)^m$$

$$\Rightarrow y' = m \left(x + \sqrt{x^2 - 1}\right)^{m-1} \left(1 + \frac{2x}{2\sqrt{x^2 - 1}}\right) = \frac{m \left(x + \sqrt{x^2 - 1}\right)^m}{\sqrt{x^2 - 1}}$$

$$\Rightarrow y' \sqrt{x^2 - 1} = my$$

$$\Rightarrow y'' y' \sqrt{x^2 - 1} + \frac{y' 2x}{2\sqrt{x^2 - 1}} = my'$$

$$\Rightarrow y''(x^2 - 1) + xy' = my' \sqrt{x^2 - 1}$$

$$\therefore \frac{y''(x^2 - 1) + xy'}{y} = m \sqrt{x^2 - 1} \frac{y'}{y}$$

$$= m \sqrt{x^2 - 1} \times \frac{m}{\sqrt{x^2 - 1}} = m^2$$

Sol 7: (C) $y^2 = P(x)$

$$2y \frac{dy}{dx} = P'(x)$$

$$\Rightarrow 2yy'' + 2(y')^2 = P''(x)$$

Multiply this equation by y^2

$$\Rightarrow 2y^3y'' + 2(yy')^2 = P''(x)y^2 = P''(x)P(x)$$

$$\therefore 2 \frac{d}{dx} \left(y^3 \frac{d^2y}{dx^2} \right) = [P''(x)P(x) - 2(yy')^2]'$$

$$= [P''(x)P(x)]' - 2 \left[\frac{[P'(x)]^2}{4} \right]'$$

$$= P'''(x)P(x) + P''(x)P'(x) - P'(x)P''(x) = P'''(x)P(x)$$

Sol 8: (D) $f(x) = -\frac{x^3}{3} + x^2 \sin 1.5a - x \cdot \sin a \cdot \sin 2a$

$$-5 \sin^{-1}(a^2 - 8a + 17)$$

$$f'(x) = -x^2 + 2x \sin 1.5a - \sin a \sin 2a$$

$$f'(\sin 8) = -\sin^2 8 + 2 \sin 8 \sin 1.5a - \sin a \sin 2a$$

$$\text{Also } -1 \leq a^2 - 8a + 17 \leq 1$$

$$-1 \leq (a - 4)^2 + 1 \leq 1$$

$$-2 \leq (a - 4)^2 \leq 0$$

$$\Rightarrow a = 4$$

$$\therefore f'(\sin 8) = -\sin^2 8 + 2 \sin 8 \sin 6$$

$$- \sin 4 \sin 8$$

$$f'(\sin 8) = -\sin^2 8 + \sin 8(2 \sin 6 - \sin 4)$$

$$\therefore |2 \sin 6| < |\sin 4| \pi < 4 < 6 < 2\pi$$

$$\text{and } \sin 8 < 0$$

$$\therefore f'(\sin 8) < 0$$

Sol 9: (B) $y = \frac{\cos 6x + 6 \cos 4x + 15 \cos 2x + 10}{\cos 5x + 5 \cos 3x + 10 \cos x}$

$$\cos 6x + \cos 4x + 5 \cos 4x + 5 \cos 2x$$

$$+ 10(\cos 2x + 1)$$

$$\Rightarrow 2 \cos \frac{10x}{2} \cos \frac{2x}{2}$$

$$+ 2 \times 5 \cos \frac{(4+2)x}{2} \cos \frac{x}{2} + 10 \times 2 \cos^2 x$$

$$= 2 \cos x [\cos 5x + 5 \cos 3x + 10 \cos x]$$

$$\therefore y = 2 \cos x$$

$$\therefore \frac{dy}{dx} = -2 \sin x$$

Sol 10: (D) $y = R(1 - \cos \theta)$

$$x = R(\theta - \sin \theta)$$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\sin \theta}{(1 - \cos \theta)}$$

$$= \frac{1 + \cos \theta}{\sin \theta} = \operatorname{cosec} \theta + \cot \theta$$

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{d\theta} \left(\frac{dy}{dx} \right)}{\frac{dx}{d\theta}}$$

$$= \frac{-\operatorname{cosec} \theta \cot \theta - \operatorname{cosec}^2 \theta}{R(1 - \cos \theta)}$$

$$= -\operatorname{cosec} \theta \frac{(1 + \cos \theta)}{\sin \theta} \times \frac{1}{R} \frac{(1 + \cos \theta)}{\sin^2 \theta}$$

$$= -\frac{1}{\sin^4 \theta} (1 + \cos \theta)^2 \times \frac{1}{R}$$

$$= -\frac{1}{R} \left(\frac{1 + \cos \theta}{\sin^2 \theta} \right)^2 = \frac{-1}{R} \left(\frac{1}{1 - \cos \theta} \right)^2$$

$$\therefore \frac{d^2 y}{dx^2} \bigg|_{\theta=\pi} = \frac{-1}{R} \left(\frac{1}{1 - (-1)} \right)^2 = -\frac{1}{4R}$$

Sol 11: (B) $f(x) = (1 + x)^n$

$$f'(x) = n(1 + x)^{n-1}$$

$$f''(x) = n(n-1)(1 + x)^{n-2}$$

$$f^n(x) = n(n-1) \dots \dots 2.1 (1+x)^0$$

$$f(0) = 1, f'(0) = n, f''(0) = n(n-1)$$

$$f(0) + f'(0) + \dots + \frac{f^n(0)}{n!}$$

$$= 1 + n + \frac{n(n-1)}{2!} \dots \dots \frac{n(n-1) \dots (2.1)}{n!}$$

$$= {}^nC_0 + {}^nC_1 + {}^nC_2 + \dots + {}^nC_n = 2^n$$

Sol 12: (D) $y = e^{4x} + 2e^{-x}$

$$\frac{dy}{dx} = 4e^{4x} - 2e^{-x}$$

$$\frac{d^2 y}{dx^2} = 16e^{4x} + 2e^{-x}$$

$$\frac{d^3 y}{dx^3} = 64e^{4x} - 2e^{-x}$$

$$\frac{d^3 y}{dx^3} - 13 \frac{dy}{dx} = 64e^{4x} - 2e^{-x} - 13(4e^{4x} - 2e^{-x})$$

$$= 12e^{4x} + 24e^{-x} = 12y$$

$$\therefore K = 12$$

Sol 13: (C) $x^4 + 3x^2y^2 + 7xy^3 + 4x^3y - 15y^4 = 0$

$$\Rightarrow (x - y)(x^3 + 5x^2y + 8xy^2 + 5y^3) = 0$$

This is in the form $f(x, y) g(x, y) = 0$ where $f(x, y) = 0$ and $g(x, y) \neq 0$ at $P(x_1, y_1)$

$$\Rightarrow f'(x, y) g(x, y) + f(x, y) g'(x, y) = 0 \Rightarrow f'(x_1, y_1) = 0$$

Also

$$f''(x, y) g(x, y) + 2 f'(x, y) g'(x, y) + f(x, y) g''(x, y) = 0$$

$$\Rightarrow f''(x_1, y_1) = 0 \Rightarrow \frac{d^2 y}{dx^2} = 0 \text{ at } (1, 1)$$

Sol 14: (C) $f(x) = e^{e^x} g(x) = f^{-1}(x), f(g(x)) = x$

$$f'(g(x)) g'(x) = 1$$

$$g'(x) = \frac{1}{f'(g(x))}$$

$$g(2) \Rightarrow 2 = e^{e^x} \Rightarrow x = \ln(\ln 2)$$

$$\therefore g(2) = \ln \ln 2$$

$$\therefore g'(2) = \frac{1}{f'(g(2))} = \frac{1}{f'(\ln \ln 2)}$$

$$f'(x) = e^x \cdot e^{e^x}$$

$$\therefore f'(\ln \ln 2) = e^{\ln \ln 2} \cdot e^{e^{\ln \ln 2}} = 2 \ln 2$$

Sol 15: (C)

$$y = \tan^{-1} \left(\frac{1 - 2 \ln |x|}{1 + 2 \ln |x|} \right) + \tan^{-1} \left(\frac{3 - 2 \ln |x|}{1 - 6 \ln |x|} \right)$$

$$\text{Let } 3 = \tan \alpha, 2 \ln |x| = \tan \beta$$

$$\therefore y = \tan^{-1} \left(\tan \left(\frac{\pi}{4} - \beta \right) \right) + \tan^{-1}(\tan(\alpha + \beta))$$

$$= \frac{\pi}{4} - \beta + \alpha + \beta$$

$$y = \frac{\pi}{4} + \alpha = \frac{\pi}{4} + \tan^{-1} 3$$

$$\therefore \frac{dy}{dx} = 0$$

Sol 16: (B) $\lim_{x \rightarrow 0^+} (x^{x^x} - x^x)$

$$x^x = e^{x \ln x}$$

$$\therefore \lim_{x \rightarrow 0} x^x = \lim_{x \rightarrow 0} e^{x \ln x} = \lim_{x \rightarrow 0} e^{\left(\frac{\ln x}{\frac{1}{x}} \right)}$$

$$= \lim_{x \rightarrow 0} e^{\frac{(\ln x)'}{\left(\frac{1}{x} \right)'}} = \lim_{x \rightarrow 0} e^{\frac{\left(\frac{1}{x} \right)}{\left(-\frac{1}{x^2} \right)}} = \lim_{x \rightarrow 0} e^{-x} = 1$$

$$\lim_{x \rightarrow 0^+} x^{x^x} = \lim_{x \rightarrow 0^+} (x)^{(x^x)} = \lim_{x \rightarrow 0^+} (x)^1 = 0$$

$$\therefore \lim_{x \rightarrow 0^+} (x^{x^x} - x^x) = 0 - 1 = -1$$

Sol 17: (B) $\lim_{x \rightarrow 0} \{ (\cot x)^x + (1 - \cos x)^{\operatorname{cosec} x} \}$

$$\lim_{x \rightarrow 0} e^{x \ln \cot x} + \lim_{x \rightarrow 0} e^{\frac{\ln(1-\cos x)}{\sin x}}$$

$$\lim_{x \rightarrow 0} e^{\frac{\ln \cot x}{\left(\frac{1}{x}\right)}} + \lim_{x \rightarrow 0} e^{\operatorname{cosec} x \ln(1-\cos x)}$$

$\lim_{x \rightarrow 0} e^{\operatorname{cosec} x \ln(1-\cos x)}$ doesn't exist as LHL \neq RHL

Multiple Correct Choice Type**Sol 18: (B, C)**

$$f(x) = \tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{\tan \theta} \right)$$

Put $x = \tan \theta$

$$\therefore f(x) = \tan^{-1} \left(\frac{\sqrt{1+\tan^2 \theta}-1}{\tan \theta} \right),$$

 $\tan \theta \neq 0$

$$= \tan^{-1} \left(\frac{\sec \theta - 1}{\tan \theta} \right) = \tan^{-1} \left(\frac{1 - \cos \theta}{\sin \theta} \right)$$

$$= \tan^{-1} \tan \left(\frac{\theta}{2} \right) = \frac{\theta}{2} = \frac{\tan^{-1} x}{2}$$

$$\therefore f(-x) = \frac{\tan^{-1}(x)}{2} = -\frac{\tan^{-1} x}{2} = -f(x)$$

 $\therefore f(x)$ is an odd function

$$\text{also } f'(x) = \left(\frac{1}{2} \tan^{-1} x \right)' = \frac{1}{2(1+x^2)}, x \in \mathbb{R} - \{0\}$$

Sol 19: (A, B, C)

$$y = \tan x \tan 2x \tan 3x$$

$$x+2x-3x = 0$$

$$\Rightarrow \tan x + \tan 2x - \tan 3x = \tan x \tan 2x \tan 3x = y$$

$$\Rightarrow y' = 3 \sec^2 3x - \sec^2 x - 2 \sec^2 2x$$

$$\text{or } y' = (\tan 3x)' \tan x \tan 2x + (\tan x)' \tan 2x \tan 3x + (\tan 2x)' \tan x \tan 3x$$

$$= 3 \sec^2 3x \tan x \tan 2x + \sec^2 x \tan 2x \tan 3x + 2 \sec^2 2x \tan x \tan 3x$$

$$= \frac{3 \tan x \tan 2x \tan 3x}{\sin 3x \cos 3x} + \frac{\tan x \tan 2x \tan 3x}{\sin x \cos x}$$

$$+ \frac{2 \tan x \tan 2x \tan 3x}{\sin 2x \cos 2x}$$

$$= 2 \tan x \tan 2x \tan 3x \times \left[\frac{3}{2 \sin 3x \cos 3x} + \frac{1}{2 \sin x \cos x} + \frac{2}{2 \sin 2x \cos 2x} \right]$$

$$= 2y [2 \operatorname{cosec} 6x + 2 \operatorname{cosec} 4x + \operatorname{cosec} 2x]$$

Sol 20: (A, C, D)

$$y = \sqrt{x \sqrt{x + \sqrt{x + \dots \infty}}}$$

$$\Rightarrow y = \sqrt{x+y}$$

$$\Rightarrow y^2 - y = x \Rightarrow (y-1) = \frac{x}{y}$$

$$\therefore 2y \frac{dy}{dx} - \frac{dy}{dx} = 1$$

$$\therefore \frac{dy}{dx} = \frac{1}{2y-1} = \frac{1}{2(y-1)+1} = \frac{1}{2 \cdot \frac{x}{y} + 1} = \frac{y}{2x+y}$$

$$\text{also } y = \frac{1 \pm \sqrt{1+4x}}{2}$$

$$\therefore \frac{dy}{dx} = \frac{1}{2 \frac{(1 \pm \sqrt{1+4x})}{2} - 1} = \frac{1}{\pm \sqrt{1+4x}}$$

Sol 21: (A, B, C, D)

$$2^x + 2^y = 2^{x+y}$$

$$2^x \ln 2 + 2^y \ln 2 \frac{dy}{dx} = 2^{x+y} \ln 2 \left(1 + \frac{dy}{dx} \right)$$

$$(2^y - 2^{x+y}) \frac{dy}{dx} = 2^{x+y} - 2^x$$

$$\frac{dy}{dx} = \frac{2^x(2^y-1)}{2^y(1-2^x)} = \frac{2^x(1-2^y)}{2^y(2^x-1)}$$

$$\text{Also } 2^{x+y} - 2^y = 2^y(2^x-1) = 2^x$$

$$\text{or } 2^{x+y} - 2^x = 2x(2^y-1) = 2^y$$

$$\therefore \frac{dy}{dx} = -\frac{2^x}{2^y} \text{ or } \frac{2^y(2^x-1)}{2^x} = 1$$

$$\therefore \frac{dy}{dx} = (1-2^y) \text{ or } \frac{2^x(2^y-1)}{2^y} = 1$$

$$\therefore \frac{dy}{dx} = -\frac{1}{(2^x-1)} = \frac{1}{1-2^x}$$

Sol 22: (A, B, C)

$$\sqrt{y+x} + \sqrt{y-x} = c$$

$$\frac{1}{2\sqrt{y+x}} \left(\frac{dy}{dx} + 1 \right) + \frac{1}{2\sqrt{y-x}} \left(\frac{dy}{dx} - 1 \right) = 0$$

$$\frac{dy}{dx} \left(\frac{\sqrt{y-x} + \sqrt{y+x}}{\sqrt{y^2 - x^2}} \right) = \frac{\sqrt{y+x} - \sqrt{y-x}}{\sqrt{y^2 - x^2}}$$

Application of Derivatives**Single Correct Choice Type**

Sol 1: (C) $3x^2 + 4xy + 5y^2 - 4 = 0$

$$y' = -\frac{(3x+2y)}{(2x+5y)}$$

$$y' = 0 \text{ when } y = -\frac{3}{2}x$$

$$\text{and } y' = \infty \text{ when } y = -\frac{2}{5}x$$

So angle is 90° .

Sol 2: (D) $x = \sec^2 t, y = \cot t$

$$\Rightarrow x = 1 + \frac{1}{y^2}$$

$$\text{At } t = \frac{\pi}{4}, x = 2, y = 1$$

$$P(2, 1)$$

$$\Rightarrow \frac{y-1}{x-2} = y' = \frac{-1}{2} \Rightarrow 2y-2=2-x \Rightarrow x+2y=4$$

$$\Rightarrow 4-2y = \frac{y^2+1}{y^2} \text{ putting the value of } x \text{ in first equation}$$

$$\Rightarrow 4y^2 - 2y^3 = y^2 + 1 \Rightarrow 2y^3 - 3y^2 + 1 = 0$$

$$\Rightarrow (y-1)(2y^2 - y - 1) = 0 \Rightarrow (y-1)(2y+1)(y-1)$$

$$y = -\frac{1}{2}; x = 5; \left(5, -\frac{1}{2}\right)$$

$$PQ = \sqrt{9 + \frac{9}{4}} = \sqrt{\frac{45}{4}} = \frac{3\sqrt{5}}{2}$$

Sol 3: (C)
$$f(x) = \begin{cases} x^{3/5} & x \leq 1 \\ -(x-2)^3 & x > 1 \end{cases}$$

$$f'(x) = \begin{cases} \frac{3}{5}x^{-2/5} & x \leq 1 \\ -3(x-2)^2 & x > 1 \end{cases}$$

$$x = 1, \frac{3}{5}, -3$$

$x = 1$ is critical point

$$f'(x) = \begin{cases} \frac{-9}{25}x^{-8/5} & x \leq 1 \\ -6(x-2) & x > 1 \end{cases}$$

$$\left. \begin{matrix} x=0 \\ x=2 \end{matrix} \right\} \text{critical point}$$

at $x = 2$, $f''(2)$ changes its sign

Sol 4: (A) $x = a(2 \cos t - \cos 2t)$

$$y = a(2 \sin t - \sin 2t)$$

$$\frac{dy}{dt} = a(2 \cos t - 2 \cos 2t)$$

$$\frac{dx}{dt} = a(-2 \sin t + 2 \sin 2t)$$

$$\Rightarrow \frac{dy}{dx} = \frac{\cos t - \cos 2t}{\sin 2t - \sin t} = \frac{\cos t - 2 \cos^2 t + 1}{2 \sin t \cos t - \sin t} = 0$$

$$\Rightarrow 2 \cos^2 t - \cos t - 1 = 0$$

$$\Rightarrow 2 \cos^2 t - 2 \cos t + \cos t - 1 = 0$$


$$\Rightarrow (2 \cos t + 1)(\cos t - 1)$$

$$\cos t = 1, \cos t = \frac{-1}{2}$$


$$t = 0, 2t, t = \frac{2\pi}{3}, \frac{4\pi}{3}$$

$t = 0$ is not possible as $\frac{dy}{dx}$ is not defined

$$t = \frac{4\pi}{3} - \frac{2\pi}{3} = \frac{2\pi}{3}$$

Sol 5: (C)

$$\frac{y-a^n}{x-a} = m = \frac{-1}{y^1} = \frac{-1}{nx^{n-1}}$$



$$y^1 = nx^{n-1}$$

$$y = a^n + \frac{a}{na^{n-1}}$$

$$b = a^n + \frac{aa^{1-n}}{n} a^n + \frac{a^{2-n}}{n} = \frac{na^{2n} + a^2}{na^n}$$

$$\lim_{a \rightarrow 0} b = \frac{2n^2 a^{2n-1} + 2a}{n^2 a^{n-1}} = \frac{2n^2 (2n-1) a^{2n-2} + 2}{n^2 (n-1) a^{n-2}},$$

value of b exist & equal to $\frac{1}{2}$

only if (n = 2)

Sol 6: (B) $f(x) = \begin{cases} -x^2 & x < 0 \\ x^2 + 8 & x \geq 0 \end{cases}$

For points (a, -a²)

$$\frac{y+a^2}{x-a} = -2a$$

$$\Rightarrow y + a^2 = 2a^2 - 2ax$$

$$\Rightarrow y = a^2 - 2ax = x^2 + 8$$

$$\Rightarrow x^2 + 2ax - a^2 + 8 \Rightarrow 4a^2 = -4(8 - a^2)$$

At a = -2

$$\Rightarrow \frac{y+4}{x+2} = +4$$

Therefore x intercept = -1

Sol 7: (D) $y = \frac{1}{2 + \cos^2 x}$

$$\Rightarrow y' = -\frac{[2\cos x \sin x]}{(2 + \cos^2 x)^2}$$

$$x = \frac{\pi}{2} | 0 | \pi \text{ For } x=0 \text{ or } x=\frac{\pi}{2}$$

$$\Rightarrow y = \frac{1}{3} \left(\text{if } x=0 \right), \frac{1}{2} \left(\text{if } x=\frac{\pi}{2} \right)$$

Sol 8: (A) At P (a, b), the equation will be

$$\frac{y-b}{x-a} = M \text{ where M is the slope}$$

$$\Rightarrow \frac{n}{a} \left(\frac{x}{a} \right)^{n-1} + \frac{n}{b} \left(\frac{y}{b} \right)^{n-1} y' = 0$$

$$\Rightarrow \frac{n}{a} + \frac{n}{a} y' = 0$$

$$\Rightarrow y' = \frac{-b}{a}$$

$$\Rightarrow \frac{y-b}{x-a} = \frac{-b}{a}$$

$$\Rightarrow ay + bx = 2ab$$

$$\Rightarrow \frac{x}{a} + \frac{y}{b} = 2$$

Multiple Correct Choice Type

Sol 9: (A, D) We can write

$$xy = k \quad \dots (i)$$

$$\frac{x}{a} + \frac{y}{b} = 1 \quad \dots (ii)$$

Solving these two equations, we get

$$x^2b + abx + ka = 0$$

$$\text{For } D = 0, (ab)^2 - 4abk = 0$$

$$ab(ab - 4k) = 0$$

$$\Rightarrow ab + 4k$$

$$\Rightarrow ab > 0$$

Hence a > 0, b > 0 or a < 0, b < 0

Sol 10: (A, B) $\sqrt{xy} = a + x$

$$\frac{(xy' + y)}{2\sqrt{xy}} = 1$$

$$y' = \frac{2\sqrt{xy} - y}{x}$$

$$\frac{y - \frac{(a+t)^2}{t}}{x-t} = \frac{2\sqrt{t} \left(\frac{a+t}{\sqrt{t}} \right) - \frac{(a+t)^2}{t}}{t}$$

$$\frac{y - \frac{(a+t)^2}{t}}{x-t} = \frac{2 \frac{(a+t)t}{t} - \frac{(a+t)(a+t)}{t}}{t}$$

$$= \frac{a+t}{t} \left[1 - \frac{a}{t} \right]$$

$$= \frac{a+t}{t^2} (t-a) = \frac{t^2 - a^2}{t^2}$$

x intercept will be

$$\Rightarrow t - \frac{(t+a)^2 t^2}{t(t^2 - a^2)} = \frac{t - (t+a)t}{(t-a)} = \frac{-2at}{t-a}$$

y intercept will be

$$= \frac{(a+t)^2}{t} - \frac{(t^2 - a^2)}{t} = \frac{2a^2 + 2at}{t} = \frac{2at}{a-t}$$

$$= a^2 - t^2 = t^2 \Rightarrow a^2 = 2t^2$$

$$t = \frac{a}{\sqrt{2}}, -\frac{a}{\sqrt{2}} [A, B]$$

Sol 11: (B, D) $2x - 3 = x^2 + px + q$

$$p + q + 1 = -1$$

$$p + q = -2$$

$$y = \left[q - \frac{p^2}{4} \right] \text{ is minimum}$$

$$= -2 - p - \frac{p^2}{4}$$

$$\Rightarrow -p - \frac{2p}{4} = 0$$

Therefore $\boxed{p = -2}$
 $\boxed{a = 0}$

Least distance is $\left| 0 - \frac{4}{4} \right| = 1$ D is correct

Sol 12: (A, B) Given that $f(x) = \frac{x^3}{3} - \frac{5x^2}{2} + 7x - 4$

$$f'(x) = x^2 - 5x + 7$$

Equation of triangle will be

$$\frac{y - \left(\frac{t^3}{3} - \frac{5t^2}{2} + 7t - 4 \right)}{x - t} = t^2 - 5t + 7$$

x intercept will be

$$x - t = \frac{-\left(\frac{t^3}{3} - \frac{5t^2}{2} + 7t - 4 \right)}{t^2 - 5t + 7}$$

$$\Rightarrow x = -\frac{\left(\frac{t^3}{3} - \frac{5t^2}{2} + 7t - 4 \right)}{t^2 - 5t + 7} + t$$

y intercept will be

$$y = -t(t^2 - 5t + 7) + \left(\frac{t^3}{3} - \frac{5t^2}{2} + 7t - 4 \right)$$

Equating (i) and (ii) and keeping them opposite in sign

$$-\left(\frac{t^3}{3} - \frac{5t^2}{2} + 7t - 4 \right) = t(t^2 - 5t + 7) - \left(\frac{t^3}{3} + \frac{5t^2}{2} - 7t + 4 \right)$$

Solving above, we get $t = 2, 3$

Therefore, co-ordinates are $\left(2, \frac{8}{3} \right) \left(3, \frac{7}{2} \right)$

Sol 13: (A, B, D)

$$y \cot x = y^3 \tan x$$

$$y^2 = \cot^2 x$$

$$\Rightarrow y = \cot x, -\cot x$$

$$\left(-\frac{\pi}{4}, -1 \right) \left(-\frac{\pi}{4}, +1 \right)$$

$$\frac{y+1}{x + \frac{\pi}{4}} = \frac{-1}{\frac{1}{2}} \Rightarrow y+1 = -2x - \frac{\pi}{2} \Rightarrow 4x + 2y = 2 + \pi$$

$$\frac{y-1}{x + \frac{\pi}{4}} = \frac{1}{\frac{1}{2}} \Rightarrow y-1 = 2x + \frac{\pi}{2} = 4x - 2y = 2 + \pi$$

Sol 14: (A, B)

$$x = a (t + \sin t \cos t)$$

$$y = a (1 + \sin t)^2$$

$$y' = \frac{2a(1 + \sin t) \cos t}{a(1 - \sin^2 t + \cos^2 t)}$$

$$= \frac{2(1 + \sin t) \cos t}{2 \cos^2 t}$$

$$\tan \theta = \frac{1 + \sin t}{\cos t}$$

$$\theta = \tan^{-1} \left(\frac{1 + \sin t}{\cos t} \right) = \frac{\pi + 2t}{4}$$

Sol 15: (B, D) $y = t^3 - 4t^2 - 3t$

$$y = 2t^2 + 3 - 5$$

$$\frac{dy}{dt} = 4t + 3$$

$$\frac{dx}{dt} = 3t^2 - 8t - 3$$

... (i)

... (ii)

$$\frac{dy}{dt} = \frac{4t+3}{3t^2-8t-3}$$

$$\frac{dy}{dx} = 0, t = \frac{-3}{4}, H=1$$

$$\frac{dy}{dx} = \text{not defined at } 3t^2 - 8t - 3 = 0$$

$$3t^2 - 9t + t - 3 = 0$$

$$(3t+1)(t-3) = 0$$

$$t = 3, \frac{-1}{3}$$

$$v = 2$$

Previous Years' Questions

Sol 1: (B, C)

Given, $xy = 1 \Rightarrow y = \frac{1}{x} \Rightarrow \frac{dy}{dx} = -\frac{1}{x^2}$

Thus, slope of normal = x^2 (Which is always positive) and it is given $ax + by + c = 0$ is normal whose slope = $-\frac{a}{b}$

$$\Rightarrow -\frac{a}{b} > 0 \text{ or } \frac{a}{b} < 0,$$

$\therefore a$ and b are of opposite sign.

Sol 2: (B, D) Given, $4x^2 + 9y^2 = 1$

On differentiating w.r.t. x , we get

$$8x + 18y \frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = -\frac{8x}{18y} = -\frac{4x}{9y}$$

The tangent at point (h, k) will be parallel to $8x = 9y$,

$$\text{then } -\frac{4h}{9k} = \frac{8}{9}$$

$$\Rightarrow h = -2k$$

Point (h, k) also lies on the ellipse

$$\therefore 4h^2 + 9k^2 = 1$$

On putting value of h in Eq. (ii), we get

$$4(-2k)^2 + 9k^2 = 1$$

$$\Rightarrow 16k^2 + 9k^2 = 1$$

$$\Rightarrow 25k^2 = 1$$

$$\Rightarrow k^2 = \frac{1}{25}; k = \pm \frac{1}{5}$$

Thus, the point where the tangents are parallel to $8x = 9y$ are $\left(-\frac{2}{5}, \frac{1}{5}\right)$ and $\left(\frac{2}{5}, -\frac{1}{5}\right)$.

Therefore, (b) and (d) are true answers.

Sol 3: Given, $y^3 - 3xy + 2 = 0$

$$\Rightarrow 3y^2 \frac{dy}{dx} - 3x \frac{dy}{dx} - 3y = 0$$

$$\Rightarrow \frac{dy}{dx} (3y^2 - 3x) = 3y$$

$$\Rightarrow \frac{dy}{dx} = \frac{3y}{3y^2 - 3x}$$

Thus, the point where tangent is horizontal. The slope of tangent is 0

$$\therefore \frac{dy}{dx} = 0 \Rightarrow \frac{3y}{3y^2 - 3x} = 0$$

$\Rightarrow y = 0$ but $y = 0$ does not satisfy the given equation of the curve therefore y cannot lie on the curve.

So, $H = \phi$ (null set)

For the point where tangent is vertical, then $\frac{dy}{dx} = \infty$

$$\Rightarrow \frac{y}{y^2 - x} = \infty$$

$$\Rightarrow y^2 - x = 0 \Rightarrow y^2 = x$$


On putting this value in the given equation of the curve, we have

Sol 4: Let the house of the swimmer be at B

$$\therefore AB = L \text{ km}$$

Let the swimmer land at C, on the shore and let

$$AC = x \text{ km}$$



$$\therefore SC = \sqrt{x^2 + d^2} \text{ and } CB = (L - x)$$

$$\therefore \text{Time} = \frac{\text{Distance}}{\text{Speed}}$$

Time from S to B = time from S to C + time from C to B.

$$\therefore T = \frac{\sqrt{x^2 + d^2}}{u} + \frac{L - x}{v}$$

$$\text{Let } f(x) = T = \frac{1}{u} \sqrt{x^2 + d^2} + \frac{L}{v} - \frac{x}{v}$$

$$\Rightarrow f'(x) = \frac{1}{u} \cdot \frac{1.2x}{2\sqrt{x^2 + d^2}} + 0 - \frac{1}{v}$$

For maximum or minimum,

$$\text{Put } f'(x) = 0$$

$$\Rightarrow v^2 x^2 = u^2(x^2 + d^2)$$

$$\Rightarrow x^2 = \frac{u^2 d^2}{v^2 - u^2}$$

$$\therefore f'(x) = 0$$

$$\text{at } x = \pm \frac{ud}{\sqrt{v^2 - u^2}}, (v > u)$$

$$\text{But } x \neq \frac{-ud}{\sqrt{v^2 - u^2}}$$

$$\therefore \text{ We consider } x = \frac{ud}{\sqrt{v^2 - u^2}}$$

$$\text{Now, } f''(x) = \frac{1}{u} \frac{d^2}{\sqrt{x^2 + d^2} (x^2 + d^2)} > 0 \text{ for all } x$$

$$\therefore f \text{ has minimum at } x = \frac{ud}{\sqrt{v^2 - u^2}}$$

$$\text{Sol 5: Given } y = \frac{x}{1+x^2}$$

$$\Rightarrow \frac{dy}{dx} = \frac{(1+x^2) \cdot 1 - x(2x)}{(1+x^2)^2} = \frac{1-x^2}{(1+x^2)^2}$$

$$\text{Let } \frac{dy}{dx} = g(x) \text{ (i.e. slope of tangent)}$$

$$\therefore g(x) = \frac{1-x^2}{(1+x^2)^2}$$

$$\begin{aligned} \Rightarrow g'(x) &= \frac{(1+x^2)^2 \cdot (-2x) - (1-x^2) \cdot 2(1+x^2) \cdot 2x}{(1+x^2)^4} \\ &= \frac{-2x(1+x^2)[(1-x^2) + 2(1-x^2)]}{(1+x^2)^4} = \frac{-2x(3-x^2)}{(1+x^2)^3} \end{aligned}$$

For greatest or least values of m we should have

$$g'(x) = 0 \Rightarrow x = 0, x = \pm\sqrt{3}$$

Now,

$$g''(x) = \frac{(1+x^2)^3(6x^2-6) - (2x^3-6x) \cdot 3(1+x^2)^2 \cdot 2x}{(1+x^2)^6}$$

$$\text{At } x = 0, g''(x) = -6 < 0$$

$$\therefore g'(x) \text{ has maximum value at } x = 0$$

$\Rightarrow (x = 0, y = 0)$ is the required point at which tangent to the curve has the greatest slope.

$$\text{Sol 6: Given, } y = \cos(x + y)$$

$$\Rightarrow \left(\frac{dy}{dx}\right) = -\sin(x + y) \cdot \left(1 + \frac{dy}{dx}\right) \quad \dots(i)$$

Since, tangent is parallel to $x + 2y = 0$,

$$\text{Then slope } \frac{dy}{dx} = -\frac{1}{2}$$

\therefore From Eq.(i),

$$-\frac{1}{2} = -\sin(x + y) \cdot \left(1 - \frac{1}{2}\right)$$

$$\Rightarrow \sin(x + y) = 1$$

Which shows $\cos(x + y) = 0$

$$\therefore y = 0$$

$$\Rightarrow x + y = \frac{\pi}{2} \text{ or } -\frac{3\pi}{2}$$

$$\therefore x = \frac{\pi}{2} \text{ or } -\frac{3\pi}{2}$$

Thus, required points are

$$\left(\frac{\pi}{2}, 0\right) \text{ and } \left(-\frac{3\pi}{2}, 0\right)$$

\therefore Equation of tangents are

$$\frac{y-0}{x-\pi/2} = -\frac{1}{2} \text{ and } \frac{y-0}{x+3\pi/2} = -\frac{1}{2}$$

$$\Rightarrow 2y = -x + \frac{\pi}{2} \text{ and } 2y = -x - \frac{3\pi}{2}$$

$$\Rightarrow x + 2y = \frac{\pi}{2} \text{ and } x + 2y = -\frac{3\pi}{2}$$

are the required equation of tangents

$$\text{Sol 7: Let } P(a \cos \theta, 2 \sin \theta) \text{ be a point on the ellipse}$$

$$4x^2 + a^2 y^2 = 4a^2 \text{ i.e., } \frac{x^2}{a^2} + \frac{y^2}{4} = 1$$

Let $A(0, -2)$ be the given point. Then,

$$(AP)^2 = a^2 \cos^2 \theta + 4(1 + \sin^2 \theta)$$

$$\Rightarrow \frac{d}{d\theta} (AP)^2 = -a^2 \sin^2 2\theta + 8(1 + \sin \theta) \cdot \cos \theta$$

$$\Rightarrow \frac{d}{d\theta} (AP)^2 = [(8 - 2a^2) \sin \theta + 8] \cos \theta$$

For maximum or minimum, we put $\frac{d}{d\theta} (AP)^2 = 0$

$$\Rightarrow [(8 - 2a^2) \sin \theta + 8] \cos \theta = 0$$

$$\Rightarrow \cos \theta = 0 \text{ or } \sin \theta = \frac{4}{a^2 - 4}$$

$$(\because 4 < a^2 < 8 \Rightarrow \frac{4}{a^2 - 4} > 1 \Rightarrow \sin \theta > 1, \text{ which is impossible})$$

$$\text{Now, } \frac{d^2}{d\theta^2} (AP)^2 = -\{(8 - 2a^2) \sin \theta + 8\} \sin \theta + (8 - 2a^2) \cdot \cos^2 \theta$$

$$\text{For } \theta = \frac{\pi}{2}, \text{ we have}$$

$$\frac{d^2}{d\theta^2} (AP)^2 = -(16 - 2a^2) < 0$$

Thus, AP^2 ie, AP is maximum when $\theta = \frac{\pi}{2}$. The point on the curve $4x^2 + a^2y^2 = 4a^2$ that is farthest from the point $A(0, -2)$ is

$$\left(a \cos \frac{\pi}{2}, 2 \sin \frac{\pi}{2} \right) = (0, 2)$$

Sol 8: Since, equation of normal to $y^2 = 4ax$ is

$$y = mx - 2am - am^3$$

Equation of normal for $y^2 = x$ is

$$y = mx - \frac{m}{2} - \frac{1}{4}m^3 \text{ which passes through } (c, 0)$$

$$\therefore 0 = m \left(c - \frac{1}{2} - \frac{m^2}{4} \right) \Rightarrow m = 0$$

$$\text{and } \frac{m^2}{4} = c - \frac{1}{2} \Rightarrow m = \pm 2 \sqrt{c - \frac{1}{2}}$$

Which gives a normal as x-axis and for other two normals


$$c - \frac{1}{2} > 0 \Rightarrow c > \frac{1}{2}$$

Now, if normals are perpendicular

$$\Rightarrow \left(2\sqrt{c - \frac{1}{2}} \right) \cdot \left(-2\sqrt{c - \frac{1}{2}} \right) = -1$$

$$\Rightarrow c - \frac{1}{2} = \frac{1}{4} \Rightarrow c = \frac{3}{4}$$

Sol 9:



Any point on the parabola $y = x^2$ is of the form (t, t^2) .

$$\text{Now, } \frac{dy}{dx} = 2x \Rightarrow \left[\frac{dy}{dx} \right]_{x=t} = 2t$$

Which is the slope of the tangent. So, the slope of the normal to $y = x^2$ at $A(t, t^2)$ is $-1/2t$.

Therefore, the equation of the normal to $y = x^2$ at $A(t, t^2)$ is

$$Y - t^2 = \left(-\frac{1}{2t} \right) (x - t) \quad \dots (i)$$

Suppose eq. (1) meets the curve again at $B(t_1, t_1^2)$

$$\text{Then, } t_1^2 - t^2 = -\frac{1}{2t} (t_1 - t)$$

$$\Rightarrow (t_1 - t) (t_1 + t) = -\frac{1}{2t} (t_1 - t)$$

$$\Rightarrow (t_1 + t) = -\frac{1}{2t}$$

$$\Rightarrow t_1 = -t - \frac{1}{2t}$$

Therefore, length of chord,

$$L = AB^2 = (t - t_1)^2 + (t^2 - t_1^2)^2$$

$$= (t - t_1)^2 + (t - t_1)^2 (t + t_1)^2$$

$$= (t - t_1)^2 [1 + (t + t_1)^2]$$

$$= \left(t + t + \frac{1}{2t} \right)^2 \left[1 + \left(t - t - \frac{1}{2t} \right)^2 \right]$$

$$\Rightarrow L = \left(2t + \frac{1}{2t} \right)^2 \left(1 + \frac{1}{4t^2} \right) = 4t^2 \left(1 + \frac{1}{4t^2} \right)^3$$

\therefore On differentiating w.r.t., we get

$$\frac{dL}{dt} = 8t \left(1 + \frac{1}{4t^2} \right)^3 + 12t^2 \left(1 + \frac{1}{4t^2} \right)^2 \left(-\frac{2}{4t^3} \right)$$

$$= 2 \left(1 + \frac{1}{4t^2} \right)^2 \left[4t \left(1 + \frac{1}{4t^2} \right) - \frac{3}{t} \right]$$

$$= 2 \left(1 + \frac{1}{4t^2} \right)^2 \left(4t - \frac{2}{t} \right) = 4 \left(1 + \frac{1}{4t^2} \right)^2 \left(2t - \frac{1}{t} \right)$$

For maxima or minima, we must have $\frac{dL}{dt} = 0$

$$\Rightarrow 2t - \frac{1}{t} = 0 \Rightarrow t^2 = \frac{1}{2} \Rightarrow t = \pm \frac{1}{\sqrt{2}}$$

Next,

$$\frac{d^2L}{dt^2} = 8 \left(1 + \frac{1}{4t^2} \right) \left(-\frac{1}{2t^3} \right) \left(2t - \frac{1}{t} \right) + 4 \left(1 + \frac{1}{4t^2} \right)^2 \left(2 + \frac{1}{t^2} \right)$$

$$\Rightarrow \left[\frac{d^2L}{dt^2} \right]_{t=\pm 1/\sqrt{2}} = 0 + 4 \left(1 + \frac{1}{2} \right)^2 (2+2) > 0$$

Therefore, L is minimum, when $t = \pm \frac{1}{\sqrt{2}}$, point A is $\left(\frac{1}{\sqrt{2}}, \frac{1}{2} \right)$ and point B is $(-\sqrt{2}, 2)$ when $t = -\frac{1}{\sqrt{2}}$, A is $\left(-\frac{1}{\sqrt{2}}, \frac{1}{2} \right)$, B is $(+\sqrt{2}, 2)$

Again, when $t = \frac{1}{\sqrt{2}}$, the equation of AB is

$$\frac{y-2}{\frac{1}{2}-2} = \frac{x+\sqrt{2}}{\frac{1}{\sqrt{2}}+\sqrt{2}}$$

$$\Rightarrow (y-2) \left\{ \left(\frac{1}{\sqrt{2}} + \sqrt{2} \right) \right\} = (x + \sqrt{2}) \left(\frac{1}{2} - 2 \right)$$

$$\Rightarrow -2y + 4 = \sqrt{2}x + 2$$

$$\Rightarrow \sqrt{2}x + 2y - 2 = 0$$

And when $t = -\frac{1}{\sqrt{2}}$, the equation of AB is $\frac{y-2}{\frac{1}{2}-2}$

$$= \frac{x - \sqrt{2}}{\left(-\frac{1}{\sqrt{2}} \right) - \sqrt{2}}$$

$$\Rightarrow (y-2) \left(-\frac{1}{\sqrt{2}} - \sqrt{2} \right) = (x - \sqrt{2}) \left(\frac{1}{2} - 2 \right)$$

$$\Rightarrow 2y - 4 = \sqrt{2}(x - \sqrt{2})$$

$$\Rightarrow \sqrt{2}x - 2y + 2 = 0$$

Sol 10: Given, $y = (1+x)^y + \sin^{-1}(\sin^2 x)$

Let $y = u + v$,

where $u = (1+x)^y$, $v = \sin^{-1}(\sin^2 x)$

On differentiating, we get

$$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

Now, $u = (1+x)^y$

Take logarithm on both sides, we get

$$\log_e u = y \log_e(1+x)$$

$$\Rightarrow \frac{1}{u} \frac{du}{dx} = \frac{y}{1+x} + \frac{dy}{dx} \{ \log_e(1+x) \}$$

$$\Rightarrow \frac{du}{dx} = (1+x)^y$$

$$\left[\frac{y}{1+x} + \frac{dy}{dx} \log_e(1+x) \right] \quad \dots (ii)$$

Again, $v = \sin^{-1}(\sin^2 x)$

$$\Rightarrow \sin v = \sin^2 x$$

$$\Rightarrow \cos v \frac{dv}{dx} = 2 \sin x \cos x$$

$$\Rightarrow \frac{dv}{dx} = \frac{1}{\cos v} (2 \sin x \cos x)$$

$$\Rightarrow \frac{dv}{dx} = \frac{2 \sin x \cos x}{\sqrt{1 - \sin^2 v}} = \frac{2 \sin x \cos x}{\sqrt{1 - \sin^4 x}} \quad \dots (iii)$$

\therefore From Eq. (i)

$$\frac{dy}{dx} = (1+x)^y \left[\frac{y}{1+x} + \frac{dy}{dx} \log_e(1+x) \right] + \frac{2 \sin x \cos x}{\sqrt{1 - \sin^4 x}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{y(1+x)^{y-1} + 2 \sin x \cos x / \sqrt{1 - \sin^4 x}}{1 - (1+x)^y \log_e(1+x)}$$

At $x = 0$,

$$y = (1+0)^y + \sin^{-1} \sin(0) = 1$$

$$\therefore \frac{dy}{dx} = \frac{1(1+0)^{1-1} + 2 \sin 0 \cos 0 / \sqrt{1 - \sin^4 0}}{1 - (1+0)^1 \log_e(1+0)}$$

$$\Rightarrow \frac{dy}{dx} = 1$$

Again, the slope of the normal is

$$m = -\frac{1}{dy/dx} = -1$$

Hence, the required equation of the normal is

$$y - 1 = (-1)(x - 0)$$

$$\text{ie, } y + x - 1 = 0$$

Sol 11: Let any point P_1 on $y = x^3$ be (h, h^3)

Then tangent at P_1 is

$$y - h^3 = 3h^2(x - h) \dots (i)$$

It meets $y = x^3$ at P_2 .

On putting the value of y in Eq. (i)

$$x^3 - h^3 = 3h^2(x - h)$$

$$\Rightarrow (x - h)(x^2 + xh + h^2) = 3h^2(x - h)$$

$$\Rightarrow x^2 + xh + h^2 = 3h^2$$

or $x = h$

$$\Rightarrow x^2 + xh + 2h^2 = 0 \Rightarrow (x - h)(x + 2h) = 0$$

$$\Rightarrow x = h \text{ or } x = -2h$$

Therefore, $x = -2h$ is the point P_2 ,

Which implies $y = -8h^3$

Hence, point $P_2 \equiv (-2h, -8h^3)$

Again, tangent at P_2 is

$$y + 8h^3 = 3(-2h)^2(x + 2h)$$

It meets $y = x^3$ at P_3

$$\Rightarrow x^3 + 8h^3 = 12h^2(x + 2h)$$

$$\Rightarrow x^3 - 2hx - 8h^2 = 0$$

$$\Rightarrow (x + 2h)(x - 4h) = 0$$

$$\Rightarrow x = 4h \Rightarrow y = 64h^3$$

Therefore, $P_3 \equiv (4h, 64h^3)$

Similarly, we get $P_4 \equiv (-8h, -8^3 h^3)$

Hence, the abscissae are

$h, -2h, 4h, -8h, \dots$ which form a GP

Let $D' = \Delta P_1 P_2 P_3$ and $D'' = \Delta P_2 P_3 P_4$

$$\frac{D'}{D''} = \frac{\Delta P_1 P_2 P_3}{\Delta P_2 P_3 P_4} = \frac{\frac{1}{2} \begin{vmatrix} h & h^3 & 1 \\ -2h & -8h^3 & 1 \\ 4h & 64h^3 & 1 \end{vmatrix}}{\frac{1}{2} \begin{vmatrix} -2h & -8h^3 & 1 \\ 4h & 64h^3 & 1 \\ -8h & -512h^3 & 1 \end{vmatrix}}$$

$$= \frac{\frac{1}{2} \begin{vmatrix} h & h^3 & 1 \\ -2h & -8h^3 & 1 \\ 4h & 64h^3 & 1 \end{vmatrix}}{\frac{1}{2} \times (-2) \times (-8) \begin{vmatrix} h & h^3 & 1 \\ -2h & -8h^3 & 1 \\ 4h & 64h^3 & 1 \end{vmatrix}}$$

$$= \frac{1}{16}$$

which is the required ratio.

Sol 12: (A, C)

Let the sides of rectangle be $15k$ and $8k$ and side of square be x then $(15k - 2x)(8k - 2x)x$ is volume.

$$v = 2(2x^3 - 23kx^2 + 60k^2x)$$

$$\left. \frac{dv}{dx} \right|_{x=5} = 0$$

$$6x^2 - 46kx + 60k^2 \big|_{x=5} = 0$$

$$6k^2 - 23k + 15 = 0$$

$$k = 3, k = \frac{5}{6}. \text{ Only } k = 3 \text{ is permissible}$$

So, the sides are 45 and 24.

Sol 13: (B)

$$P \rightarrow \frac{\cos(\tan^{-1} y) + y \sin(\tan^{-1} y)}{\cot(\sin^{-1} y) + \tan(\sin^{-1} y)}$$

$$= \frac{\frac{1}{\sqrt{1+y^2}} + \frac{y^2}{\sqrt{1+y^2}}}{\frac{\sqrt{1-y^2}}{y} + \frac{y}{\sqrt{1-y^2}}} = \frac{\sqrt{1+y^2}}{\frac{1}{y\sqrt{1-y^2}}} = y\sqrt{1-y^4}$$

$$\Rightarrow \frac{1}{y^2} \left(\frac{\cos(\tan^{-1} y) + y \sin(\tan^{-1} y)}{\cot(\sin^{-1} y) + \tan(\sin^{-1} y)} \right)^2 + y^4$$

$$= \frac{1}{y^2} (y^2 (1 - y^4)) + y^4 = 1 - y^4 + y^4 = 1$$

$$Q \rightarrow \cos x + \cos y + \cos z = 0$$

$$\sin x + \sin y + \sin z = 0$$

$$\cos x + \cos y = -\cos z \quad \dots (i)$$

$$\sin x + \sin y = -\sin z \quad \dots (ii)$$

$$(1)^2 + (2)^2$$

$$1 + 1 + 2(\cos x \cos y + \sin x \sin y) = 1$$

$$2 + 2 \cos(x - y) = 1$$

$$2 \cos(x - y) = -1$$

$$\cos(x - y) = -\frac{1}{2}$$

$$\Rightarrow 2 \cos^2 \left(\frac{x-y}{2} \right) - 1 = -\frac{1}{2} \Rightarrow 2 \cos^2 \left(\frac{x-y}{2} \right) = \frac{1}{2}$$

$$\Rightarrow \cos^2 \left(\frac{x-y}{2} \right) = \frac{1}{4} \Rightarrow \cos \left(\frac{x-y}{2} \right) = \frac{1}{2}$$

$$R \rightarrow \cos \left(\frac{\pi}{4} - x \right) \cos 2x + \sin x \sin 2x \sec x$$

$$= \cos x \sin 2x \sec x + \cos \left(\frac{\pi}{4} + x \right) \cos 2x$$

$$\left[\cos \left(\frac{\pi}{4} - x \right) - \cos \left(\frac{\pi}{4} + x \right) \right] \cos 2x$$

$$= (\cos x \sin 2x - \sin x \sin 2x) \sec x$$

$$\frac{2}{\sqrt{2}} \sin x \cos 2x = (\cos x - \sin x) \sin 2x \sec x$$

$$\sqrt{2} \sin x \cos 2x = (\cos x - \sin x) 2 \sin x$$

$$\frac{1}{\sqrt{2}} = \frac{1}{\cos x + \sin x} \Rightarrow x = \frac{\pi}{4}$$

$$\sec x = \sec \frac{\pi}{4} = \sqrt{2}$$

$$S \rightarrow \cot \left(\sin^{-1} \sqrt{1-x^2} \right)$$

$$\cot \alpha = \frac{x}{\sqrt{1-x^2}}$$

$$\tan^{-1}(x\sqrt{6}) = \phi$$

$$\sin \phi = \frac{x\sqrt{6}}{\sqrt{6x^2+1}}$$

$$\Rightarrow \frac{x}{\sqrt{1-x^2}} = \frac{x\sqrt{6}}{\sqrt{6x^2+1}}$$

$$6x^2+1=6-6x^2$$

$$12x^2=5$$

$$x = \sqrt{\frac{5}{12}} = \frac{1}{2}\sqrt{\frac{5}{3}}$$